

## RESEARCH ARTICLE

# Risk factors, changes in serum inflammatory factors, and clinical prevention and control measures for puerperal infection

Hongbi Song | Keli Hu  | Xuyuan Du | Jiao Zhang | Shu Zhao

Department of Obstetrics, Guizhou  
Provincial People's Hospital, Guiyang, China

**Correspondence**

Keli Hu, Department of Obstetrics, Guizhou  
Provincial People's Hospital, No.1, Baoshan  
South Road, Guiyang 550002, Guizhou,  
China.  
Email: kelihu6lv@163.com

**Funding information**

Special fund for clinical research of Wu  
Jieping medical fund, Grant/Award Number:  
320.6750.17173

**Abstract**

**Background:** To investigate the risk factors and changes in serum inflammatory factors in puerperal infection, and propose clinical prevention measures.

**Methods:** A total of 240 subjects with suspected puerperal infection treated in our hospital from January 2017 to December 2017 were collected, among which puerperal infection was definitely diagnosed in 40 cases, and it was excluded in 40 cases. Levels of interleukin-6 (IL-6), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), and high-sensitivity C-reactive protein (hs-CRP) were compared between the two groups, and the change trends of IL-6 and hs-CRP were recorded.

**Results:** Levels of IL-6, hs-CRP, and TNF- $\alpha$  in puerperal infection group were higher than those in non-infection group ( $P < .05$ ). Levels of IL-6 and hs-CRP at enrollment and 1-3 days after enrollment in infection group were higher than those in non-infection group ( $P < .05$ ). The body mass index  $>25$ , placenta previa, placenta accreta, postpartum hemorrhage, premature rupture of membrane, gestational diabetes mellitus, and anemia during pregnancy were relevant and independent risk factors for puerperal infection. Puerperal infection occurred in uterine cavity, vagina, pelvic peritoneum, pelvic tissue, incision, urinary system, etc, and gram-negative (G $+$ ) bacteria were dominated in pathogens.

**Conclusion:** The inflammatory response of patients with puerperal infection is significantly enhanced.

**KEYWORDS**

inflammatory factors, prevention measures, puerperal infection, risk factors

## 1 | INTRODUCTION

In clinic, puerperal infection mainly refers to the reproductive tract infection occurring after delivery, which, as a kind of complication seriously threatening delivery quality and life safety of puerperae, has a certain influence on postpartum recovery and even neonatal feeding.<sup>1</sup> Birth canal injury is caused due to fetal delivery via reproductive tract during the puerperium, and the body's immunity of pregnant women

significantly declines during the puerperium. As a result, pathogenic microorganisms invade the human body, leading to infection, and even septicopyemia and threatening the maternal life.<sup>2</sup> At the same time, the drug resistance of pathogenic bacteria of puerperal infection has significantly increased nowadays, and the incidence rate of puerperal infection caused by uncommon pathogenic bacteria in the past also increases, so the traditional antibacterial drugs, such as penicillin, fail to control the infection effectively in clinical experiential medication.<sup>3</sup>

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2019 The Authors. *Journal of Clinical Laboratory Analysis* Published by Wiley Periodicals, Inc.

Meanwhile, levels of inflammation-related cytokines in patients with puerperal infection are also significantly increased. Interleukin-6 (IL-6), as the most important inflammation-related cytokine in the body, mainly regulates the cell function and is involved in the body immunity.<sup>4</sup> High-sensitivity C-reactive protein (hs-CRP), as the most widely used inflammation-related cytokine currently, has a definite correlation with the severity of tissue damage and infection, which is an important index in clinical observation in acute infection stage.<sup>5</sup>

In this study, in order to better investigate the occurrence of puerperal infection, the analysis of common pathogenic factors of puerperal infection started with changes in the body's common inflammatory factors after puerperal infection, and common infection sites and pathogen distribution were explored, hoping to provide corresponding guidance for clinical diagnosis, treatment, and prevention.

## 2 | DATA AND METHODS

### 2.1 | General data

A total of 240 subjects with suspected puerperal infection treated in our hospital from January 2017 to December 2017 were collected. Another 40 subjects without puerperal infection during the same period were selected as control group. Before enrollment, all subjects signed and agreed to be enrolled, and this survey research was approved by the Ethics Committee of hospital. Inclusion criteria: primipara who displayed signs of infection in any part of the body within 48 hours before enrollment, underwent vaginal delivery, and had relevant medical data. Subject who was associated with pain abdomen, malodorous lochia, abdominal distention, uterine tenderness, pelvic abscess, peritonitis, mechanical or foreign body injury. Exclusion criteria: subjects who took antibacterial drugs or immunosuppressors within 48 hours before enrollment, or complicated with malignant tumor, severe dysfunction in the heart, lung, liver, kidney, mental diseases, fever due to medical causes, wound/surgical site infection, mastitis, urinary tract infection or thrombophlebitis.

### 2.2 | Investigation methods

Clinical relevant data of subjects with puerperal infection were analyzed, and all clinical data were collected by obstetricians and related investigators receiving unified training. Patients and their authorizers should be fully cooperated in the collection of investigation data, the investigation was conducted anonymously, the patients' privacy should be protected during the process, and the question raised by patients was answered. The relevant investigation results obtained should not be disclosed to any organization or individual without the permission of subjects enrolled and their authorizers. At enrollment and 30 days after the first investigation, investigators filled out the same questionnaire and completed the data of subjects, and the correlation coefficients obtained in the two times were set as the stability coefficients. In this study, the reliability coefficient ( $\alpha = 0.1$ ) was used to evaluate

the reliability level, and it was 0.921 after calculation. Clinical data obtained were checked alternatively by two people and entered into the EpiData software data analysis system, followed by statistical processing via Statistical Product and Service Solutions (SPSS) 21.0.

### 2.3 | Detection of inflammatory factor levels and standard reference values

IL-6 was detected via enzyme-linked immunosorbent assay, whose standard reference value is 0.37-0.46 ng/L. TNF- $\alpha$  was detected via double-antibody single-step sandwich method, whose standard reference value is 5-100 ng/L, and hs-CRP was detected via latex-enhanced immunoturbidimetry, whose standard reference value is  $\leq 10$  mg/L.

### 2.4 | Identification of pathogen species of puerperal infection

Pathogen culture and identification were based on the *National Clinical Laboratory Procedures*, and all operations were performed in strict accordance with the above operating procedures. Bacteria were cultured using the Kirby-Bauer (K-B) diffusion method, while fungi were cultured using the glucose peptone agar medium. All operations were conducted strictly according to the CLSI 2008-2010 procedures, and data were analyzed using WHONET (version 5.3-5.4). All the above operations were performed by the medical laboratory technicians with working experience in clinical detection for 5 years or above strictly according to the operation procedures and regulations.

### 2.5 | Observation indexes

Levels of inflammatory cytokines (IL-6, TNF- $\alpha$  and hs-CRP) in subjects with and without puerperal infection were compared, and the change trends of IL-6 and hs-CRP in both groups were recorded at enrollment and 1-3 days after enrollment. Univariate and multivariate analyses were performed for relevant clinical data, such as age, body mass index, gestational week, placenta previa, placenta accreta, postpartum hemorrhage, premature rupture of membrane, gestational hypertension, gestational diabetes mellitus, and anemia during pregnancy, and the infection sites and pathogen distribution in subjects with puerperal infection were recorded.

### 2.6 | Statistical analysis

SPSS 21.0 (IBM) statistical software was used. Univariate analysis was performed first for puerperal infection, followed by multivariate logistic regression analysis. Measurement data were presented as mean  $\pm$  standard deviation ( $\bar{x} \pm s$ ), and chi-square test was used for the intergroup comparison of rate.  $P < .05$  suggested that the difference was statistically significant.

### 3 | RESULTS

#### 3.1 | Comparisons of inflammatory factor levels between the two groups

A total of 240 subjects with suspected puerperal infection treated in our hospital from January 2017 to December 2017 were collected and 40 women with severe maternal sepsis were finally enrolled for


**TABLE 1** Comparisons of inflammatory factor levels between the two groups ( $\bar{x} \pm s$ )

	IL-6 (ng/L)	hs-CRP (mg/L)	TNF- $\alpha$ (ng/L)
Infection group	1.05 $\pm$ 0.15	16.5 $\pm$ 1.0	130.0 $\pm$ 7.4
Non-infection group	0.41 $\pm$ 0.03	4.1 $\pm$ 0.1	71.3 $\pm$ 4.0
t	26.461	78.035	44.134
P	.000	.000	.000

the study according to the inclusion and exclusion criteria and result of pathogen culture. They were aged 19-42 years old ([28.5  $\pm$  1.2] years old on average) with the gestational week of 34-41 weeks ([38.1  $\pm$  0.6] weeks on average), including 21 cases receiving lateral episiotomy. Another 40 subjects in control group were aged 19-42 years old ([28.6  $\pm$  1.1] years old on average) with the gestational week of 34-41 weeks ([38.0  $\pm$  0.6] weeks on average), including 20 cases receiving lateral episiotomy. Levels of inflammatory factors (IL-6, hs-CRP and TNF- $\alpha$ ) were detected in both groups and the levels in puerperal infection group were significantly higher than those in non-infection group ( $P < .05$ ; Table 1).

#### 3.2 | Change trends of main inflammatory factor (IL-6) level in the two groups

At enrollment and 1-3 days after enrollment, the IL-6 level was (1.05  $\pm$  0.15) ng/L, (0.89  $\pm$  0.06) ng/L, (0.63  $\pm$  0.05) ng/L, and (0.51  $\pm$  0.03) ng/L, respectively, in infection group, and (0.41  $\pm$  0.03)



**FIGURE 1** Levels of inflammatory factors and infection sites. Change trends of main inflammatory factors IL-6 (A) and hs-CRP (B) levels in the two groups by ELISA. C, Main infection sites in patients with puerperal infection include uterine cavity, vagina, pelvic peritoneum, pelvic tissue, incision, and urinary system

ng/L, (0.39 ± 0.03) ng/L, (0.28 ± 0.02) ng/L, and (0.26 ± 0.02) ng/L, respectively, in non-infection group. Levels of IL-6 in infection group were evidently higher than those in non-infection group at different time points ( $t = 26.461, 47.140, 41.105, \text{ and } 43.853, P < .05$ ; Figure 1A).

### 3.3 | Change trends of main inflammatory factor (hs-CRP) level in the two groups

At enrollment and 1-3 days after enrollment, the hs-CRP level was (16.5 ± 1.0) mg/L, (12.5 ± 0.8) mg/L, (11.7 ± 0.6) mg/L, and (9.9 ± 0.3) mg/L, respectively, in infection group, and (4.1 ± 0.1) mg/L, (4.0 ± 0.1) mg/L, (3.8 ± 0.1) mg/L, and (4.0 ± 0.1) mg/L, respectively, in non-infection group. Levels of hs-CRP in infection group were obviously higher than those in non-infection group at different time points ( $t = 78.035, 66.679, 82.140, \text{ and } 118.0000, P < .05$ ; Figure 1B).

### 3.4 | Univariate analysis of relevant risk factors in puerperal infection group

According to the univariate analysis, the incidence rate of puerperal infection was significantly increased in subjects with the body mass index >25, placenta previa, placenta accreta, postpartum hemorrhage, premature rupture of membrane, gestational diabetes mellitus, and anemia during pregnancy ( $P < .05$ ), and they were relevant risk factors for puerperal infection (Table 2).

### 3.5 | Multivariate Logistic regression analysis of relevant risk factors in puerperal infection group

According to the multivariate logistic regression analysis, the body mass index >25, placenta previa, placenta accreta, postpartum hemorrhage, premature rupture of membrane, gestational diabetes mellitus, and anemia during pregnancy were independent risk factors for puerperal infection (Table 3).

### 3.6 | Analysis of main infection sites

Among 40 cases of puerperal infection, there were 14 cases (35.0%) of uterus cavity infection, 8 cases (20.0%) of vaginal infection, 7 cases (17.5%) of pelvic peritoneal infection, 4 cases (10.0%) of pelvic tissue infection, 3 cases (7.5%) of incision infection, 2 cases (5.0%) of urinary system infection, and 2 cases (5.0%) of infection in other sites (Figure 1C).

### 3.7 | Pathogen distribution in subjects with puerperal infection

Among 40 cases of puerperal infection, G<sup>-</sup> bacteria were detected in 24 cases (60.0%), G<sup>+</sup> bacteria in 14 cases (35.0%), and fungi in

**TABLE 2** Univariate analysis of relevant risk factors in puerperal infection group (n)

	Case surveyed	Case of puerperal infection	$\chi^2$	P
Age				
>30 y old	115	19	0.003	.954
<30 y old	125	21		
Body mass index				
>25	135	31	8.808	.003
<25	105	9		
Gestational week				
>38 wk	130	21	0.054	.817
<38 wk	110	19		
Placenta previa				
Yes	116	35	29.486	.000
No	124	5		
Placenta accreta				
Yes	112	34	31.228	.000
No	128	4		
Postpartum hemorrhage				
Yes	125	30	10.101	.001
No	115	10		
Premature rupture of membrane				
Yes	130	29	6.498	.011
No	110	11		
Gestational hypertension				
Yes	110	19	0.054	.817
No	130	21		
Gestational diabetes mellitus				
Yes	112	32	18.552	.000
No	128	8		
Anemia during pregnancy				
Yes	123	30	10.837	.001
No	117	10		

two cases (5.0%). Specifically, *Staphylococcus aureus* accounts for 12.5%, *Staphylococcus epidermidis* 2.5%, *Enterococcus* 15.0%, *Streptococcus* 5.0%, *Gardnerella vaginalis* 12.5%, *Escherichia coli* 27.5%, *Pseudomonas aeruginosa* 10.0%, *Acinetobacter baumannii* 5.0%, *Klebsiella pneumoniae* 5.0%, *Fungi* 5.0%, *Candida albicans* 2.5%, and *Candida tropicalis* 2.5%.

## 4 | DISCUSSION

Puerperal infection is a common and frequently occurring disease in clinic,<sup>6</sup> which is also a common postpartum complication in puerperae. Its clinical manifestations mainly include elevated postpartum

**TABLE 3** Multivariate Logistic regression analysis of relevant risk factors in puerperal infection group

	$\beta$	Standard error (SE)	W	OR	P	95% CI
Body mass index	2.34	0.46	9.92	12.81	.01	1.21-34.40
Placenta previa	2.92	0.54	4.46	6.86	.04	1.59-49.24
Placenta accreta	1.82	0.50	10.82	4.81	.01	1.58-14.38
Postpartum hemorrhage	1.83	0.57	4.82	1.96	.04	1.03-7.45
Premature rupture of membrane	0.89	0.53	9.47	2.44	.00	1.38-4.31
Gestational diabetes mellitus	1.12	15.50	6.38	3.07	.00	1.76-5.34
Anemia during pregnancy	1.71	33.60	7.11	5.52	.00	3.10-9.83

temperature accompanied by chills, lower abdominal pain, and tenderness, and severe infection displayed in biochemical examination.<sup>7</sup> During the delivery process, changes in the physiological structure of pregnant women, hemorrhage, delivery injury, etc. lead to the decline in the body's immunity, resulting in puerperal infection under the action of some risk factors.<sup>8</sup> Subjects with mild inflammatory response mostly suffer from the local infection in the reproductive system, while those with severe inflammatory response may have sepsis and even septicopyemia, threatening the life.<sup>9</sup> Therefore, effective measures should be actively taken in clinic to ensure the clinical therapeutic effect once puerperal infection occurs, thereby improving the prognosis of pregnant women, and minimizing the impact of puerperal infection on puerperae and neonates.<sup>10</sup>

In this study, in order to better investigate the occurrence, development, and prevention of puerperal infection, levels of inflammatory factors were compared between subjects with and without puerperal infection. It was found that levels of inflammatory factors (IL-6, hs-CRP, and TNF- $\alpha$ ) in puerperal infection group were significantly higher than those in non-infection group, and levels of IL-6 and hs-CRP at enrollment and 1-3 days after enrollment in infection group were also obviously higher than those in non-infection group, indicating that levels of inflammatory cytokines in the body of subjects with puerperal infection are remarkably increased, and the body's inflammatory response is significant. At the same time, analyses of risk factors for puerperal infection showed that the body mass index >25, placenta previa, placenta accreta, postpartum hemorrhage, premature rupture of membrane, gestational diabetes mellitus, and anemia during pregnancy were relevant and independent risk factors for puerperal infection, suggesting that subjects with the body mass index >25, placenta previa, placenta accreta, postpartum hemorrhage, premature rupture of membrane, gestational diabetes mellitus, and anemia during pregnancy should be paid attention to in clinic, and related measures should be taken to prevent puerperal infection. Finally, the study on puerperal infection sites and pathogenic bacteria manifested that the infection mainly occurred in reproductive tract, such as uterine cavity and vagina, followed by pelvic cavity. Gram-negative bacteria were dominated in

the pathogenic bacteria, followed by gram-positive bacteria (about 30%). In terms of antibacterial drugs selected in experiential therapy, therefore, it is recommended that gram-negative bacteria be the main target with consideration to gram-positive bacteria.

Prevention measures of puerperal infection should be taken before pregnancy and during pregnancy, delivery, and puerperium. First, before pregnancy, it is suggested that women of child-bearing age strengthen physical exercise, pay attention to nutritional regulation, actively prevent and control the reproductive system diseases, especially inflammatory diseases, reduce the frequency of uterine curettage, and make good preparation for pregnancy.<sup>11</sup> During pregnancy, it is suggested that early pregnancy be determined in time, health management during pregnancy be actively implemented,<sup>12</sup> the standard maternal manual be created before 13 weeks of gestation, the comprehensive physical examination be given,<sup>13</sup> and high-risk pregnancy and pathologic pregnancy be screened to improve the safety of pregnancy. During delivery, sterile operations should be strictly performed, the digital anal examination is strictly forbidden,<sup>14</sup> the frequency of vaginal examination should be reduced as far as possible to reduce the incidence rate of retrograde infection,<sup>15</sup> indications of cesarean section and lateral episiotomy should be strictly grasped, antibacterial drugs can be used for a short time for those undergoing cesarean section to prevent the infection, and the maternal management should be strengthened<sup>16</sup> to improve the hospital delivery rate,<sup>17</sup> prevent postpartum hemorrhage,<sup>18</sup> and reduce the injury of birth canal due to delivery as far as possible.<sup>19</sup> During puerperium, it is recommended that puerperae get enough sleep, strengthen nutrition reasonably, and improve the body's immunity, the health management be strengthened, and prevention measures be actively taken for puerperae with high-risk factors, thereby improving the prognosis of patients and reducing the incidence of puerperal infection.<sup>20</sup> The limitation in our study still exists that a relatively larger study population from different regions requires to be included to confirm our finding and even reflect data at national level, which may provide fundamental leads for the diagnosis and treatment of puerperal infection.

## 5 | CONCLUSION

In conclusion, the inflammatory response of patients with puerperal infection is significantly enhanced, with elevation of IL-6 and hs-CRP levels. Subjects with the body mass index >25, placenta previa, placenta accreta, postpartum hemorrhage, premature rupture of membrane, gestational diabetes mellitus, and anemia during pregnancy should be paid attention to, and prevention measures for puerperal infection should be actively taken. In experiential medication, it is recommended that gram-negative bacteria be the main target with consideration to gram-positive bacteria.

## ACKNOWLEDGMENTS

This work was supported by special fund for clinical research of Wu Jieping medical fund (320.6750.17173).

## ORCID

Keli Hu  <https://orcid.org/0000-0002-5873-9663>

## REFERENCES

- Nenke MA, Zeng A, Meyer EJ, et al. Differential effects of estrogen on corticosteroid-binding globulin forms suggests reduced cleavage in pregnancy. *J Endocr Soc*. 2017;1:202-210.
- Bonet M, Ota E, Chibueze CE, Oladapo OT. Routine antibiotic prophylaxis after normal vaginal birth for reducing maternal infectious morbidity. *Cochrane Database Syst Rev*. 2017;13:21-37.
- Mascarello KC, Horta BL, Silveira MF. Maternal complications and cesarean section without indication: systematic review and meta-analysis. *Rev Saude Publica*. 2017;51:116-118.
- Gudu W, Abdulahi M. Labor, delivery and postpartum complications in nulliparous women with female genital mutilation admitted to karamara hospital. *Ethiop Med J*. 2017;55:11-17.
- Bonet M, Ota E, Chibueze CE, Oladapo OT. Antibiotic prophylaxis for episiotomy repair following vaginal birth. *Cochrane Database Syst Rev*. 2017;2:11-18.
- Tardieu SC, Schmidt E. Group A streptococcus septic shock after surgical abortion: a case report and review of the literature. *Case Rep Obstet Gynecol*. 2017;20:63-67.
- Orazulike NC, Alegbeleye JO, Obiorah CC, Nyengidiki TK, Uzoigwe SA. A 3-year retrospective review of mortality in women of reproductive age in a tertiary health facility in Port Harcourt, Nigeria. *Int J Womens Health*. 2017;16:769-775.
- Wu X, Wang C, Li Y, et al. Cervical dilation balloon combined with intravenous drip of oxytocin for induction of term labor: a multi-center clinical trial. *Arch Gynecol Obstet*. 2018;297:77-83.
- Rosenbloom JI, Stout MJ, Tuuli MG, et al. New labor management guidelines and changes in cesarean delivery patterns. *Am J Obstet Gynecol*. 2017;217:689.
- Gon G, Ali SM, Towriss C, et al. Unpacking the enabling factors for hand, cord and birth-surface hygiene in Zanzibar maternity units. *Health Policy Plan*. 2017;32:1220-1228.
- Easter SR, Molina RL, Venkatesh KK, Kaimal A, Tuomala R, Riley LE. Clinical risk factors associated with peripartum maternal bacteremia. *Obstet Gynecol*. 2017;130:710-717.
- Cobo F, Rodríguez-Granger J, Sampedro A, Navarro-Mari JM. Infected breast cyst due to *Prevotella buccae* resistant to metronidazole. *Anaerobe*. 2017;48:177-178.
- Liabsuetrakul T, Choobun T, Peeyananjarassri K, Islam QM. Antibiotic prophylaxis for operative vaginal delivery. *Cochrane Database Syst Rev*. 2017;5:8-12.
- Jena P, Sheela CN, Venkatachala RP, Devarbhavi H. Obstetric outcome in women with chronic liver disease. *J Obstet Gynaecol India*. 2017;67:263-269.
- Marwah S, Topden SR, Sharma M, Mohindra R, Mittal P. Severe puerperal sepsis-a simmering menace. *J Clin Diagn Res*. 2017;11:QC04-QC08.
- Kulkarni GB, Mirza AM, Ramakrishnan S, Mustare V. Preliminary data on utility of subcutaneous unfractionated heparin in patients with deep cerebral venous thrombosis. *J Thromb Thrombolysis*. 2017;44:247-253.
- Cobo F, Rodríguez-Granger J, Sampedro A, Navarro-Mari JM. Breast abscess due to *Finegoldia magna* in a non-puerperal women. *Anaerobe*. 2017;47:183-184.
- Kobayashi N, Ahmed S, Sumi A, Urushibara N, Kawaguchiya M, Aung MS. Collaborative research on puerperal infections in Bangladesh. *Nihon Eiseigaku Zasshi*. 2017;72:106-111.
- Subramaniam A, Owen J, Campbell SB, Harper LM, Fitzwater JL, Edwards RK. Maternal body mass index and oxytocin exposure in nulliparous women: is there an interaction associated with maternal and neonatal morbidities? *J Matern Fetal Neonatal Med*. 2017;6:1-6.
- Ehrmann Feldman D, Vinet É, Sylvestre MP, et al. Postpartum complications in new mothers with juvenile idiopathic arthritis: a population-based cohort study. *Rheumatology (Oxford)*. 2017;56:1378-1385.

**How to cite this article:** Song H, Hu K, Du X, Zhang J, Zhao S. Risk factors, changes in serum inflammatory factors, and clinical prevention and control measures for puerperal infection. *J Clin Lab Anal*. 2020;34:e23047. <https://doi.org/10.1002/jcla.23047>