



Cochrane
Library

Cochrane Database of Systematic Reviews

Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews (Review)

Anderson L, Taylor RS

Anderson L, Taylor RS.

Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews.

Cochrane Database of Systematic Reviews 2014, Issue 12. Art. No.: CD011273.

DOI: [10.1002/14651858.CD011273.pub2](https://doi.org/10.1002/14651858.CD011273.pub2).

www.cochranelibrary.com

Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews (Review)

Copyright © 2014 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	3
OBJECTIVES	4
METHODS	4
RESULTS	6
Figure 1.	7
DISCUSSION	10
AUTHORS' CONCLUSIONS	11
ACKNOWLEDGEMENTS	12
REFERENCES	13
ADDITIONAL TABLES	17
APPENDICES	39
WHAT'S NEW	40
HISTORY	40
CONTRIBUTIONS OF AUTHORS	40
DECLARATIONS OF INTEREST	40
SOURCES OF SUPPORT	41
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	41
INDEX TERMS	41

[Overview of Reviews]

Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews

Lindsey Anderson¹, Rod S Taylor¹

¹Institute of Health Research, University of Exeter Medical School, Exeter, UK

Contact: Rod S Taylor, rod.taylor@gl.a.ac.uk.

Editorial group: Cochrane Heart Group.

Publication status and date: Edited (no change to conclusions), published in Issue 12, 2021.

Citation: Anderson L, Taylor RS. Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews. *Cochrane Database of Systematic Reviews* 2014, Issue 12. Art. No.: CD011273. DOI: [10.1002/14651858.CD011273.pub2](https://doi.org/10.1002/14651858.CD011273.pub2).

Copyright © 2014 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Overviews are a new approach to summarising evidence and synthesising results from related systematic reviews.

Objectives

To conduct an overview of Cochrane systematic reviews to provide a contemporary review of the evidence for delivery of cardiac rehabilitation, to identify opportunities for merging or splitting existing Cochrane reviews, and to identify current evidence gaps to inform new cardiac rehabilitation systematic review titles.

Methods

We searched The Cochrane Database of Systematic Reviews (2014, Issue 10) to identify systematic reviews that addressed the objectives of this overview. We assessed the quality of included reviews using the Revised Assessment of Multiple Systematic Reviews (R-AMSTAR) measurement tool and the quality of the evidence for reported outcomes using the GRADE framework. The focus of the data presentation was descriptive with detailed tabular presentations of review level and trial level characteristics and results.

Main results

We found six Cochrane systematic reviews and judged them to be of high methodological quality. They included 148 randomised controlled trials (RCTs) in 98,093 participants. Compared with usual care alone, the addition of exercise-based cardiac rehabilitation in low-risk people after myocardial infarction or percutaneous coronary intervention or with heart failure appeared to have no impact on mortality, but did reduce hospital admissions and improved health-related quality of life. Psychological- and education-based interventions alone appeared to have little or no impact on mortality or morbidity but may have improved health-related quality of life. Home- and centre-based programmes were equally effective in improving quality of life outcomes at similar healthcare costs. Selected interventions can increase the uptake of cardiac rehabilitation programmes whilst there is currently only weak evidence to support interventions that improve adherence to cardiac rehabilitation programmes. The quality of the primary RCTs in the included systematic reviews was variable, and limitations in the methodological quality of the RCTs led to downgrading of the quality of the evidence, which varied widely by review and by outcome.

Authors' conclusions

Exercise-based cardiac rehabilitation is an effective and safe therapy to be used in the management of clinically stable people following myocardial infarction or percutaneous coronary intervention or who have heart failure. Future RCTs of cardiac rehabilitation need to improve their reporting methods and reflect the real world practice better including the recruitment of higher risk people and consideration of contemporary models of cardiac rehabilitation delivery, and identify effective interventions for enhancing adherence to rehabilitation.

PLAIN LANGUAGE SUMMARY

Participation in rehabilitation programmes that include regular exercise, can improve the quality of life for people with heart disease

Background

Cardiac rehabilitation (CR) seeks to improve the function, health-related quality of life and well-being of people with heart disease through a combination of activities, in particular exercise training alongside educational and psychological support. Since the mid-2000s, the number of published Cochrane reviews has grown to six systematic reviews/meta-analyses of CR. These reviews assessed the impact of CR on different types of heart disease (e.g. following a heart attack, heart surgery or heart failure) or different ways of providing CR (e.g. in a hospital- or home-based setting, exercise only programmes or exercise in combination with an educational or psychological intervention or both). The aim of the overview was to review the current CR Cochrane reviews to provide a 'friendly front end' to this 'portfolio' of reviews.

Study characteristics

We searched for Cochrane reviews that analysed the data from randomised controlled trials (RCT; experiments that randomly allocate participants to one of two or more treatment groups), which looked at the effectiveness of CR in adults with heart disease and compared patient outcomes with a no-exercise control group. This overview summarised the findings from these reviews.

Key results

We found six high-quality Cochrane reviews that included 148 RCTs in 98,093 people who primarily had experienced a heart attack, had undergone cardiac surgery or had chronic heart failure. The findings of this overview showed important benefits of CR participation that included a reduction in the risk of hospital admissions, as well as improvements in health-related quality of life compared with not undertaking rehabilitation.

Quality of the evidence

The quality of the RCTs in the included systematic reviews was variable, and limitations in their methodological quality led to downgrading of the quality of the evidence, which varied widely by review and outcome. We make the following recommendations for the future conduct and reporting of systematic reviews of CR.

- The scope of CR reviews needs to reflect current guidelines that recommend that CR should be based on an individually prescribed programme of exercise training with appropriate co-interventions.
- Future CR reviews need to explore the complexity of CR using appropriate approaches to explore the association between intervention characteristics and outcomes across trials.
- Future Cochrane CR reviews need to standardise their methods and reporting.

BACKGROUND

Description of the condition

Heart disease is a broad term used to describe a range of diseases that affect the heart, including diseases of heart blood vessels (coronary artery disease), heart rhythm problems (arrhythmias), heart infections and congenital heart defects. Coronary heart disease (CHD) is the most common type of heart disease and its common symptoms are chest pain (angina) and myocardial infarction (MI). Acute coronary syndrome refers to a range of acute CHD states and includes unstable angina (chest pain at rest), non-ST segment elevation MI (ST segment elevation generally absent) and ST segment elevation infarction (persistent ST segment elevation usually present). CHD can result in difficulties in functionality and performance of everyday activities and can impair sexual function (Racca 2010), contributing to a reduction in health-related quality of life (HRQoL) (Gravely-Witte 2007).

CHD is now considered the leading cause of global mortality. According to the World Health Organization (WHO), CHD accounted for 12.9% of all deaths (seven million deaths) and 5.8% of total disability-adjusted life years globally in 2011 (WHO 2014). The situation is worse in high-income countries, and it has been estimated that CHD accounted for 24.8% of all deaths in Europe in 2011 (WHO 2014). However, despite the overall increase in CHD burden in high-income countries, age-adjusted mortality for this disease is declining and over half of people diagnosed now survive (Allender 2008). This is driven largely by preventive interventions, treatments to prevent death during an acute disease manifestation and rehabilitation interventions that prolong survival (Gaziano 2010). Conversely, morbidity is rising, with an increasing number of survivors of MI (Mathers 2008), and an associated number of cases of chronic heart failure (HF) (Kostis 1997).

The most common cause of HF is CHD. Non-ischaemic causes of HF include hypertension and atrial fibrillation. HF is a complex clinical syndrome that results from any structural or functional impairment of ventricular filling or ejection of blood. It has been increasingly recognised that HF has two sub-categories: 1. impaired left ventricular contraction, which results in a reduced ejection fraction (less than 35% to 50%), known as HF with reduced ejection fraction (HFREF) or 'systolic HF'; and 2. HF with preserved ejection fraction (HFPEF) with an ejection fraction of greater than 35% to 50% and also known as 'diastolic HF'. People with HF experience marked reductions in their exercise capacity, which has detrimental effects on their activities of daily living, HRQoL, and their hospital admission rate and mortality (Go 2014). In high-income countries, about 2% of adults have HF, but in people over the age of 65 years, this increases to 6% to 10% (McMurray 2005; Dickstein 2008). The prevalence and incidence of HF is steadily increasing, with approximately 825,000 new cases annually in the US (Go 2014). HF has a poor prognosis, with 30% to 40% of people diagnosed dying within one year, although thereafter the mortality is less than 10% per year (Cowie 2000; Hobbs 2007). However, as with CHD, survival after HF diagnosis has also improved (Go 2014), and in the UK there is evidence of a trend of improved prognosis, with the six-month mortality rate decreasing from 26% in 1995 to 14% in 2005 (Mehta 2009).

Description of the interventions

Many definitions of cardiac rehabilitation (CR) have been proposed. The following definition encompasses the key concepts of CR: "The coordinated sum of activities required to influence favourably the underlying cause of cardiovascular disease, as well as to provide the best possible physical, mental and social conditions, so that the patients may, by their own efforts, preserve or resume optimal functioning in their community and through improved health behaviour, slow or reverse progression of disease" (BACPR 2012). While exercise training is the foundation of CR, it is recommended that 'comprehensive' programmes also include education (e.g. provision of information about a healthy lifestyle) and psychological intervention (e.g. counselling to reduce stress). CR has many of the characteristics of a 'complex intervention' as defined by in the Medical Research Council 2008 guidance for developing and evaluating complex interventions, that is, 1. number of interacting components, 2. number and difficulty of behaviours required by people delivering or receiving the intervention, 3. number and variability of outcomes and 4. degree of flexibility or tailoring of the intervention permitted (non-standardisation/reproducibility) (Craig 2008).

Patient education is the process by which health professionals impart information to patients that will alter their health behaviours or improve their health status (Koongstvedt 2001). There is substantial variation in the delivery of patient education for cardiac patients; it may be classroom- or home-based, group or individual, tailored or generic. Duration and reinforcement of education also differs between programmes. Some programmes are developed according to validated educational theory and by trained professionals while others are delivered by peers.

Interventions that specifically aim to influence psychological or psychosocial outcomes are varied and may range from organisational efforts to improve patient communication and support (e.g. Jolly 1998), to empirically supported psychotherapies used to target diagnosed psychopathology in cardiac patients (e.g. Black 1998). Furthermore, psychological/psychosocial interventions may incorporate other elements of CR such as diet and lifestyle advice, or exercise. In some cases, the intervention may be described as 'psychological' only to the extent that psychological techniques are used to further other treatment goals.

The patient groups routinely recommended for CR include people with post-MI, post-revascularisation procedure and HF. Traditionally, CR programmes have been offered in a supervised centre-based setting. However, many people do not receive rehabilitation (Bethell 2008), and with uptake of CR for both CHD and HF currently at sub-optimal levels (Tierney 2011; Dalal 2012; NICE 2013), home-based CR programmes have been increasingly introduced to widen access and participation. In addition to uptake, maintaining longer-term adherence to CR is also a key challenge (Daly 2002; Moore 2003), and therefore, interventions aimed at improving patient uptake and adherence to CR programmes have been adopted and will be investigated in this overview.

Based on current evidence, national and international guidelines on the management of CHD and HF including those by the American College of Cardiology (ACC)/American Heart Association (AHA), European Society of Cardiology (ESC) and National Institute for Health and Care Excellence (NICE, UK), consistently recommend CR

as an effective and safe intervention (McMurray 2012; NICE 2013; Yancy 2013).

How the intervention might work

The mechanism by which CR may work depends on the patient group and the component of rehabilitation being considered. Most mechanistic evidence is for exercise training.

For people with CHD, exercise training has direct benefits on the heart and coronary vasculature, including myocardial oxygen demand, endothelial function, autonomic tone, coagulation and clotting factors, inflammatory markers and the development of coronary collateral vessels (Clausen 1976; Hambrecht 2000). However, findings of the original Cochrane review of exercise-based CR for CHD (Jolliffe 2001) supported the hypothesis that reductions in mortality may also be mediated via the indirect effects of exercise through improvements in the risk factors for atherosclerotic disease (i.e. lipids, smoking and blood pressure) (Taylor 2006).

The precise mechanism(s) through which exercise training benefits people with HF remains unclear. One explanation, applicable to people with ischaemic causes of HF, is that exercise training improves myocardial perfusion by alleviating endothelial dysfunction therefore dilating coronary vessels and by stimulating new vessel formation by way of intermittent ischaemia (Piepoli 2004). Indeed, Belardinelli and colleagues have demonstrated that aerobic training improves myocardial contractility and diastolic filling (Belardinelli 1998). One meta-analysis by Haykowsky et al, demonstrated the benefits of exercise training on cardiac remodelling as measured by ejection fraction, end-diastolic volume and end-systolic volume (Haykowsky 2007). Regardless of cause, there are important neurohormonal and musculoskeletal abnormalities in HF. Exercise training may reduce adrenergic tone and increase vagal tone, as suggested by an assessment of variability in heart rate. Skeletal muscle dysfunction and wasting may also respond to exercise training (Piepoli 2004). Hambrecht et al. have demonstrated that regular physical activity in people with HF stimulates vasodilation in the skeletal muscle vasculature (Hambrecht 1998).

The benefits of education and psychological interventions depend on changing people's behaviour including improvements in healthy lifestyle and changes in mood, such as reductions in depression and anxiety.

Why it is important to do this overview

In 2001, Jolliffe et al. published the first Cochrane review of CR, summarising the evidence of 32 randomised controlled trials (RCTs) in 8440 post-MI and revascularisation patients, and confirming a mortality benefit of exercise-based CR (Jolliffe 2001). With the funding support of the National Institute of Health Research (NIHR, UK), since the mid-2000s, the number of published Cochrane reviews has grown to six systematic reviews/meta-analyses.

- Exercise-based cardiac rehabilitation for coronary heart disease (Heran 2011).
- Exercise-based rehabilitation for heart failure (Taylor 2014b).
- Psychological interventions for coronary heart disease (Whalley 2011).

- Patient education in the management of coronary heart disease (Brown 2011).
- Home-based versus centre-based cardiac rehabilitation (Taylor 2014a).
- Promoting patient uptake and adherence in cardiac rehabilitation (Karmali 2014).

The development of the portfolio of Cochrane reviews has reflected many of the key areas of evolution in the provision of CR and how this model of service delivery can differ across international healthcare jurisdictions. These include the shift from emphasis on exercise therapy alone to comprehensive secondary prevention including risk factor and dietary education and management of psychological factors; the expansion of the population of cardiac patients receiving CR services to include HF; the development of alternative settings of CR delivery that include home provision in addition to the traditional supervised hospital- or centre-based programmes; and the need to broaden the consideration of the outcomes of CR to inform the needs of healthcare policy makers (e.g. impacts on hospital admission, HRQoL and healthcare costs). This Cochrane CR review portfolio remains dynamic, with three reviews having undergone an update in the last 12 months (Karmali 2014; Taylor 2014a; Taylor 2014b).

The portfolio of Cochrane reviews has played an important role in informing evidence-based policy for CR nationally and internationally, and the reviews have been cited in several key clinical guidelines including those by the ACC/AHA, ESC and NICE, which consistently recommend CR as a safe and effective intervention (Balady 2011; Perk 2012; McKelvie 2013; NICE 2013b; Task Force Members 2013; Yancy 2013).

Overviews of systematic reviews are a new approach to summarising evidence, synthesising results from multiple systematic reviews into a single usable document (Becker 2011). By providing a single synthesis of all relevant evidence in a particular area, overviews may be useful for therapeutic and policy decision-making, providing a comprehensive 'friendly front end' to the evidence, so that the reader does not have to assimilate the data from separate systematic reviews. Overviews can also help inform the strategic direction of conduct and structuring of future systematic reviews. For example, the latest version of the Cochrane review of exercise-based CR for CHD included 47 RCTs in over 10,000 participants and may, therefore, benefit from being organised into sub-reviews ('splitting') according to CHD indications (i.e. post-MI, revascularisation and angina). Finally, overviews provide an opportunity to identify potential 'evidence gaps' and, therefore, inform areas in which new Cochrane reviews should be prioritised.

OBJECTIVES

To conduct an overview of Cochrane systematic reviews to provide a contemporary review of the evidence for delivery of cardiac rehabilitation, to identify opportunities for merging or splitting existing Cochrane reviews, and to identify current evidence gaps to inform new cardiac rehabilitation systematic review titles.

METHODS

We conducted this overview in accordance with the recommendations for Cochrane overviews (Becker 2011).

Criteria for considering reviews for inclusion

We initially included the portfolio of six Cochrane CR reviews that were already known to us. In addition, we sought to include any other Cochrane reviews that may inform the aims of this overview including those that assessed the efficiency of CR services or that compared the delivery of CR across different settings.

Types of reviews

We included Cochrane reviews and protocols currently published in *The Cochrane Library* that examined the impact of CR. Given the targeted aims of this overview, we did not consider non-Cochrane systematic reviews.

Types of participants

We included adults aged 18 or over, with heart disease, regardless of aetiology.

Types of interventions

For the purposes of this review, we defined CR as: exercise with or without education with or without psychological intervention, delivered to people with heart disease, in a hospital community or a home-based setting.

Types of outcome

Patient-related outcomes

- Mortality:
 - cardiovascular mortality and non-cardiovascular mortality.
- Morbidity:
 - MI (total MI, fatal MI and non-fatal MI);
 - total revascularisations (coronary artery bypass graft (CABG), percutaneous transluminal coronary angioplasty (PTCA) and re-stenting);
 - total hospitalisations (cardiovascular hospitalisations and other hospitalisations);
 - HRQoL assessed using validated instruments (e.g. 36-item Short Form (SF-36), EQ5D).

Process-related outcomes

- Measure of uptake of, or adherence to, CR.
- Costs and cost-effectiveness.

Search methods for identification of reviews

We searched The Cochrane Database of Systematic Reviews (2014, Issue 10) using the search strategy listed in [Appendix 1](#). We applied no date or language restrictions. Where reviews had been updated, we sought only the most recent version. We sought full Cochrane reviews or protocols currently published that:

- examined the impact or delivery of CR;
- included adults with heart disease, regardless of aetiology;
- included exercise training interventions either alone or in combination with an educational or psychological intervention or both, delivered in a hospital community or a home-based setting.

Data collection and analysis

Selection of reviews

Two authors (LA, RST) independently screened the titles and abstracts of all of the Cochrane systematic reviews identified as a result of the search for inclusion, and coded them as 'retrieve' (eligible or potentially eligible/unclear) or 'do not retrieve'. We retrieved the full-text of selected reviews and two authors (LA, RST) independently screened the full-text and identified reviews for inclusion, and identified and recorded reasons for exclusion of the ineligible reviews. We resolved any disagreements through discussion.

Data extraction and management

We used standardised data collection forms to extract characteristics of reviews and included studies and outcome data. We piloted these forms on one review included in the overview. One author (LA) extracted review and study characteristics and outcome data from included reviews and a second author (RST) checked all extracted data for accuracy. We resolved disagreements by consensus. If study level information within a published review was unclear or missing, we clarified this by reference to the published reports of the individual RCT. One author (LA) transferred extracted data into the Review Manager 5 ([RevMan 2014](#)), and a second author (RST) spot-checked data for accuracy against the systematic reviews.

We extracted the following information from included Cochrane reviews: review objectives or question, search time frame, inclusion criteria (study design, population, intervention, comparator and outcomes), source of funding and stated conflicts of interest of review authors.

We extracted the following characteristics of the RCTs included in each of the Cochrane reviews: number of included trials, year of publication, population, intervention and comparator, primary and secondary outcomes specified and collected, total duration of study, number of study centres and location.

We sought the following outcome data:

- all-cause and disease-specific mortality;
- morbidity: fatal and non-fatal MI; percutaneous coronary intervention (PCI); hospitalisation: overall and disease-specific;
- HRQoL assessed using validated instruments (e.g. SF-36, EQ5D);
- measures of uptake of, or adherence to, CR; and
- costs and cost-effectiveness.

We did not re-assess the risk of bias of included studies within reviews, but instead reported according to the review authors' assessment using The Cochrane Collaboration 'Risk of bias' tool (in accordance with the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#))). The standard 'Risk of bias' items include: random sequence generation and allocation concealment, description of drop-outs and withdrawals, blinding of outcome assessment and presence of selective reporting. In addition, we sought evidence that the groups were balanced at baseline, that an intention-to-treat analysis was undertaken and that groups received comparable care (apart from the intervention). Where a 'Risk of bias' element was not reported within the review, one author (LA) assessed the original included study publication and a second author (RST) checked the details.

Assessment of methodological quality of included reviews

Quality of included Cochrane reviews

One author (LA) independently assessed the methodological quality of the included reviews using the 'Revised Assessment of Multiple Systematic Reviews' (R-AMSTAR) measurement tool (Kung 2010), where the 11 domains of the original AMSTAR tool (Shea 2009) were scored between 1 and 4 and the R-AMSTAR total score ranged from 11 to 44. We resolved any disagreements by discussion. A second author (RST) checked the assessment.

Quality of evidence in included reviews

One author (LA) used GRADEProfiler software to assess the quality of evidence for outcomes reported in, and extracted from, each of the reviews (GRADEpro 2008), based on the following factors: indirectness of evidence, unexplained heterogeneity, publication bias, risk of bias due to study design limitations and imprecision of results (Balslem 2011). A second author (RST) checked the assessment.

Assessment of bias in conducting the overview

We conducted the overview according to the published protocol and we have reported any deviations from it in the [Differences between protocol and review](#) section of this overview.

Data synthesis

The unit of analysis for this overview is the systematic reviews (and not individual trials). The focus of the data presentation was descriptive, with detailed tabular presentations of the extracted review- and trial-level information outlined above. We conducted no de novo data analysis of trial-level outcomes for this overview. We have tabulated review-level summaries for all the outcomes listed above from each of the included reviews. Where outcomes

were meta-analysed within a review, we extracted and reported pooled effect sizes. Where no quantitative pooling of effect sizes was reported, or where outcomes were reported descriptively by single studies, we reported these results by vote counting (Bushman 1984), or using standardised language indicating direction of effect and statistical significance. For continuous outcomes, we summarised data using the standardised mean difference (SMD) or mean difference (MD) with 95% confidence interval (CI) as reported in the included reviews. For dichotomous outcomes, we presented the risk ratio (RR) or odds ratio (OR) and 95% CI as appropriate.


Due to the heterogeneity of populations, interventions and outcomes in the included systematic reviews, we did not seek to compare either CR interventions directly across reviews (e.g. exercise CR versus education for CHD) or to compare interventions across review populations (e.g. exercise CR for CHD versus exercise CR for HF). For this reason, we did not attempt to compare outcome results across trials using indirect network meta-analysis methods.

RESULTS

Identification of reviews

Figure 1 summarises the review selection process in a flow diagram. Our database search yielded 92 titles from which we identified one published Cochrane review (in addition to the previously identified six reviews) and four Cochrane review protocols that we judged to meet the inclusion criteria. On review of the full text, we excluded the published Cochrane review (Hulzebos 2012), as it evaluated physical therapy with an exercise component for elective cardiac surgery patients and included only one RCT of exercise training, while the other RCTs assessed inspiratory muscle training (Appendix 2). We judged the four Cochrane protocols to meet the inclusion criteria (see Appendix 3). The remainder of this overview focused on presenting the six Cochrane CR reviews.

Figure 1. Study flow diagram.



Description of included reviews

The characteristics of the six included Cochrane reviews are summarised in [Table 1](#) and included RCTs are summarised [Table 2](#).

All included reviews ran searches from the inception of the electronic databases to 2013, and were published between 2011 and 2014. In all reviews, searches were limited to an RCT design and in three cases the inclusion was limited to RCTs with follow-up of six months or longer ([Brown 2011](#); [Heran 2011](#); [Taylor 2014b](#)). In total, the included reviews contained 148 RCTs and 97,486 participants. Four RCTs were included in more than one review ([Stern 1983](#); [Miller 1984](#); [PRECOR 1991](#); [Lisspers 1999](#)). Most included RCTs were published since the mid-1990 (1970 to 1979: 4 RCTs; 1980 to 1989: 16 RCTs; 1990 to 1999: 40 RCTs; 2000 to 2009: 72 RCTs, 2010 and later: 16 RCTs). The median sample size of RCTs ranged widely from only 16 participants ([Duncan 2003](#)) to 46,606 participants ([Esposito 2008](#)). Most RCTs were undertaken in either Europe (69%) or North America (35%) and were mainly single centre (79%).

Search methods

All six reviews searched the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, MEDLINE In-Process, EMBASE and CINAHL. In addition, four of the reviews searched PsycINFO ([Brown 2011](#); [Heran 2011](#); [Karmali 2014](#); [Taylor 2014a](#)). Three reviews also undertook searches for ongoing RCTs using trial registers (International Standard Randomized Controlled Trial Number (ISRCTN) registry (www.controlled-trials.com) and ClinicalTrials.gov (clinicaltrials.gov/)) and all reviews searched for additional RCTs by manually checking the reference lists of included studies.

Participants

The types of participants included in this overview varied between reviews. The scope of two reviews included all adults with heart disease, regardless of indication ([Karmali 2014](#); [Taylor 2014a](#)), three reviews were limited to people with CHD (post-MI and PCI) ([Brown 2011](#); [Heran 2011](#); [Whalley 2011](#)), and one review was limited to HF ([Taylor 2014b](#)). Although 78% of the RCTs that reported gender included women, the median proportion of men included in RCTs ranged from 60% to 88% across reviews. The mean age of participants in RCTs ranged from 46 to 87 years.

Interventions

Two of the reviews included exercise training or exercise training alongside other interventions that included education or psychological support or both ([Heran 2011](#); [Taylor 2014b](#)). One review focused on psychological interventions ([Whalley 2011](#)), although it included several RCTs that also incorporated an educational component. One review included only RCTs with an educational focus ([Brown 2011](#)), one included interventions to increase the uptake and adherence to CR ([Karmali 2014](#)), and one review compared the delivery of CR in home- and centre-based settings ([Taylor 2014a](#)).

Outcome measures

All reviews pre-specified outcome measures that consistently included all-cause mortality and HRQoL. Although all reviews sought morbidity outcomes, the definition and breadth of these outcomes varied across reviews. For example, the review by Heran et al. stated that they sought MI (total, fatal,

non-fatal), revascularisations (total, CABG, PTCA, stenting) and hospitalisations ([Heran 2011](#), while the review by Karmali et al. reported "CHD event rates" ([Karmali 2014](#)). Four reviews sought economic outcomes ([Brown 2011](#); [Heran 2011](#); [Taylor 2014a](#); [Taylor 2014b](#)), and two reviews reported collected uptake or adherence data ([Karmali 2014](#); [Taylor 2014a](#)).

Data analysis

Five of the six reviews included meta-analyses of mortality and morbidity outcomes. The review by Karmali et al. pre-stated that heterogeneity (participants, interventions and outcomes), together with the small number of studies identified, precluded undertaking meta-analysis ([Karmali 2014](#)). Given the heterogeneity in measures, only one review used meta-analysis to pool HRQoL data across RCTs ([Taylor 2014b](#)), the other reviews used a descriptive or vote counting approach to summarise outcomes. This was also the case for uptake and adherence and economic outcomes. Two of the reviews undertook meta-regression analyses to explore how the impact of interventions varied across participant and RCT characteristics ([Heran 2011](#); [Taylor 2014b](#)).

Methodological quality of included reviews

Based on our assessments using the R-AMSTAR tool, all included reviews scored between 35 and 41 (out of a possible maximum of 44) and we deemed them of high methodological quality (see [Table 3](#)). None of the reviews stated that journals were hand or manually searched and only one stated that searches were supplemented by consulting books or experts in the field. Two reviews were marked down based on inadequate reporting of the publication status of their included studies. None of the reviews rated the quality of evidence based on a characterised instrument such as GRADE, and while all reviews used The Cochrane Collaboration 'Risk of bias' tool, most were marked down as they did not refer to the quality of included studies in formulating recommendations. The two weaknesses identified across reviews by R-AMSTAR were the lack of an explicit statement on the impact of findings on clinical practice guidelines and the failure to assess the sources of support or conflict of interest in the included RCTs.

Risk of bias of included randomised controlled trials

All six Cochrane reviews used the core items of The Cochrane Collaboration 'Risk of bias' tool (see [Table 4](#)). A consistent finding across reviews was that the included RCTs often did not give enough detail to assess the adequacy of their potential risk of bias. Where details were reported, the quality of RCTs appeared to vary considerably across the risk of bias items. Across all reviews, only a minority of RCTs were judged to be 'adequate' in terms of sequence generation (31%), sequence concealment (29%) and outcome blinding (24%). Other aspects of RCT quality (baseline balance, selective reporting, loss of follow-up, intention-to-treat analysis and groups receiving same intervention) were judged to be better (greater than 50% of all included RCTs achieving adequacy).

Quality of evidence from randomised controlled trials in included reviews

The quality of the evidence reported by the RCTs in the included reviews was rated using the GRADE method. The quality of the evidence varied widely (by review and by outcome) and ranged from very low to moderate. See [Table 5](#); [Table 6](#); [Table 7](#); [Table 8](#); and [Table 9](#) for details.

Effect of interventions

Table 10 summarises the outcome results across included Cochrane reviews.

Exercise-based cardiac rehabilitation for coronary heart disease (Heran 2011)

The Heran et al. review undertook database searches up to December 2009 with the inclusion of RCTs with six months or more of follow-up comparing CR with no CR control (Heran 2011). The review included 47 RCTs with 10,794 participants who were mainly post-MI or post-PCI, predominantly men (median 88%) and with a median mean age of 55.0 years (see Table 2). CR programmes differed considerably across RCTs in duration (range one to 30 months), frequency (one to seven sessions/week) and session length (20 to 90 minutes/session), and included both exercise-only CR programmes and comprehensive CR programmes (exercise plus psychological or education intervention, or both). We judged this review to be of good methodological quality, with an R-AMSTAR score of 39.

With follow-up of six to 12 months, there was a trend towards a reduction in total mortality (RR 0.82; 95% CI 0.67 to 1.01; low GRADE rating) and no difference was seen between groups in cardiovascular mortality (RR 0.93; 95% CI 0.71 to 1.21; low GRADE rating). However, with follow-up of 12 months or more, CR reduced overall (RR 0.87; 95% CI 0.75 to 0.99; moderate GRADE rating) and cardiovascular mortality (RR 0.74; 95% CI 0.63 to 0.87; moderate GRADE rating). There was no evidence of a difference in risk of reinfarction or PCI between CR and control. Ten of the included studies (2379 participants) reported hospital admissions. In the shorter term (less than 12 months' follow-up), hospital admissions were reduced compared with control (RR 0.69; 95% CI 0.51 to 0.93; moderate GRADE rating) but there was no evidence of a reduction in the longer term (greater than 12 months' follow-up) (RR 0.98; 95% CI 0.87 to 1.11; low GRADE rating). There was no evidence of heterogeneity of effect across RCTs for any of the mortality or morbidity outcomes. Univariate meta-regression showed no differences in intervention effects across various participant and RCT characteristics in mortality or morbidity outcomes. In seven out of 10 RCTs, there was evidence of a significantly higher level of HRQoL with CR than control. Three of the included studies reported data on patient costs, their direct comparison limited by differences in currencies and the time when the studies were conducted.

Exercise-based rehabilitation for heart failure (Taylor 2014b)

The Taylor et al. review was updated with searches up to March 2013 and included 33 RCTs with six months or more of follow-up comparing CR with no CR control in 4676 participants with HF (Taylor 2014b). Participants were predominantly men (median 80%) with a median age of 60 years, had a reduced ejection fraction (HFREF less than 40% to 45%) and New York Heart Association classification I to III (see Table 2). The exercise regimen ranged widely across RCTs from a session duration of 15 to 120 minutes, from one to seven sessions/week, and from intensity of 40% to 80% of maximal heart rate (or equivalent) over a period from one to 120 months and included both exercise-only CR programmes and comprehensive CR programmes (exercise plus psychological or education (or both) intervention). We judged this review to be of good methodological quality, with an R-AMSTAR score of 39.

There was no evidence of difference in pooled mortality between intervention and controls at six- to 12-month follow-up (RR 0.93; 95% CI 0.69 to 1.27; low GRADE rating). However, in the six RCTs with more than 12 months' follow-up, there was a trend towards a reduction in all-cause mortality with exercise (RR 0.88; 95% CI 0.75 to 1.02; low GRADE rating). Compared with control, exercise training reduced the risk of overall hospitalisation (RR 0.75; 95% CI 0.62 to 0.92; moderate GRADE rating) and HF-specific hospitalisation (RR 0.61; 95% CI 0.46 to 0.80; moderate GRADE rating) although there was no difference in all hospital admissions at beyond 12-month follow-up (RR 0.92; 95% CI 0.66 to 1.29; very low GRADE rating). Exercise resulted in a clinically important improvement in the Minnesota Living with Heart Failure (MLWHF) questionnaire (MD -5.8 points, -9.2 to -2.4; very low GRADE rating) although there was evidence of high levels of statistical heterogeneity ($I^2 = 70%$). Univariate meta-regression analysis showed that these benefits in hospitalisation and HRQoL were independent of participant characteristics (age, gender, left ventricular ejection fraction), type of CR (exercise only versus comprehensive CR), exercise-based CR regimen, length of follow-up, overall risk of bias, RCT publication date, single versus multicentre RCT or CR setting (home versus centre-based). There was limited evidence to support CR for people with HF with HFPEF (three RCTs, undefined participant number) and when exclusively delivered in a home-based setting (five RCTs, 521 participants).

Three RCTs reported economic data. Although no group differences in costs or outcomes across these three studies achieved statistical significance, two studies indicated CR to be cost-effective (USD1773 per life-year saved; Georgiou 2001), and a mean gain in quality-adjusted life-year (QALY) of 0.03 at an additional mean cost of USD1161 per person (Flynn 2009).

Psychological interventions for coronary heart disease (Whalley 2011)

The Whalley et al. review undertook searches up to January 2009 with the inclusion of RCTs of psychological interventions compared with usual care in people with a diagnosis coronary artery disease (Whalley 2011). The review included 24 RCTs in 9296 participants who were predominantly low-risk post-MI or PCI, male (median 84%) with a median mean age of 57 years (see Table 2). The review authors reported substantial variability in the intensity of treatments offered across RCTs; the mean number of hours spent in treatment was 26.1 hours (2.4 to 96). Included trials applied both psychological-only CR programmes and comprehensive CR programmes (psychological and education interventions).

Most interventions were based on group therapy sessions or comprised a mix of group and individual session; only four RCTs used treatments that were delivered only on an individual basis. We judged the review to be of good methodological quality, with an R-AMSTAR score of 39. There was evidence of a trend towards a reduction in all-cause mortality (RR 0.89; 95% CI 0.75 to 1.05; low GRADE rating) and fewer cardiac deaths with psychological intervention (RR 0.80; 95% CI 0.64 to 1.00; low GRADE rating). There were significant effects on occurrence of revascularisation (RR 0.95; 95% CI 0.80 to 1.13; moderate GRADE rating) and non-fatal reinfarction (RR 0.87; 95% CI 0.67 to 1.13; low GRADE rating). One of seven studies reported superiority in HRQoL with psychological intervention compared with usual care.

Patient education in the management of coronary heart disease (Brown 2011)

The Brown et al. review undertook searches up to August 2010 with the inclusion of RCTs (with follow-up of six months or more) of patient education interventions compared with usual care (Brown 2011). The review included 13 RCTs in 68,556 participants with HF, stable angina, and post-MI and PCI who were predominantly male (median 60%) with a median mean age of 62.0 years (see Table 2). Interventions varied considerably across RCTs, with some providing group sessions, some individualised education and others both. Educational regimen ranged from two clinic visits to a four-week residential stay with 11 months of follow-up sessions. All included trials were limited to educational interventions and did not use other CR interventions of exercise or psychological support. We judged this review to be of good methodological quality, with an R-AMSTAR score of 41. There was no evidence of a significant difference in total mortality (RR 0.79; 95% CI 0.55 to 1.13; moderate GRADE rating) or morbidity (MI: RR 0.63; 95% CI 0.26 to 1.48, low GRADE rating; CABG: RR 0.58; 95% CI 0.19 to 1.71, low GRADE rating; hospitalisation: RR 0.83; 95% CI 0.65 to 1.07, moderate GRADE rating). Across the 11 studies that reported HRQoL, while there was no consistent difference in HRQoL total or domain score at follow-up between intervention and control, five RCTs demonstrated statistically significant differences in some domains in favour of intervention. Five RCTs reported healthcare utilisation and costs. Given the small number of included RCTs, the authors deemed meta-regression analysis inappropriate.

Home-based versus centre-based cardiac rehabilitation (Taylor 2014a)

The Taylor et al. review was updated with searches up to November 2012 and sought to include RCTs comparing home-based and centre-based CR (Taylor 2014a). Home-based CR was defined as "a structured programme with clear objectives for the participants, including monitoring, follow up visits, letters or telephone calls from staff, or at least self-monitoring diaries" and centre-based CR was defined as "based in a variety of settings (e.g. hospital physiotherapy department, university gymnasium, community sports centre)". The review included 17 RCTs in 2172 people with stable angina, HF and post-MI and PCI who were predominantly male (median 80%) with a median mean age of 60 years (see Table 2). Most RCTs compared comprehensive programmes (i.e. exercise training plus education or psychological (or both) interventions) with the exercise components differing considerably across RCTs in duration (range 1.5 to six months), frequency (one to five sessions per week) and session length (20 to 60 minutes per session). Included trials applied both exercise-only CR programmes and comprehensive CR programmes (exercise plus psychological or education (or both) intervention). We judged the review to be of good methodological quality, with an R-AMSTAR score of 40. There was evidence of a difference in mortality at three to 12 months' follow-up between home and centre CR (RR 0.79; 95% CI 0.43 to 1.47; low GRADE rating). Four studies reported cardiac events, but no pooling of data was possible due to differences in the nature of the reported events. There was no evidence of difference between the two settings in overall or domain HRQoL scores in individual RCTs. Four out of the 14 studies reporting adherence found superior adherence in the home-based compared with centre-based CR setting. There was no consistent difference in the healthcare costs associated with the two forms of CR, although difference in currencies and timing of studies meant that it was not

possible to compare costs directly across studies. In three of the four studies, the healthcare costs associated with CR were lower for the home-based than centre-based programmes, although this was significantly lower in only one study (GBP170 per participant versus GBP200 per participant; difference of -GBP 30, 95% CI -45 to -12; P value < 0.0001; Dalal 2007). Jolly et al. found that home-based CR was more expensive than centre-based CR (GBP198 per participant versus GBP157 per participant; P value < 0.05; Jolly 2007), although the costs of two would be the same if participant travel costs and travel time were included. Given the small number of included RCTs, the authors deemed meta-regression analysis inappropriate.

Promoting participant uptake and adherence in cardiac rehabilitation (Karmali 2014)

The Karmali et al. review was updated with searches up to January 2013 and sought to include RCTs of interventions to increase CR uptake (participants attendance or enrolment in CR programmes) or adherence (extent to which the participant's behaviour concurred with the advice given by health professional, e.g. to attend CR meetings or to undertake independent exercise) (Karmali 2014). The review included 18 RCTs in 2505 participants with HF, stable angina, and post-MI and PCI who were predominantly male (median 84%). We judged this review to be of good methodological quality, with an R-AMSTAR score of 35. Meta-analysis and meta-regression was not undertaken due to heterogeneity in outcome definition across RCTs. Of the 10 RCTs (1658 participants) evaluating the effectiveness of interventions to increase uptake of CR, eight reported higher rates of CR uptake in the intervention group (range 11% to 46%). Uptake was variously defined in these studies as enrolment in CR, attendance at a variety of time points or by number of sessions over a 12-week period. Interventions that improved uptake of CR included: structured nurse- or therapist-led contacts, early appointments after discharge, motivational letters, gender-specific programmes and intermediate-phase programmes for elderly people. Three out of eight RCTs (1167 participants) found significant improvements in adherence to CR although there was no evidence of an improvement in HRQoL. Interventions that improved adherence included self monitoring of activity, action planning and tailored counselling by CR staff. Although data were limited, there was no evidence of a difference in mortality or morbidity with uptake or adherence interventions. No RCTs reported on costs or cost-effectiveness.

DISCUSSION

Summary of main results

CR programmes have become an integral part of the standard of care for people with heart disease. The scope of contemporary CR has shifted from exercise interventions alone to more comprehensive secondary prevention programmes that include risk factor education and psychological support. This overview identified six Cochrane systematic reviews of RCTs that have assessed the outcomes of various aspects of the delivery of CR and its component interventions. The key outcome findings of our overview were:

- exercise-based CR in low-risk people with HF and after MI or PCI, is safe, with no increase in short-term mortality, and effective in terms of reductions in the risk of hospital admission and improvements in patient HRQoL, compared with control. While there was considerable evidence of heterogeneity

across included primary studies in both the characteristics of the evaluated CR programmes and also across the included participants, the outcome benefits of CR in terms of HRQoL and reduced hospitalisation appeared to be independent of these programme and participant characteristics;

- psychological-based and education-based interventions alone appear to have little or no impact on mortality or hospitalisation, but may improve HRQoL of people with CHD in comparison with usual care alone;
- home-based and centre-based programmes seem to be equally effective in improving the outcomes of exercise-based CR in low-risk people after MI or post-revascularisation or with HF. Healthcare costs of the two forms of CR were similar, presumably as any cost reduction in delivering the intervention in the home was offset by the associated costs of delivering individual nursing care; and
- uptake of CR programmes was only weakly supported by interventions designed to improve adherence to CR programmes.

Overall completeness and applicability of evidence

There are a number of published non-Cochrane systematic reviews of CR (Oldridge 1988; O'Connor 1989; Brown 2003; Piepoli 2004; Haykowsky 2007; Hwang 2009; Lawler 2011; Oldridge 2012). Given that our focus was Cochrane reviews, we acknowledge that this overview cannot be regarded as an all-inclusive summary of the evidence base for CR. However, by focusing on high-quality Cochrane reviews, we believe this overview potentially provides a least biased estimate of the impact of CR.

Quality of the evidence

The included Cochrane systematic reviews were generally of high quality and three had been updated with a literature search since 2011 (Karmali 2014; Taylor 2014a; Taylor 2014b). However, the quality of the primary RCTs in the included systematic reviews was variable. The main sources of bias in the primary studies were inadequate reporting of allocation concealment and randomisation methods and lack of outcome blinding. These limitations in the methodological quality led to the downgrading of the quality of the evidence, which varied by outcome within each review. Other reasons for downgrading the evidence included heterogeneity or inconsistency of effect, and imprecision of results. Another potential source of inconsistency that was not reported in the reviews was differential use of outcome data by RCTs (i.e. some studies analysed only post-interventional data while others measured pre-post change).

Potential biases in the overview process

This overview included RCTs conducted between 1974 and 2013. During this time, there have been major advances in medical management, such as the increased use of statins since the mid-1990s. Indeed, it has been hypothesised that major advances in post-MI medical management since the mid-2000s has led to a reduction in the incremental effect on mortality of CR compared with usual care alone (Taylor 2012). This decrement in mortality benefit associated with CR was supported by the Rehabilitation After Myocardial Infarction Trial (RAMIT), which was published after the search cut of the exercise-based CR for CHD Cochrane review. This trial randomised 1813 participants in 14 hospitals in England and Wales to receive either comprehensive CR or usual care and

found no difference in all-cause mortality at two years (RR 0.98; 95% CI 0.74 to 1.30) or after seven to nine years (RR 0.99; 95% CI 0.85 to 1.15) (West 2012). This RCT was published after the search cut off of the exercise-based CR for CHD Cochrane review.

A potential strength of an overview is that it can provide an opportunity to undertake indirect comparisons across interventions that might not be included in single systematic reviews using mixed treatment comparisons and network meta-analysis methods (Becker 2011; Mills 2013). In brief, an indirect comparison involves the comparison of two (or more) interventions via one or more common comparator. For example, we may seek to compare the impact of exercise-based interventions and psychological-based interventions via the combination of RCTs of exercise-based intervention versus usual care with RCTs of psychological-based intervention versus usual care. However, for the intervention effect determined using an indirect comparison to be valid and equivalent to the intervention effect measured using a direct comparison, the sets of RCTs used to obtain the indirect comparison need to be sufficiently similar in their characteristics (i.e. patient population, intervention, comparator and outcomes across trials need to be similar - the transitivity assumption) (Cipriani 2013). Given the substantial heterogeneity in the populations of the included CR RCTs, not only between, but also within the included CR systematic reviews, we deemed indirect comparisons as inappropriate in the case of this overview. Based on the same reasoning, readers of this overview need to apply considerable caution in taking an informal indirect comparison approach and comparing the results for a given outcome across reviews.

Agreements and disagreements with other studies or reviews

In 2012, Oldridge undertook an overview of meta-analyses of CR in people with CHD (Oldridge 2012). Given that this overview included both Cochrane and non-Cochrane meta-analyses published since 2000, there is considerable overlap in findings and conclusions with the present overview. One important difference between the two overviews is the conclusion of a reduction in all-cause and cardiovascular mortality with CR in the overview by Oldridge (Oldridge 2012). This mortality benefit was primarily seen in three non-Cochrane meta-analyses (Taylor 2004; Clark 2005; Lawler 2011), while the Cochrane review found a statistically significant reduction in all-cause and cardiac mortality only at follow-up of greater than 12 months (Heran 2011).

AUTHORS' CONCLUSIONS

Implications for practice

The evidence compiled by this overview supports current international clinical guidelines that state that the addition of cardiac rehabilitation (CR) to medical management is effective (improving health-related quality of life (HRQoL) and reducing the risk of future hospitalisations) and safe (with no increase in short-term mortality), compared with a no exercise training control, for clinically stable participants following myocardial infarction (MI) or percutaneous coronary intervention (PCI) or who have heart failure (Balady 2011; Perk 2012; McKelvie 2013; Task Force Members 2013; NICE 2013; Yancy 2013). Future randomised controlled trials (RCTs) of CR need to improve their reporting methods and better reflect the real world practice including the recruitment of higher-

risk participants and consideration of contemporary models of CR delivery, and identify effective interventions for enhancing adherence to rehabilitation.

Implications for research

Based on this overview, and taking account of recent guidelines for the conduct of systematic review of complex interventions (Weir 2012; Petticrew 2013), we make the following recommendations for the conduct of future CR systematic reviews:

- **Scope of reviews:** the scope of CR reviews needs to reflect current guidelines that consistently recommend that CR should be based on an individually prescribed programme of exercise training with appropriate co-intervention including psychological or educational interventions (BACPR 2012; McMurray 2012; NICE 2013; Yancy 2013).
- **Handling of the complexity of CR:** given that CR is a complex intervention, a key challenge of systematic reviews of CR is taking account of the potential heterogeneity in CR interventions (content and methods of delivery) and the population of people who receive CR. Future reviews of CR need to explore this complexity using approaches that include stratification ('splitting') of outcome results by patient indication (e.g. post-MI versus post-PCI) or intervention type (i.e. exercise training only versus comprehensive CR interventions); reporting within RCT subgroup analyses; and use of meta-regression to explore the association between intervention characteristics and outcomes across trials. Consideration should also be given to the appropriate use of indirect comparison methods (Bucher 1997) in reviews or broadening the inclusion criteria of reviews to include active comparator arms of RCTs that would allow assessment of the comparative effectiveness of different CR interventions (or both). Theory-based approaches to systematic reviews of CR are also needed (Gardner 2010).
- **Consistency in review conduct and reporting:** to facilitate comparison across CR systematic reviews and the efficient future update of this overview, future Cochrane CR reviews need to standardise their methods and reporting, including the reporting of included RCT characteristics, risk of bias assessment criteria, outcomes and evidence synthesis approaches.

In addition to the current Cochrane CR reviews that are in protocol and title stage (Devi 2011; Euler 2013; Sibilitz 2013; Mechta-Nielsen 2014; Risom 2014), consideration should be given to new Cochrane titles to fill the CR evidence gaps identified by this overview, including 'exercise-based CR for post-cardiac transplantation' and 'exercise-based CR for congenital heart disease'.

This overview also highlights several potential areas for consideration in the conduct of future RCTs of CR. RCT recruitment criteria need to reflect the real world of CR delivery better, which includes people at higher risk who are older, female and from a broader range of ethnicities and socio-economic groups. Reporting of trial methods should be improved, with greater details of the process of randomisation and outcome blinding, more precise description of the intervention, and consistency in the collection and reporting of outcome measures, including the use of validated HRQoL instruments, cardiac-related events, re-admissions and costs. Finally, as noted by Clark and colleagues, future RCTs need to "open the black box" of CR better (Clark 2013). In other words, to determine the incremental benefits of the various components of CR requires future RCTs to provide more precise descriptions of their CR interventions so these comparisons can be more explicitly and reliably undertaken in future systematic reviews. One publication provides researchers and clinicians with a framework to improved reporting of intervention detail (Hoffmann 2014). In addition, the design of future RCTs should consider 'head-to-head' comparisons of different combinations of CR interventions (e.g. an 'exercise-only' CR intervention compared with 'exercise plus' CR intervention).

ACKNOWLEDGEMENTS

We are grateful for the help of Nicole Martin at the Cochrane Heart Group, for her help in developing the search strategy for this overview. We are extremely grateful to our external reference group members (Dr Mike Knapton, Prof Gill Furze, Dr Neil Smart, Prof Patrick Doherty, Dr Ann-Dorthe Zwisler, Prof Neil Oldridge, Dr Phil Adams, Dr Christopher O'Conner, Dr Juan Pablo-Casas and Dr Mark Huffman) for their invaluable advice and support in the preparation of this manuscript.

REFERENCES

References to included reviews

Brown 2011

Brown JP, Clark AM, Dalal H, Welch K, Taylor RS. Patient education in the management of coronary heart disease. *Cochrane Database of Systematic Reviews* 2011, Issue 12. Art. No: CD008895. [DOI: [10.1002/14651858.CD008895.pub2](https://doi.org/10.1002/14651858.CD008895.pub2)]

Heran 2011

Heran BS, Chen JM, Ebrahim S, Moxham T, Oldridge N, Rees K, et al. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database of Systematic Reviews* 2011, Issue 7. Art. No: CD001800. [DOI: [10.1002/14651858.CD001800.pub2](https://doi.org/10.1002/14651858.CD001800.pub2)]

Karmali 2014

Karmali K, Davies P, Taylor F, Beswick A, Martin N, Ebrahim S. Promoting patient uptake and adherence in cardiac rehabilitation. *Cochrane Database of Systematic Reviews* 2014, Issue 6. Art. No: CD007131. [DOI: [10.1002/14651858.CD007131.pub3](https://doi.org/10.1002/14651858.CD007131.pub3)]

Taylor 2014a

Taylor RS, Dalal H, Jolly K, Zawada A. Home-based versus centre-based cardiac rehabilitation. *Cochrane Database of Systematic Reviews* 2014, Issue in press. Art. No: CD007130. [DOI: [10.1002/14651858.CD007130.pub2](https://doi.org/10.1002/14651858.CD007130.pub2)]

Taylor 2014b

Taylor RS, Sager VA, Davies EJ, Briscoe S, Coats AJ, Dalal H, et al. Exercise-based rehabilitation for heart failure. *Cochrane Database of Systematic Reviews* 2014, Issue 4. Art. No: CD003331. [DOI: [10.1002/14651858.CD003331.pub4](https://doi.org/10.1002/14651858.CD003331.pub4)]

Whalley 2011

Whalley B, Rees K, Davies P, Bennett P, Ebrahim S, Liu Z, et al. Psychological interventions for coronary heart disease. *Cochrane Database of Systematic Reviews* 2011, Issue 8. Art. No: CD002902. [DOI: [10.1002/14651858.CD002902.pub3](https://doi.org/10.1002/14651858.CD002902.pub3)]

References to excluded reviews

Hulzebos 2012

Hulzebos EHJ, Smit Y, Helden PPJM, van Meeteren NLU. Preoperative physical therapy for elective cardiac surgery patients. *Cochrane Database of Systematic Reviews* 2012, Issue 11. Art. No: CD010118. [DOI: [10.1002/14651858.CD010118.pub2](https://doi.org/10.1002/14651858.CD010118.pub2)]

Additional references

Allender 2008

Allender S, Scarborough P, O'Flaherty M, Capewell S. Patterns of coronary heart disease mortality over the 20th century in England and Wales: possible plateaus in the rate of decline. *BMC Public Health* 2008;**8**:148.

BACPR 2012

British Association for Cardiovascular Prevention and Rehabilitation. The BACPR standards and core components

for cardiovascular disease prevention and rehabilitation, 2nd edition, 2012. www.bacpr.com/resources/46C_BACPR_Standards_and_Core_Components_2012.pdf (accessed 7 December 2014).

Balady 2011

Balady GJ, Ades PA, Bittner VA, Franklin BA, Gordon NF, Thomas RJ, et al. Referral, enrolment, and delivery of cardiac rehabilitation/secondary prevention programs at clinical centers and beyond: a presidential advisory from the American Heart Association. *Circulation* 2011;**124**(25):2951-60.

Balshem 2011

Balshem H, Helfand M, Schunemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *Journal of Clinical Epidemiology* 2011;**64**:401-6.

Becker 2011

Becker LA, Oxman AD. Chapter 22: overviews of reviews. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Belardinelli 1998

Belardinelli R, Georgiou D, Ginzton L, Cianci G, Purcaro A. Effects of moderate exercise training on thallium uptake and contractile response to low-dose dobutamine of dysfunctional myocardium in patients with ischemic cardiomyopathy. *Circulation* 1998;**97**:553-61.

Bethell 2008

Bethell H, Lewin R, Evans J, Turner S, Allender S, Petersen S. Outpatient cardiac rehabilitation attendance in England: variability by region and clinical characteristics. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2008;**28**(6):386-91.

Black 1998

Black JL, Allison TG, Williams DE, Rummans TA, Gau GT. Effect of intervention for psychological distress on rehospitalization rates in cardiac rehabilitation patients. *Psychosomatics* 1998;**39**:134-43.

Brown 2003

Brown A, Taylor R, Noorani H, Stone JA, Skidmore B. Exercise-based cardiac rehabilitation programs for coronary artery disease: a systematic clinical and economic review. Canadian Coordinating office for Health Technology Assessment 2003;**Technology report 34**.

Bucher 1997

Bucher HC, Guyatt GH, Griffith LE, Walter SD. The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *Journal of Clinical Epidemiology* 1997;**50**:683-91.

Bushman 1984

Bushman BJ. Vote-counting procedures in meta-analysis. In: Cooper H, Hedges LV, editors(s). *The Handbook of Research Synthesis*. New York: Russell Sage Foundation, 1984:193-213.

Cipriani 2013

Cipriani A, Higgins JP, Geddes JR, Salanti G. Conceptual and technical challenges in network meta-analysis. *Annals of Internal Medicine* 2013;**159**:130-7.

Clark 2005

Clark AM, Hartling L, Vandermeer B, McAlister FA. Meta-analysis: secondary prevention programs for patients with coronary artery disease. *Annals of Internal Medicine* 2005;**143**:659-72.

Clark 2013

Clark AM. What are the components of complex interventions in healthcare? Theorizing approaches to parts, powers and the whole intervention. *Social Science & Medicine* 2013;**93**:185-93.

Clausen 1976

Clausen JP. Circulatory adjustments to dynamic exercise and effect of physical training in normal subjects and in patients with coronary artery disease. *Progress in Cardiovascular Disease* 1976;**18**:459-95.

Cowie 2000

Cowie MR, Wood DA, Coats AJ, Thompson SG, Suresh V, Poole-Wilson PA, et al. Survival of patients with a new diagnosis of heart failure: a population based study. *Heart* 2000;**83**:505-10.

Craig 2008

Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008;**337**:a1655.

Dalal 2007

Dalal HM, Evans PH, Campbell JL, Taylor RS, Watt A, Read KL, et al. Home-based versus hospital-based rehabilitation after myocardial infarction: a randomized trial with preference arms - Cornwall Heart Attack Rehabilitation Management Study (CHARMS). *International Journal of Cardiology* 2007;**119**:202-11.

Dalal 2012

Dalal HM, Wingham J, Palmer J, Taylor R, Petre C, Lewin R, et al. Why do so few patients with heart failure participate in cardiac rehabilitation? A cross-sectional survey from England, Wales and Northern Ireland. *BMJ Open* 2012;**2**:e000787.

Daly 2002

Daly J, Sindone AP, Thompson DR, Hancock K, Chang E, Davidson P. Barriers to participation in and adherence to cardiac rehabilitation programs: a critical literature review. *Progress in Cardiovascular Nursing* 2002;**17**:8-17.

Devi 2011

Devi R, Igbinedion E, Powell J, Singh S, Rees K. Internet based interventions for the secondary prevention of coronary heart disease. *Cochrane Database of Systematic Reviews* 2011, Issue 10. Art. No: CD009386. [DOI: [10.1002/14651858.CD009386](https://doi.org/10.1002/14651858.CD009386)]

Dickstein 2008

Dickstein K, Cohen-Solal A, Filippatos G, McMurray JJ, Ponikowski P, Poole-Wilson PA, et al. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the diagnosis and treatment of acute and chronic heart failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *European Journal of Heart Failure* 2008;**10**:933-89.

Duncan 2003

Duncan K, Pozehl B. Effects of an exercise adherence intervention on outcomes in patients with heart failure. *Rehabilitation Nursing* 2003;**28**:117-22.

Esposito 2008

Esposito D, Brown R, Chen A, Schore J, Shapiro R. Impacts of a disease management program for dually eligible beneficiaries. *Health Care Financing Review* 2008;**30**:27-45.

Euler 2013

Euler U, Wegewitz UE, Schmitt J, Adams J, van Dijk JL, Seidler A. Interventions to support return-to-work for patients with coronary heart disease. *Cochrane Database of Systematic Reviews* 2013, Issue 9. Art. No: CD010748. [DOI: [10.1002/14651858.CD010748](https://doi.org/10.1002/14651858.CD010748)]

Flynn 2009

Flynn KE, Pina IL, Whellan DJ, Lin L, Blumenthal JA, Ellis SJ, et al. Effects of exercise training on health status in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA* 2009;**301**:1451-9.

Gardner 2010

Gardner B, Whittington C, McAteer J, Eccles MP, Michie S. Using theory to synthesise evidence from behaviour change interventions: the example of audit and feedback. *Social Science & Medicine* 2010;**70**:1618-25.

Gaziano 2010

Gaziano TA, Bitton A, Anand S, Abrahams-Gessel S, Murphy A. Growing epidemic of coronary heart disease in low- and middle-income countries. *Current Problems in Cardiology* 2010;**35**:72-115.

Georgiou 2001

Georgiou D, Chen Y, Appadoo S, Belardinelli R, Greene R, Parides MK, et al. Cost-effectiveness analysis of long-term moderate exercise training in chronic heart failure. *American Journal of Cardiology* 2001;**87**:984-8.

Go 2014

Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Baha MJ, et al. Heart disease and stroke statistics - 2014 update: a report from the American Heart Association. *Circulation* 2014;**129**:e28-e292.

GRADEpro 2008 [Computer program]

GRADE Working Group GRADEprofiler. Version 3.2 for Windows. Brozek J, Oxman A, Schünemann H. GRADE Working Group, 2008.

Gravely-Witte 2007

Gravely-Witte S, De Gucht V, Heiser W, Grace SL, Van Elderen T. The impact of angina and cardiac history on health-related quality of life and depression in coronary heart disease patients. *Chronic Illness* 2007;**3**:66-76.

Hambrecht 1998

Hambrecht R, Fiehn E, Weigl C, Gielen S, Hamann C, Kaiser R, et al. Regular physical exercise corrects endothelial dysfunction and improves exercise capacity in patients with chronic heart failure. *Circulation* 1998;**98**(24):2709-15.

Hambrecht 2000

Hambrecht R, Gielen S, Linke A, Fiehn E, Yu J, Walther C, et al. Effects of exercise training on left ventricular function and peripheral resistance in patients with chronic heart failure: a randomized trial. *JAMA* 2000;**283**:3095-101.

Haykowsky 2007

Haykowsky MJ, Liang Y, Pechter D, Jones LW, McAlister FA, Clark AM. A meta-analysis of the effect of exercise training on left ventricular remodeling in heart failure patients: the benefit depends on the type of training performed. *Journal of the American College of Cardiology* 2007;**49**:2329-36.

Higgins 2011

Higgins JPT. Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Hobbs 2007

Hobbs FD, Roalfe AK, Davis RC, Davies MK, Hare R, Midlands Research Practices Consortium. Prognosis of all-cause heart failure and borderline left ventricular systolic dysfunction: 5 year mortality follow-up of the Echocardiographic Heart of England Screening Study (ECHOES). *European Heart Journal* 2007;**28**:1128-34.

Hoffmann 2014

Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;**348**:g1687.

Hwang 2009

Hwang R, Marwick T. Efficacy of home-based exercise programmes for people with chronic heart failure: a meta-analysis. *European Journal of Cardiovascular Prevention and Rehabilitation* 2009;**16**:527-35.

Jolliffe 2001

Jolliffe JA, Rees K, Taylor RS, Thompson D, Oldridge N, Ebrahim S. Exercise-based rehabilitation for coronary heart

disease. *Cochrane Database of Systematic Reviews* 2001, Issue 1. Art. No: CD001800. [DOI: [10.1002/14651858.CD001800](https://doi.org/10.1002/14651858.CD001800)]

Jolly 1998

Jolly K, Bradley F, Sharp S, Smith H, Mant D. Follow-up care in general practice of patients with myocardial infarction or angina pectoris: initial results of the SHIP trial. Southampton Heart Integrated Care Project. *Family Practice* 1998;**15**:548-55.

Jolly 2007

Jolly K, Taylor R, Lip GY, Greenfield S, Raftery J, Mant J, et al. The Birmingham Rehabilitation Uptake Maximisation Study (BRUM). Home-based compared with hospital-based cardiac rehabilitation in a multi-ethnic population: cost-effectiveness and patient adherence. *Health Technology Assessment* 2007;**11**:1-118.

Koongstvedt 2001

Koongstvedt PR. *The Managed Health Care Handbook*. 4th edition. New York: Aspen Publishers, 2001.

Kostis 1997

Kostis JB, Davis BR, Cutler J, Grimm RH Jr, Berge KG, Cohen JD, et al. Prevention of heart failure by antihypertensive drug treatment in older persons with isolated systolic hypertension. SHEP Cooperative Research Group. *JAMA* 1997;**278**:212-6.

Kung 2010

Kung J, Chiappelli F, Cajulis OO, Avezova R, Kossan G, Chew L, et al. From systematic reviews to clinical recommendations for evidence-based health care: validation of Revised Assessment of Multiple Systematic Reviews (R-AMSTAR) for grading of clinical relevance. *Open Dentistry Journal* 2010;**4**:84-91.

Lawler 2011

Lawler PR, Filion KB, Eisenberg MJ. Efficacy of exercise-based cardiac rehabilitation post-myocardial infarction: a systematic review and meta-analysis of randomized controlled trials. *American Heart Journal* 2011;**162**:571-84.

Lisspers 1999

Lisspers J, Sundin O, Hofman-Bang C, Nordlander R, Nygren A, Ryden L, et al. Behavioral effects of a comprehensive, multifactorial program for lifestyle change after percutaneous transluminal coronary angioplasty: a prospective, randomized controlled study. *Journal of Psychosomatic Research* 1999;**46**:143-54.

Mathers 2008

Mathers C, Fat D, Boerma JT. *The global burden of disease: 2004 update*. World Health Organization, 2008.

McKelvie 2013

McKelvie R S, Moe G W, Ezekowitz J A, Heckman G A, Costigan J, Ducharme A, et al. The 2012 Canadian Cardiovascular Society heart failure management guidelines update: focus on acute and chronic heart failure. *Can J Cardiol* 2013;**29**:168-81.

McMurray 2005

McMurray JJ, Pfeffer MA. Heart failure. *Lancet* 2005;**365**:1877-89.

McMurray 2012

McMurray John JV, Adamopoulos Stamatias, Anker Stefan D, Auricchio Angelo, Böhm Michael, Dickstein Kenneth, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012 The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *European heart journal* 2012;**33**:1787-847.

Mechta-Nielsen 2014

Mechta-Nielsen. Exercise-based cardiac rehabilitation for adult patients with ICD [personal communication]. Email to: L Anderson 9 May 2014.

Mehta 2009

Mehta PA, Dubrey SW, McIntyre HF, Walker DM, Hardman SM, Sutton GC, et al. Improving survival in the 6 months after diagnosis of heart failure in the past decade: population-based data from the UK. *Heart* 2009;**95**:1851-6.

Miller 1984

Miller NH, Haskell WL, Berra K, DeBusk RF. Home versus group exercise training for increasing functional capacity after myocardial infarction. *Circulation* 1984;**70**:645-9.

Mills 2013

Mills EJ, Thorlund K, Ioannidis JP. Demystifying trial networks and network meta-analysis. *BMJ* 2013;**346**:f2914.

Moore 2003

Moore SM, Dolansky MA, Ruland CM, Pashkow FJ, Blackburn GG. Predictors of women's exercise maintenance after cardiac rehabilitation. *Journal of Cardiopulmonary Rehabilitation* 2003;**23**:40-9.

NICE 2013

NICE. Clinical guideline 172. MI – secondary prevention. Secondary prevention in primary and secondary care for patients following a myocardial infarction. www.nice.org.uk/guidance/cg172 2013.

NICE 2013b

NICE. Commissioning guides 40. Cardiac rehabilitation services. www.nice.org.uk/guidance/cm40 2013.

O'Connor 1989

O'Connor GT, Buring JE, Yusuf S, Goldhaber SZ, Olmstead EM, Paffenbarger RS Jr, et al. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. *Circulation* 1989;**80**:234-44.

Oldridge 1988

Oldridge NB, Guyatt GH, Fischer ME, Rimm AA. Cardiac rehabilitation after myocardial infarction. Combined experience of randomized clinical trials. *JAMA* 1988;**260**:945-50.

Oldridge 2012

Oldridge N. Exercise-based cardiac rehabilitation in patients with coronary heart disease: meta-analysis outcomes revisited. *Future of Cardiology* 2012;**8**:729-51.

Perk 2012

Perk J, De Backer G, Gohlke H, Graham I, Reiner Z, Verschuren M, et al. European Guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts). *European Heart Journal* 2012;**33**(13):1635-701.

Petticrew 2013

Petticrew M, Anderson L, Elder R, Grimshaw J, Hopkins D, Hahn R, et al. Complex interventions and their implications for systematic reviews: a pragmatic approach. *Journal of Clinical Epidemiology* 2013;**66**:1209-14.

Piepoli 2004

Piepoli MF, Davos C, Francis DP, Coats AJ, ExTra Match Collaborative. Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ* 2004;**328**:189.

PRECOR 1991

PRECOR, Leizoroviez A, Saint-Pierre A, Vasselon C, Boissel JP. Comparison of a rehabilitation programme, a counselling programme and usual care after an acute myocardial infarction: results of a long-term randomized trial. *European Heart Journal* 1991;**12**:612-6.

Racca 2010

Racca V, Spezzaferrri R, Modica M, Mazzini P, Jonsdottir J, De Maria R, et al. Functioning and disability in ischaemic heart disease. *Disability and Rehabilitation* 2010;**32 Suppl 1**:S42-9.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Risom 2014

Risom SS, Zwisler A-D, Johansen PP, Sibilitz KL, Lindschou J, Taylor RS, et al. Exercise-based cardiac rehabilitation for adults with atrial fibrillation. *Cochrane Database of Systematic Reviews* 2014, Issue 7. Art. No: CD011197. [DOI: [10.1002/14651858.CD011197](https://doi.org/10.1002/14651858.CD011197)]

Shea 2009

Shea BJ, Hamel C, Wells GA, Bouter LM, Kristjansson E, Grimshaw J, et al. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *Journal of Clinical Epidemiology* 2009;**62**:1013-20.

Sibilitz 2013

Sibilitz KL, Berg SK, Tang LH, Risom SS, Gluud C, Lindschou J, et al. Exercise-based cardiac rehabilitation for adults after heart valve surgery. *Cochrane Database of Systematic Reviews* 2013, Issue 12. Art. No: CD010876. [DOI: [10.1002/14651858.CD010876](https://doi.org/10.1002/14651858.CD010876)]

Stern 1983

Stern MJ, Gorman PA, Kaslow L. The group counselling v exercise therapy study. A controlled intervention with subjects

following myocardial infarction. *Archives of Internal Medicine* 1983;**143**:1719-25.

Task Force Members 2013

Task Force Members, Montalescot G, Sechtem U, Andreotti F, Arden C, Budaj A, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *European Heart Journal* 2013;**34**(38):2949-3003.

Taylor 2004

Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *American Journal of Medicine* 2004;**116**:682-92.

Taylor 2006

Taylor RS, Unal B, Critchley JA, Capewell S. Mortality reductions in patients receiving exercise-based cardiac rehabilitation: how much can be attributed to cardiovascular risk factor improvements? *European Journal of Cardiovascular Prevention and Rehabilitation* 2006;**13**:369-74.

Taylor 2012

Taylor RS, Cochrane Review Authors. The RAMIT trial: its results in the context of 2012 Cochrane review. *Heart* 2012;**98**:672-3.

Tierney 2011

Tierney S, Mamas M, Skelton D, Woods S, Rutter MK, Gibson M, et al. What can we learn from patients with heart failure about

exercise adherence? A systematic review of qualitative papers. *Health Psychology* 2011;**30**:401-10.

Weir 2012

Weir MC, Grimshaw, JM, Mayhew A, Fergusson D. Decisions about lumping vs. splitting of the scope of systematic reviews of complex interventions are not well justified: a case study in systematic reviews of health care professional reminders. *Journal of Clinical Epidemiology* 2012;**65**:756-63.

West 2012

West RR, Jones DA, Henderson AH. Rehabilitation After Myocardial Infarction Trial (RAMIT): multi-centre randomised controlled trial of comprehensive cardiac rehabilitation in patients following acute myocardial infarction. *Heart* 2012;**98**:637-44.

WHO 2014

World Health Organization (WHO). Global health observatory data repository. apps.who.int/gho/data/node.main (accessed 7 December 2014).

Yancy 2013

Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE, Drazner MH, et al. 2013 ACCF/AHA Guideline for the Management of Heart Failure. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology* 2013;**62**:e147-239.

ADDITIONAL TABLES

Table 1. Summary of included Cochrane review characteristics

Review short title (reference)	Exercise for CHD (Heran 2011)	Exercise for HF (Taylor 2014b)	Psychological for CHD (Whalley 2011)	Education for CHD (Brown 2011)	Home vs. centre (Taylor 2014a)	Uptake and adherence (Karmali 2014)
Main objective	To determine the effectiveness of exercise-based CR (exercise training alone or in combination with psychosocial or educational interventions) on mortality, morbidity and HRQoL of people with CHD	To determine the effectiveness of exercise-based interventions compared with usual medical care by focusing on mortality, hospital admission rate, morbidity and HRQoL in	To determine the independent effects of psychological interventions in people with CHD	To assess the effects of patient education on mortality, morbidity, HRQoL and healthcare costs in people with CHD	To determine the effectiveness of home-based CR programmes compared with supervised centre-based CR on mortality and morbidity, HRQoL and modifiable cardiac risk factors in people with CHD	To determine the harms and benefits of interventions to increase patient uptake of, and adherence to, CR

Table 1. Summary of included Cochrane review characteristics (Continued)

	people with HF					
Search time frame	November 2000 to December 2009	2008 to March 2013	2001 to January 2009	1990 to August 2010	2008 to November 2012	2008 to January 2013
Study design	RCTs (follow-up \geq 6 months)	RCTs (follow-up \geq 6 months)	RCTs (no minimum follow-up)	RCTs (follow-up \geq 6 months)	RCTs (no minimum follow-up)	RCTs (no minimum follow-up)
Population	<i>Inclusion</i> Post-MI Post revascularisation CHD defined by angiography <i>Exclusion</i> Heart valve surgery HF Heart transplantation CRT or ICD implant	<i>Inclusion</i> HF Previous CR <i>Exclusion</i> None	<i>Inclusion</i> Post-MI Post revascularisation Angina CHD defined by angiography <i>Exclusion</i> None	<i>Inclusion</i> Post-MI Post revascularisation Angina CHD defined by angiography	<i>Inclusion</i> Post-MI Post revascularisation Angina HF <i>Exclusion</i> Heart transplantation CRT or CD implant Previous CR	<i>Inclusion</i> Post-MI Post revascularisation Angina HF CHD <i>Exclusion</i> Heart transplantation CRT or ICD implant
Intervention	Exercise training with or without the addition of psychosocial or educational interventions (or both)	Exercise training with or without the addition of psychosocial or educational interventions (or both)	Psychological interventions delivered by healthcare workers with specific training in psychological techniques	Patient education interventions involving direct contact with a health professional and including structured knowledge transfer about CHD	CR programmes delivered in a home-based setting	CR plus any intervention with the specific aim of increasing patient uptake of, or adherence to, CR or any of its component parts
Comparator	No exercise training control that could include psychological, educational interventions, standard medical care or a combination	No exercise training control that could include psychological, educational interventions, standard medical care or a combination	No psychological intervention control that could include exercise interventions or standard medical care	No education intervention control that could include exercise interventions or standard medical care	CR programmes delivered in a centre-based setting	CR programmes without the intervention
Outcomes	<ul style="list-style-type: none"> Mortality (total, CV, non-CV) MI (total, fatal, non-fatal) 	<ul style="list-style-type: none"> Mortality (total, HF and sud- 	<ul style="list-style-type: none"> Mortality (total and CV) Morbidity (non-fatal MI) 	<ul style="list-style-type: none"> Mortality (total, CV and non-CV) 	<ul style="list-style-type: none"> Mortality (total and CV) Morbidity (reinfarction, revascularisa- 	Uptake of, or adherence to, CR (primary)

Table 1. Summary of included Cochrane review characteristics (Continued)

	<ul style="list-style-type: none"> Revascularisations (total, CABG, PTCA, re-stenting) Hospitalisations (total, CV, other) HRQoL Economic (costs and cost-effectiveness) 	<ul style="list-style-type: none"> den death Hospitalisation (total, HF) HRQoL Economic (costs and cost-effectiveness) 	<ul style="list-style-type: none"> Revascularisation (CABG and PTCA) Psychological well-being anxiety, depression, stress and Type A Behaviour/hostility HRQoL 	<ul style="list-style-type: none"> Total CV events MI (fatal or non-fatal, or both) Other fatal or non-fatal (or both) CV events Revascularisations (CABG, PTCA with or without stenting) Hospitalisations (cardiac-related) HRQoL Withdrawals/drop-outs Economic (healthcare costs and cost-effectiveness) 	<ul style="list-style-type: none"> tion, cardiac-associated hospitalisation) Exercise capacity Risk factors (smoking behaviour, blood lipid levels, blood pressure) HRQoL Adverse events (withdrawal from the exercise programme) Adherence to rehabilitation Economic (health service use, costs and cost-effectiveness) 	<ul style="list-style-type: none"> Mortality (total) Morbidity Risk factors (smoking behaviour, blood lipid levels, blood pressure) HRQoL Economic (healthcare costs and cost-effectiveness) Any beneficial or adverse events
Funding source	NIHR, UK Cochrane Collaboration Programme Grant, UK	None specified	Department of Social Medicine, University of Bristol, UK Health Services Research Focus, University of Wales College of Medicine, UK British Heart Foundation, UK ESCR, UK NIHR, UK Cochrane Collaboration Heart Programme Grant, UK	NIHR, UK Cochrane Collaboration Programme Grant, UK	NIHR Cochrane Heart Programme grant, UK Transparency of the National Health System Drug Reimbursement Decisions, Poland, EU	NIHR programme grant, UK
Authors' declarations of interest	Authors were authors of the original Cochrane review. RST was a co-investigator on a number of CR RCTs	-	None declared	None declared	RST was a co-author of the original Cochrane review and was a co-investigator on a number of CR RCTs	None declared

CABG: coronary artery bypass graft; CAD: coronary artery disease; CHD: coronary heart disease; CR: cardiac rehabilitation; CRT: cardiac resynchronisation therapy; CV: cardiovascular; ESCR: Economic and Social Research Council; HF: heart failure; HRQoL: health-related

quality of life; ICD: implantable cardioverter defibrillator; MI: myocardial infarction; NIHR: National Institute of Health Research; PTCA: percutaneous transluminal coronary angioplasty; RCT: randomised controlled trial.

Table 2. Summary of characteristics of included RCTs

Review short title (reference)	Exercise for CHD (Heran 2011)	Exercise for HF (Taylor 2014b)	Psycho- logical for CHD (Whalley 2011)	Education for CHD (Brown 2011)	Home vs. centre (Taylor 2014a)	Uptake and adherence (Karmali 2014)
RCTs (participants)						
Number	47 RCTs (10,794)	33 RCTs (4740)	24 RCTs (9296)	13 RCTs (68,556)	17 RCTs (2172)	18 RCTs (2505)
Nature of intervention*						
Exercise only	17	21	0	0	6	Interventions aimed at increasing patient up-take of CR (10 RCTs)
Psychological only	0	0	14	0	0	
Education only	0	0	0	13	0	Interventions designed to increase adherence to exercise (7 RCTs) or supervised CR (1 RCT)
> 1 intervention	29*	12	10 (psycho-logical and education)	0	11	
Sample size						
Median (range)	142 (28 to 2304)	54 (19 to 2331)	133 (44 to 2481)	288 (87 to 46,606)	104 (20 to 525)	110 (16 to 597)
Intervention duration [months]						
Median (range) months	3 (1 to 30)	6 (1 to 120)	NR	6 (1 to 30)	3 (1.5 to 6)	NR
Publication year (number of RCTs)						
1970-1979	2	0	2	0	0	0
1990-1999	11	0	4	0	1	2
1990-1999	20	5	8	4	2	3
2000-2009	14	20	10	9	11	8
2010+	0	8	0	0	3	5
% male						
Median (range)	88 (0 to 100)	80 (36 to 100)	84 (0 to 100)	60 (0 to 100)	80 (60 to 100)	84 (0 to 100)

Table 2. Summary of characteristics of included RCTs (Continued)

% white						
Median (range)	NR	85 (60 to 100) from 8 RCTs	NR	86 (55 to 97) from 6 RCTs	80 from 1 RCT	79 (43 to 95) from 6 RCTs
Age (years)						
Median (range)	55 (49 to 70)	60 (51 to 81)	57 (51 to 62)	62 (51 to 73)	60 (52 to 69)	62 (51 to 77)
Indication (number of RCTs)						
MI only	28	0	10	2	4	4
Angina only	1	0	1	1	0	0
Revascularisation only	1	0	4	2	4	0
MI or revascularisation (or both)	4	0	4	1	5	3
MI or angina	4	0	2	0	0	3
Mixed CHD	9	0	2	4	0	7
HF	0	33	0	3 CHD or HF	3	1
Arrhythmia	0	0	1	0	1	0
Study location (number of RCTs (%))						
Europe	20 (43)	20 (64)	11 (46)	7 (54)	10 (58)	6 (33)
North America	3 (6)	11 (30)	11 (46)	6 (46)	5 (29)	11 (61)
Asia/Australia	7 (15)	1 (3)	2 (8)	0	1 (6)	1 (6)
Other	-	1 (3)	0	0	1 (6)	0
NR	17 (36)	0	0	0	0	0
Single centre						
Number of RCTs (%)	23 (49)	30 (91)	8 (33)	4 (31)	15 (88)	10/16 (63)**
Follow-up duration [months]						
Median (range)	24 (6 to 120)	6 (6 to 120)	NR	18 (6 to 60)	6 (2 to 72)	3 (1.5 to 12)

CHD: coronary heart disease; HF: heart failure; MI: myocardial infarction; NR: not reported; RCT: randomised controlled trial.

* 1 RCT randomly assigned to exercise-only or comprehensive intervention.

** 2 studies were unavailable to us as they were unpublished degree dissertations.

Table 3. R-AMSTAR assessment of included systematic reviews

Review short title (reference)	Exercise for CHD (Heran 2011)	Exercise for HF (Taylor 2014b)	Psycho- logical for CHD (Whalley 2011)	Education for CHD (Brown 2011)	Home vs. centre (Taylor 2014a)	Uptake and adher- ence (Karmali 2014)
1. Was an 'a priori' design provided?						
(A) 'a priori' design	Yes	Yes	Yes	Yes	Yes	Yes
(B) Statement of inclusion criteria	Yes	Yes	Yes	Yes	Yes	Yes
(C) PICO/PIPO research question (popula- tion, intervention, comparison, prediction, outcome)	Yes	Yes	Yes	Yes	Yes	Yes
Score	4	4	4	4	4	4
2. Was there duplicate study selection and data extraction?						
(A) There should be at least 2 independent data extractors as stated or implied	Yes	*Yes	*Yes	*Yes	*Yes	Yes
(B) Statement of recognition or awareness of consensus procedure for disagreements	Yes	Yes	Yes	Yes	Yes	Yes
(C) Disagreements among extractors re- solved properly as stated or implied	Yes	Yes	Yes	Yes	Yes	Yes
Score	4	4	4	4	4	Yes
3. Was a comprehensive literature search performed?						
(A) At least 2 electronic sources should be searched	Yes	Yes	Yes	Yes	Yes	Yes
(B) The report must include years and databases used (e.g. CENTRAL, MEDLINE, EMBASE)	Yes	Yes	Yes	Yes	Yes	Yes
(C) Key words or MESH terms (or both) must be stated AND where feasible the search strategy outline should be provid- ed such that one can trace the filtering process of the included articles	Yes	Yes	Yes	Yes	Yes	Yes
(D) In addition to the electronic databases (PubMed, MEDLINE, EMBASE), all search- es should be supplemented by consulting current contents, reviews, textbooks, spe- cialised registers, or experts in the particu- lar field of study, and by reviewing the ref- erences in the studies found	No	Yes	Yes	Yes	Yes	No
(E) Journals were "hand-searched" or "manual searched" (i.e. identifying highly relevant journals and conducting a manu-	No	No	No	No	No	No

Table 3. R-AMSTAR assessment of included systematic reviews (Continued)

al, page-by-page search of their entire contents looking for potentially eligible studies)

Score	3	4	4	4	4	3
4. Was the status of publication (i.e. grey literature) used as an inclusion criterion?						
(A) The authors should state that they searched for reports regardless of their publication type	*No	*No	No	*No	Yes	*No
(B) The authors should state whether or not they excluded any reports (from the systematic review), based on their publication status, language, etc.	Yes	Yes	Yes	Yes	Yes	Yes
(C) "Non-English papers were translated" or readers sufficiently trained in foreign language	Yes	No	Yes	Yes	No	No
(D) No language restriction or recognition of non-English articles	Yes	Yes	Yes	Yes	Yes	Yes
Score	4	3	4	4	4	3
5. Was a list of studies (included and excluded) provided?						
(A) Table/list/figure of included studies, a reference list does not suffice	Yes	Yes	Yes	Yes	Yes	Yes
(B) Table/list/figure of excluded studies, either in the article or in a supplemental source (i.e. online). (Excluded studies refers to those studies seriously considered on the basis of title and/or abstract, but rejected after reading the body of the text)	Yes	Yes	Yes	Yes	Yes	Yes
(C) Author satisfactorily/sufficiently stated the reason for exclusion of the seriously considered studies	Yes	Yes	Yes	Yes	Yes	Yes
(D) Reader was able to retrace the included and the excluded studies anywhere in the article bibliography, reference or supplemental source	Yes	Yes	Yes	Yes	Yes	Yes
Score	4	4	4	4	4	4
6. Were the characteristics of the included studies provided?						
(A) In an aggregated form such as a table, data from the original studies should be provided on the participants, interventions AND outcomes	Yes	Yes	Yes	Yes	Yes	Yes
(B) Provide the ranges of relevant characteristics in the studies analysed (e.g. age, race, sex, relevant socioeconomic data,	Yes	Yes	Yes	Yes	Yes	Yes

Table 3. R-AMSTAR assessment of included systematic reviews (Continued)

disease status, duration, severity or other diseases should be reported)						
(C) The information provided appears to be complete and accurate (i.e. there was a tolerable range of subjectivity here. Is the reader left wondering? If so, state the needed information and the reasoning)	Yes	Yes	Yes	Yes	Yes	Yes
Score	4	4	4	4	4	4
7. Was the scientific quality of the included studies assessed and documented?						
(A) 'A priori' methods of assessment should be provided (e.g. for effectiveness studies if the author(s) chose to include only randomised, double-blind, placebo-controlled studies, or allocation concealment as inclusion criteria); for other types of studies alternative items will be relevant	Yes	Yes	Yes	Yes	Yes	Yes
(B) The scientific quality of the included studies appeared to be meaningful	Yes	Yes	Yes	Yes	Yes	Yes
(C) Discussion/recognition/awareness of level of evidence	Yes	Yes	Yes	Yes	Yes	Yes
(D) Quality of evidence should be rated/ranked based on characterised instruments. (Characterised instrument is a created instrument that ranks the level of evidence, e.g. GRADE (Grading of Recommendations Assessment, Development and Evaluation))	No	No	No	No	No	No
Score	3	3	3	3	3	3
8. Was the scientific quality of the included studies used appropriately in formulating conclusions?						
(A) The results of the methodological rigor and scientific quality should be considered in the analysis and the conclusions of the review	Yes	Yes	Yes	Yes	Yes	Yes
(B) The results of the methodological rigor and scientific quality were explicitly stated in formulating recommendations	No	No	No	No	No	Yes
(C) To have conclusions integrated/drives towards a clinical consensus statement	Yes	Yes	Yes	Yes	Yes	Yes
(D) This clinical consensus statement drives towards revision or confirmation of clinical practice guidelines	No	No	No	Yes	No	No
Score	2	2	2	3	2	3
9. Were the methods used to combine the findings of studies appropriate?						

Table 3. R-AMSTAR assessment of included systematic reviews (Continued)

(A) Statement of criteria that were used to decide that the studies analysed were similar enough to be pooled?	Yes	Yes	No	Yes	Yes	Yes
(B) For the pooled results, a test should be done to ensure the studies were combinable, to assess their homogeneity (i.e. Chi ² test for homogeneity, I ² statistic)	Yes	Yes	Yes	Yes	Yes	NA
(C) Is there a recognition of heterogeneity or lack of thereof	Yes	Yes	Yes	Yes	Yes	Yes
(D) If heterogeneity exists a "random-effects model" should be used or the rationale (i.e. clinical appropriateness) of combining should be taken into consideration (i.e. is it sensible to combine?), or stated explicitly (or both)	Yes	Yes	Yes	Yes	Yes	NA
(E) If homogeneity exists, author should state a rationale or a statistical test	Yes	Yes	NA	NA	Yes	NA
Score	4	4	3	4	4	2
10. Was the likelihood of publication bias (a.k.a. "file drawer" effect) assessed?						
(A) Recognition of publication bias or file-drawer effect	Yes	Yes	Yes	Yes	Yes	Yes
(B) An assessment of publication bias should include graphical aids (e.g. funnel plot, other available tests)	Yes	Yes	Yes	Yes	Yes	No
(C) Statistical tests (e.g. Egger regression test)	Yes	Yes	Yes	Yes	Yes	No
Score	4	4	4	4	4	2
11. Was the conflict of interest stated?						
(A) Statement of sources of support	Yes	Yes	Yes	Yes	Yes	Yes
(B) No conflict of interest. This is subjective and may require some deduction or searching	Yes	Yes	Yes	Yes	Yes	Yes
(C) An awareness/statement of support or conflict of interest in the primary inclusion studies	No	No	No	No	No	No
Score	3	3	3	3	3	3
Total score (n/44)	39	39	39	41	40	35

CHD: coronary heart disease; HF: heart failure.

* Studies were screened independently by 2 review authors. Data were extracted by 1 review author and checked by a second review author.

** While the authors did not explicitly state that they searched for reports regardless of publication type, it was clear from the included studies or text (or both) that a search of grey literature was conducted.

Table 4. Risk of bias of included randomised controlled trials

Review short title (reference)	Exercise for CHD (Heran 2011)	Exercise for HF (Taylor 2014b)	Psychologi- cal for CHD (Whalley 2011)	Education for CHD (Brown 2011)	Home vs. centre (Taylor 2014a)	Uptake and adherence (Karmali 2014)	Total
Number of RCTs with low risk of bias (%)							
Random sequence generation	8 (17)	10 (30)	7 (29)	9 (69)	4 (24)	9 (50)	47 (31)
Allocation concealment	7 (15)	6 (18)	7 (29)	7 (54)	7 (41)	8 (44)	41 (27)
Groups balanced at baseline	^a 27 (57)	32 (97)	^a 10 (42)	12 (92)	14 (82)	ab ⁹ (56)	103 (68)
Outcome blinding	4 (9)	11 (33)	5 (21)	4 (31)	7 (41)	5 (28)	36 (24)
Selective reporting	0 (0)	31 (94)	16 (67)	12 (92)	16 (94)	15 (83)	90 (59)
Loss to follow-up < 20%	33 (70)	29 (88)	13 (54)	10 (77)	11 (65)	4 (22)	99 (65)
Intention-to-treat analysis	^a 19 (40)	29 (88)	22 (92)	11 (85)	14 (82)	ab ⁷ (44)	101 (66)
Groups received same treatment apart from intervention*	^a 21 (45)	21 (64)	^a 16 (67)	11 (85)	15 (88)	ab ¹⁵ (94)	100 (66)

CHD: coronary heart disease; HF: heart failure; RCT: randomised controlled trial.

^a Risk of bias was not reported within the review, but was assessed by the authors of this overview.

^b Denominator = 16 as 2 studies were unavailable to us as they were unpublished degree dissertations.

Table 5. Exercise-based cardiac rehabilitation for coronary heart disease

Exercise-based cardiac rehabilitation for coronary heart disease						
Patient or population: people with CHD						
Settings:						
Intervention: exercise-based CR						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Exercise-based CR				
Total mortality Follow-up: 6-12 months	Study population		RR 0.82 (0.67 to 1.01)	6000 (19 studies)	⊕⊕⊕⊖ low ^{1,2}	-
	65 per 1000	53 per 1000 (43 to 65)				
	Moderate					
-	-	-	-	-	-	-
Total mortality Follow-up: 12-120 months	Study population		RR 0.87 (0.75 to 0.99)	5790 (16 studies)	⊕⊕⊕⊖ moderate ¹	-
	126 per 1000	109 per 1000 (94 to 125)				
	Moderate					
-	-	-	-	-	-	-
Cardiovascular mortality Follow-up: 6-12 months	Study population		RR 0.93 (0.71 to 1.21)	4130 (9 studies)	⊕⊕⊕⊖ low ^{1,2}	-
	51 per 1000	48 per 1000 (36 to 62)				
	Moderate					
-	-	-	-	-	-	-
Cardiovascular mortality Follow-up: 12-120 months	Study population		RR 0.74 (0.63 to 0.87)	4757 (12 studies)	⊕⊕⊕⊖ moderate ¹	-
	129 per 1000	96 per 1000 (81 to 112)				
	Moderate					
-	-	-	-	-	-	-
Hospitalisations Follow-up: 6-12 months	Study population		RR 0.69 (0.51 to 0.93)	463 (4 studies)	⊕⊕⊕⊖ moderate ¹	-
	324 per 1000	224 per 1000 (165 to 302)				
	Moderate					
-	-	-	-	-	-	-

Table 5. Exercise-based cardiac rehabilitation for coronary heart disease (Continued)

Hospitalisations Follow-up: 12-48 months	Study population		RR 0.98 (0.87 to 1.11)	2009 (7 studies)	⊕⊕⊕⊖ low ^{1,3}	-
	342 per 1000	335 per 1000 (297 to 379)				
	Moderate					
-	-					
MI Follow-up: 6-12 months	Study population		RR 0.92 (0.7 to 1.22)	4216 (12 studies)	⊕⊕⊕⊖ very low ^{1,2,4}	-
	45 per 1000	41 per 1000 (32 to 55)				
	Moderate					
-	-					
MI Follow-up: 12-120 months	Study population		RR 0.97 (0.82 to 1.15)	5682 (16 studies)	⊕⊕⊕⊖ low ^{1,4}	-
	89 per 1000	87 per 1000 (73 to 103)				
	Moderate					
-	-					
CABG Follow-up: 6-12 months	Study population		RR 0.91 (0.67 to 1.24)	2312 (14 studies)	⊕⊕⊕⊖ low ^{1,2}	-
	67 per 1000	61 per 1000 (45 to 83)				
	Moderate					
-	-					
CABG Follow-up: 12-120 months	Study population		RR 0.93 (0.68 to 1.27)	2189 (9 studies)	⊕⊕⊕⊖ low ^{1,2}	-
	69 per 1000	64 per 1000 (47 to 88)				
	Moderate					
-	-					
PTCA Follow-up: 6-12 months	Study population		RR 1.02 (0.69 to 1.5)	1328 (7 studies)	⊕⊕⊕⊖ low ^{1,2}	-
	69 per 1000	71 per 1000 (48 to 104)				
	Moderate					
-	-					

Table 5. Exercise-based cardiac rehabilitation for coronary heart disease (Continued)

PTCA	Study population		RR 0.89	1322	⊕⊕○○	-
Follow-up: 12-48 months			(0.66 to 1.19)	(6 studies)	low ^{1,2}	
	124 per 1000	110 per 1000 (82 to 147)				
	Moderate					
	-	-				

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CABG: coronary artery bypass graft; **CHD:** coronary heart disease; **CI:** confidence interval; **CR:** cardiac rehabilitation; **MI:** myocardial infarction; **PTCA:** percutaneous transluminal coronary angioplasty; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Random sequence generation and allocation concealment were poorly described; bias likely.

² The 95% CIs include both no effect and appreciable benefit or harm (i.e. RR < 0.75 or > 1.25).

³ Moderate heterogeneity ($I^2 > 50\%$).

⁴ Funnel plots or Egger test (or both) suggest evidence of asymmetry.

Table 6. Exercise-based cardiac rehabilitation for heart disease
Exercise-based cardiac rehabilitation for heart failure

Patient or population: people with HF

Settings:

Intervention: exercise-based CR

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Exercise-based CR				
Total mortality	Study population		RR 0.93	1871	⊕⊕○○	-
Follow-up: 6-12 months			(0.69 to 1.27)	(25 studies)	low ^{1,2}	
	75 per 1000	70 per 1000 (52 to 96)				
	Moderate					
	-	-				
Total mortality	Study population		RR 0.88	2845	⊕⊕○○	-
				(6 studies)	low ^{1,2}	

Table 6. Exercise-based cardiac rehabilitation for heart disease (Continued)

Follow-up: 12-120 months	196 per 1000	173 per 1000 (147 to 200)	(0.75 to 1.02)			
	Moderate					
	-	-				
Hospitalisations Follow-up: 6-12 months	227 per 1000	170 per 1000 (141 to 209)	RR 0.75 (0.62 to 0.92)	1328 (15 studies)	⊕⊕⊕⊖ moderate ¹	-
	Moderate					
	-	-				
Hospitalisations Follow-up: 12-74 months	604 per 1000	556 per 1000 (399 to 779)	RR 0.92 (0.66 to 1.29)	2722 (5 studies)	⊕⊖⊖⊖ very low ^{1,2,3}	-
	Moderate					
	-	-				
Hospitalisations (HF-specific admissions) Follow-up: 12-120 months	182 per 1000	111 per 1000 (84 to 145)	RR 0.61 (0.46 to 0.8)	1036 (12 studies)	⊕⊕⊕⊖ moderate ^{1,2}	-
	Moderate					
	-	-				
HRQoL MLWHF score Follow-up: 6-12 months	-	The mean HRQoL in the intervention groups was 5.8 lower (9.21 to 2.44 lower)	-	1270 (13 studies)	⊕⊖⊖⊖ very low ^{1,3,4}	-
HRQoL All HRQoL measures Follow-up: 12-120 months	-	The mean HRQoL in the intervention groups was 0.46 lower (0.66 to 0.26 lower)	-	3240 (13 studies)	⊕⊖⊖⊖ very low ^{1,2,3,4}	-
HRQoL MLWHF Follow-up: 6-120 months	-	The mean HRQoL in the intervention groups was 9.49 lower (17.48 to 1.5 lower)	-	329 (20 studies)	⊕⊖⊖⊖ very low ^{1,2,3,4}	-

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **CR:** cardiac rehabilitation; **HF:** heart failure; **HRQoL:** health-related quality of life; **MLWHF:** Minnesota Living with Heart Failure questionnaire; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Table 6. Exercise-based cardiac rehabilitation for heart disease (Continued)

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Random sequence generation and allocation concealment were poorly described; bias likely.

² The 95% CIs include both no effect and appreciable benefit or harm (i.e. RR < 0.75 or > 1.25).

³ Moderate heterogeneity ($I^2 > 50\%$).

⁴ Funnel plots or Egger test (or both) suggest evidence of asymmetry.

Table 7. Psychological-based interventions for coronary heart disease
Psychological-based interventions for coronary heart disease

Patient or population: people with CHD

Settings:

Intervention: psychological-based interventions

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Psychological-based interventions				
Total mortality Follow-up: 6-12 months	Study population		RR 0.89 (0.75 to 1.05)	6852 (17 studies)	⊕⊕⊕⊕ low ^{1,2}	-
	93 per 1000	83 per 1000 (70 to 98)				
	Moderate					
	-	-				
Cardiovascular mortality Follow-up: 6-15 months	Study population		RR 0.80 (0.64 to 1)	3893 (5 studies)	⊕⊕⊕⊕ low ^{1,2}	-
	85 per 1000	68 per 1000 (55 to 85)				
	Moderate					
	-	-				
MI (non-fatal) Follow-up: 6-15 months	Study population		RR 0.87 (0.67 to 1.13)	7534 (12 studies)	⊕⊕⊕⊕ low ^{1,2}	-
	83 per 1000	72 per 1000 (55 to 94)				
	Moderate					
	-	-				

Table 7. Psychological-based interventions for coronary heart disease (Continued)

Revascularisation (CABG and PTCA combined) Follow-up: 6-15 months	Study population		RR 0.95 (0.8 to 1.13)	6670 (12 studies)	⊕⊕⊕⊖ moderate ¹	-
	121 per 1000	115 per 1000 (97 to 137)				
	Moderate					
-	-	-	-	-	-	-

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CABG: coronary artery bypass graft; **CHD:** coronary heart disease; **CI:** confidence interval; **MI:** myocardial infarction; **PTCA:** percutaneous transluminal coronary angioplasty; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Random sequence generation and allocation concealment were poorly described; bias likely.

² The 95% CIs include both no effect and appreciable benefit or harm (i.e. RR < 0.75 or > 1.25).

Table 8. Education-based interventions for coronary heart disease
Education-based interventions for coronary heart disease

Patient or population: people with CHD

Settings:

Intervention: education-based interventions

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Education-based interventions				
Total mortality deaths Follow-up: median 18 months	Study population		RR 0.79 (0.55 to 1.13)	2330 (6 studies)	⊕⊕⊕⊖ moderate ¹	-
	96 per 1000	76 per 1000 (53 to 108)				
	Moderate					
-	-	-	-	-	-	-
Hospitalisations	Study population		RR 0.83 (0.65 to 1.07)	12,905 (4 studies)	⊕⊕⊕⊖ moderate ¹	-
	64 per 1000	53 per 1000 (41 to 68)				
	Moderate					
-	-	-	-	-	-	-

Table 8. Education-based interventions for coronary heart disease (Continued)

		Moderate			
		-	-		
MI	Study population			RR 0.63	209
				(0.26 to 1.48)	(2 studies)
	118 per 1000	74 per 1000			
		(31 to 174)		$\oplus\oplus\oplus\oplus$	very low ²
	Moderate				
		-	-		
CABG	Study population			RR 0.58	209
				(0.19 to 1.71)	(2 studies)
	78 per 1000	45 per 1000			
		(15 to 134)		$\oplus\oplus\oplus\oplus$	low ²
	Moderate				
		-	-		
All-cause withdrawal	Study population			RR 1.03	2862
				(0.83 to 1.27)	(8 studies)
	181 per 1000	186 per 1000			
		(150 to 230)		$\oplus\oplus\oplus\oplus$	moderate ¹
	Moderate				
		-	-		

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CABG: coronary artery bypass graft; **CHD:** coronary heart disease; **CI:** confidence interval; **MI:** myocardial infarction; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ The 95% CIs include both no effect and appreciable benefit or harm (i.e. RR < 0.75 or > 1.25).

² The 95% CIs include both no effect and substantial benefit or harm (i.e. RR < 0.50 or > 1.50).

Table 9. Home-based cardiac rehabilitation compared with centre-based cardiac rehabilitation for heart disease
Home-based cardiac rehabilitation compared with centre-based cardiac rehabilitation for heart disease

Patient or population: people with heart disease

Settings:

Intervention: home-based CR

Comparison: centre-based CR

Table 9. Home-based cardiac rehabilitation compared with centre-based cardiac rehabilitation for heart

disease Outcomes (Continued)	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Centre-based CR	Home-based CR				
Total mortality Follow-up: 3-12 months	Study population		RR 0.79 (0.43 to 1.47)	1166 (7 studies)	⊕⊕⊕⊖ low ^{1,2}	-
	27 per 1000	22 per 1000 (12 to 40)				
	Moderate					
	-	-				
All-cause withdrawal Follow-up: median 6 months	Study population		RR 1.04 (1.01 to 1.07)	1984 (18 studies)	⊕⊕⊕⊖ moderate ¹	-
	874 per 1000	909 per 1000 (883 to 936)				
	Moderate					
	-	-				

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **CR:** cardiac rehabilitation; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Random sequence generation and allocation concealment were poorly described; bias likely.

² The 95% CIs include both no effect and appreciable benefit or harm (i.e. RR < 0.75 or > 1.25).

Table 10. Summary of outcome results across Cochrane systematic reviews

Review short title (reference)	Exercise for CHD (Heran 2011)	Exercise for HF (Taylor 2014b)	Psychological for CHD (Whalley 2011)	Education for CHD (Brown 2011)	Home vs. centre (Taylor 2014a)	Uptake and adherence (Karmali 2014)
Total mortality	Follow-up < 12 months 19 RCTs (6000 participants),	Follow-up < 12 months 25 RCTs (1871 participants)	17 RCTs (6852 participants)	6 RCTs (2330 participants) RR 0.79; 95% CI 0.55 to 1.13	Follow-up < 12 months 7 RCTs (1166 participants)	3 RCTs (211 participants) 0/3 RCTs reported a significant dif-

Table 10. Summary of outcome results across Cochrane systematic reviews (Continued)

	RR 0.82; 95% CI 0.67 to 1.01 $I^2 = 0\%$	RR 0.93; 95% CI 0.697 to 1.27 $I^2 = 0\%$	RR 0.89; 95% CI 0.75 to 1.05 $I^2 = 2\%$	$I^2 = 16\%$	RR 0.79; 95% CI 0.43 to 1.47 $I^2 = 0\%$	ference between intervention and control groups (no pooling of data)
	Follow-up > 12 months 16 RCTs (5790 participants) RR 0.87; 95% CI 0.75 to 0.99 $I^2 = 0\%$	Follow-up > 12 months 6 RCTs (2845 participants) RR 0.88; 95% CI 0.75 to 1.02 $I^2 = 34\%$			Follow-up > 12 months 1 RCT (525 participants) RR 1.99; 95% CI 0.50 to 7.88	
Cardiovascular mortality	Follow-up < 12 months 9 RCTs (4130 participants) RR 0.93; 95% CI 0.71 to 1.21 $I^2 = 0.0\%$	<i>"Studies did not consistently report deaths due to heart failure or sudden death"</i>	5 RCTs (3893 participants) RR 0.80; 95% CI 0.6 to 1.00 $I^2 = 0.0\%$	NR	NR	NR
	Follow-up > 12 months 12 RCTs (4757) RR 0.74; 95% CI 0.63 to 0.87 $I^2 = 0\%$					
Hospitalisation	Follow-up < 12 months 4 RCTs (463 participants) RR 0.69; 95% CI 0.51 to 0.93 $I^2 = 12\%$	Follow-up < 12 months 15 RCTs (1328 participants) RR 0.75; 95% CI 0.62 to 0.92 $I^2 = 0\%$	NR	At end of follow-up period 4 RCTs (12,905 participants) RR 0.83; 95% CI 0.65 to 1.07 $I^2 = 32\%$	1 RCT No difference between home-based and centre-based CR	3 RCTs (numbers NR) No significant difference between intervention and control groups (no pooling of data)
	Follow-up > 12 months 7 RCTs (2009 participants) RR 0.98; 95% CI 0.87 to 1.11 $I^2 = 56\%$	Follow-up > 12 months 5 RCTs (2722 participants) RR 0.92; 95% CI 0.66 to 1.29 $I^2 = 63\%$				
HF-specific admissions	NR	Follow-up > 12 months 12 RCTs (1036 participants)	NR	1 RCT Participants in the intervention group had 41% fewer (P value = 0.05) and 61% fewer	NR	NR

Table 10. Summary of outcome results across Cochrane systematic reviews (Continued)

	RR 0.89; 95% CI 0.66 to 1.19					
	$I^2 = 20\%$					
HRQoL	10 RCTs	20 RCTs	7 RCTs	Across 11 RCTs, 81 HRQoL outcome scores/sub-scores reported:	10 RCTs	2 RCTs
	7/10 RCTs reported evidence of a significantly higher level of HRQoL with intervention at follow-up	<p>Follow-up < 12 months</p> <p>13 RCTs (1270 participants)</p> <p>MLWHF score: MD -5.8; 95% CI -9.2 to -2.4</p> <p>$I^2 = 70\%$</p> <p>Follow up > 12 months</p> <p>3 RCTs (329 participants)</p> <p>MD -9.5; 95% CI -17.54 to -1.5</p> <p>$I^2 = 73\%$</p> <p>All HRQoL measures pooled</p> <p>20 RCTs (3240 participants)</p> <p>SMD -0.5; 95% CI -0.7 to -0.3</p> <p>$I^2 = 79\%$</p>	1/7 RCTs reported evidence of a significantly higher level of HRQoL with intervention at follow-up	<p>14/81 in favour of intervention compared to control</p> <p>67/81 no significant difference between intervention and control</p> <p>5/11 RCTs reported evidence of a significantly higher level of some HRQoL domains with intervention at follow-up</p> <p>No consistent difference in HRQoL total or domain score at follow-up between intervention and control</p>	8/10 RCTs reported improvements in HRQoL at follow-up with both home-based and centre-based CR compared with baseline	1/2 RCTs reported improvement in HRQoL with intervention (not significant)
					No strong evidence of difference in overall HRQoL outcomes or domain score at follow up between home-based and centre-based CR	1/2 RCTs reported improvement in both groups but no significant difference between intervention and control
Economics	Costs	3 RCTs	NR	5 RCTs reported healthcare utilisation costs	3/4 RCTs reported healthcare costs associated with CR were lower for the home-based than centre-based programmes	NR
Costs	3 RCTs	2 studies undertook a cost effectiveness analysis and 1 reported costs		2/5 RCTs reported an overall mean net saving of USD965 per participant at 6 months follow-up and USD1420 per participant at 24 months follow-up		
Cost-effectiveness	2/3 studies reported total healthcare costs were not statistically significantly different between groups	There was no evidence of significantly different costs or outcomes		1/5 RCTs reported an increase in mean net costs of USD52 per participant	1/4 RCTs reported that home-based CR was more costly than centre-based CR but costs would be the same if participant travel costs and trav-	
	Cost-effectiveness					
	1 RCT					
	Authors concluded that rehabilitation was an efficient use of healthcare resources and may be economically justified					

Table 10. Summary of outcome results across Cochrane systematic reviews (Continued)

				2/5 RCTs reported no difference between groups		el time were included	
				No RCTs reported cost-effectiveness		8 studies reported different aspects of consumption of healthcare resources	
						No significant between group differences were seen	
All-cause withdrawal/drop-out at follow-up	NR	NR	NR	At follow-up	At follow-up	NR	
				8 RCTs (2862 participants)	18 (1894 participants)		
				RR 1.03; 95% CI 0.83 to 1.27	RR 1.04; 95% CI 1.00 to 1.08		
				$I^2 = 34%$	$I^2 = 44%$		
Uptake	NR	NR	NR	NR	NR		10 RCTs (1338 participants)
							8/10 RCTs reported uptake was significantly higher in intervention group
Adherence	NR	NR	NR	NR	14 RCTs		8 RCTs (1150 participants)
					*3/14 RCTs reported adherence was significantly higher in home-based CR		3/8 RCTs reported adherence was significantly higher in intervention group

CABG: coronary artery bypass graft; CHD: coronary heart disease; CR: cardiac rehabilitation; HF: heart failure; HRQoL: health-related quality of life; MD: mean difference; MI: myocardial infarction; MLWHF: Minnesota Living with Heart Failure questionnaire; NR: not reported; PTCA: percutaneous transluminal coronary angioplasty; RCT: randomised controlled trial; RR: risk ratio; SMD: standardised mean difference.

* As reported in the 'Summary of findings' table. Effects of interventions section states 4/14.

APPENDICES

Appendix 1. Search strategy

The Cochrane Library

#1 cardiac near/4 rehab*

#2 cardiac near/4 exercise*

#1 OR #2

Appendix 2. Excluded systematic reviews

Author	Title	Reason for exclusion from overview
Hulzebos 2012	Preoperative physical therapy for elective cardiac surgery patients	Only 1/8 included randomised controlled trials compared cardiorespiratory exercise training with a non-exercise control

Appendix 3. Protocols identified

Author	Title
Devi 2011	Internet-based interventions for the secondary prevention of coronary heart disease
Euler 2013	Interventions to support return-to-work for patients with coronary heart disease
* Mechta-Nielsen 2014	Exercise-based cardiac rehabilitation for adult patients with ICD
Risom 2014	Exercise-based cardiac rehabilitation for adults with atrial fibrillation
Sibilitz 2013	Exercise-based cardiac rehabilitation for adults after heart valve surgery

*We are aware of this proposed title through personal communication with the authors.

WHAT'S NEW

Date	Event	Description
22 December 2021	Amended	The editorial team, in agreement with the lead author Rod Taylor, have decided that no update of this review is needed currently as the topic is covered sufficiently in recent non-Cochrane publications.

HISTORY

Protocol first published: Issue 8, 2014

Review first published: Issue 12, 2014

CONTRIBUTIONS OF AUTHORS

Both authors were involved in the conception and design of the protocol and the review; undertook the study selection, data extraction and risk of bias assessment; and drafted the manuscript.

DECLARATIONS OF INTEREST

Rod Taylor was a co-author on five of the included systematic reviews.

Lindsey Anderson has no known conflict of interest.

Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews (Review)

Copyright © 2014 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

SOURCES OF SUPPORT

Internal sources

- University of Exeter Medical School, UK

External sources

- No sources of support provided

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

There were no differences between the protocol and the review.

INDEX TERMS

Medical Subject Headings (MeSH)

Coronary Disease [*rehabilitation]; Exercise Therapy [*methods]; Heart Failure [*rehabilitation]; Myocardial Infarction [*rehabilitation]; Percutaneous Coronary Intervention [*rehabilitation]; Randomized Controlled Trials as Topic; Review Literature as Topic

MeSH check words

Humans