

Neurovirulence of Herpes Simplex Virus Types 1 and 2 Isolates in Diseases of the Central Nervous System

T. Bergström^{1*}, K. Alestig², B. Svennerholm¹, P. Horal¹, B. Sköldenberg³, A. Vahlne¹

Herpes simplex virus (HSV) isolates derived from the central nervous system of ten patients with HSV-1-induced encephalitis, one patient with multiple sclerosis, and 14 patients with HSV-2-induced meningitis were investigated for neurovirulence by assaying the LD₅₀ after nose and intracerebral (i.c.) inoculation of mice. HSV-1 encephalitis strains were significantly more virulent after nose inoculation (i.e. neuroinvasive) when compared with HSV-1 isolates from patients with oral lesions only, whereas HSV-2 meningitis strains were significantly more virulent after i.c. inoculation when compared with HSV-2 isolates from patients with genital lesions only. No correlation between high neurovirulence (defined as low LD₅₀ for both routes of infection) and replication in cell cultures of neuronal and non-neuronal cell lines was found, but the weakly neurovirulent HSV-1 strain isolated from a patient with multiple sclerosis gave low replication yields. After nose inoculation, a highly neuroinvasive HSV-1 laboratory reference strain replicated to high titers in nose tissue, the trigeminal ganglia and brainstem, while a strain with low neuroinvasiveness but high i.c. virulence replicated less well in the brainstem. Neuroinvasiveness of the virus strain might be one factor of relevance in the pathogenesis of HSV-1 encephalitis in man.

When infecting the central nervous system (CNS) herpes simplex virus type 1 (HSV-1) can cause focal, necrotizing encephalitis, this virus being considered the most important cause of acute encephalitis in immunocompetent patients (1-3). The sporadic occurrence of this disease and the fact that as yet no pathogenetic factor has been singled out as playing a decisive role might indicate that in a particular patient a number of unfortunate circumstances concur. Of relevance to the pathogenesis could be predisposing factors in the patient himself, the virus dose, or the site of replication at primary infection, as well as neurovirulent properties of the virus.

Data have been presented in support of HSV-1 strain variability of neurovirulence in mouse models (4-7) and in rabbits (8). Attempts have been made to pinpoint the HSV-1 genes regulating neurovirulence by the use of recombinant strains (9-12) or deletion mutants (13), but results indicate that several genes might be in-

involved. However, whether the pathogenesis in HSV-1 encephalitis in man is influenced by the neurovirulence of HSV-1 strains is still not known.

Herpes simplex virus type 2 (HSV-2) is a rare cause of encephalitis in man (2), and in most cases reported the patients have been immunocompromised (14, 15). Instead, HSV-2 may sometimes induce a serous meningitis when HSV-2 infection spreads from the genital tract to the CNS (16, 17). Symptoms of meningitis are far more common in primary HSV-2 infection, where the surface area of involvement is larger and the duration of virus shedding longer, as compared with secondary HSV-2 infection (i.e. the first episode of HSV-2 infection preceded by an HSV-1 infection) (18). These findings indicate possible relevance of virus load from the periphery in the development of HSV-2 meningitis, but the potential virulence properties of HSV-2 strains causing meningitis have not yet been described.

We used a mouse model to assay the LD₅₀ after infection by different routes with CNS-derived HSV-1 isolates from ten patients with encephalitis and one patient with multiple sclerosis, and HSV-2 isolates from 14 patients with meningitis, in comparison with clinical reference strains isolated from oral lesions (HSV-1) and genital

¹Department of Clinical Virology, University of Göteborg, Guldhedsgatan 10b, 413 46 Göteborg, Sweden.

²Department of Infectious Diseases, University of Göteborg, Östra Hospital, 416 85 Göteborg, Sweden.

³Department of Infectious Diseases, Karolinska Institute, Danderyd Hospital, 182 88 Danderyd, Sweden.

lesions (HSV-2) in patients without neurological symptoms, and with known laboratory reference strains. The mouse is permissive to CNS infections with both HSV-1 and HSV-2, also by peripheral routes (19). It has been reported that the neurovirulence of recombinant mouse-adapted poliovirus strains in this animal is in concordance with neurovirulence in man (20).

Materials and Methods

Virus Isolation. Clinical samples of brain material, cerebrospinal fluid (CSF), and secretion from oral and genital lesions were inoculated in cell cultures of green monkey kidney (GMK AH-1 cells) or Vero cells. Isolates were typed as HSV-1 or HSV-2 by EIA using monoclonal antibodies (21, 22).

Virus Strains. Ten HSV-1 strains were isolated from brains of patients with encephalitis (HSV-1 ENC). Eight of these strains were derived from biopsies (patients no. 1–8, Table 1) performed during a trial of antiviral treatment of HSV encephalitis (3). All biopsies were taken before the commencement of antiviral treatment. In three of these patients (no. 1, 2 and 3), serological data were consistent with a probable primary HSV-1 infection (low level or absence of HSV-1 IgG antibodies and presence of HSV IgM antibodies in the first serum sample) at the time of the encephalitis, while four had serologic evidence of reactivated HSV-1 infection. In one patient, serological data were missing. Two HSV-1 strains were isolated from brain material obtained at autopsy of two patients who died of encephalitis before receiving antiviral treatment (patients no. 8 and 10, Table 1) and in whom serologic data (high levels of HSV-1 IgG antibodies at onset) indicated reactivation. As controls, we used ten HSV-1 strains isolated from oral lesions in patients without neurological symptoms seen at the Department of Dermatology, University of Göteborg. These strains are referred to as HSV-1 REF strains. Also included is an HSV-1 strain which we isolated from the CSF of a patient during the first bout of multiple

sclerosis. The isolate was designated HSV-1 BAN and was recently described in more detail (23). Laboratory reference strains (HSV-1 LABREF) used were F (B. Roizman, Chicago), McIntyre (A. Nahmias, Atlanta) and our own strain KJ 502 (24).

In fourteen patients with meningitis, HSV-2 was isolated from CSF (HSV-2 MEN) at either the Department of Virology, University of Göteborg (six strains) or at the Department of Virology, Stockholm County Central Microbiological Laboratory (eight strains). HSV-2 clinical reference strains (HSV-2 REF) were isolated from genital lesions in ten patients with HSV-2 infection without neurological symptoms seen at the Department of Dermatology, University of Göteborg. HSV-2 laboratory reference strains (HSV-2 LABREF) used were B4327UR and 7875 (S. Jeansson, Göteborg).

Preparation and Titration of Virus Stocks. In all experiments, low (second or third) passages of virus were used, and virus stocks were prepared by limited dilution of HSV strains in GMK AH-1 cell-tube cultures followed by low multiplicity infection of monolayers of GMK AH-1 cell in 200 cm² glass flasks with Eagle's medium and incubated at 37 °C. Flasks were frozen at –70 °C when an almost confluent cytopathic effect was observed. The HSV-infected cell cultures were thawed, homogenized, centrifuged at low-speed, aliquoted in 2 ml vials, and stored at –70 °C. In connection with the experiments, plaque titrations of the virus strains were performed on GMK AH-1 cells grown in 5 cm plastic dishes with 1 % methyl cellulose as overlaying medium, and the virus strains were quantified by counting plaque forming units (PFUs). HSV titers were then expressed as PFUs/ml.

LD₅₀ in Mice. Outbred Swiss albino mice were used. We preferred an outbred strain, since clear differences in susceptibility of several inbred mouse strains to HSV have been reported (25). Groups of five mice of 3 to 4 weeks of age were infected by intracerebral (i.c.) injection of 0.025 ml in the left temporal region or, when assessing neuroinvasiveness by nose inoculation, by scraping the region of the vibrissae bilaterally, and inoculation (2 × 0.05 ml) of 10 log dilutions in Hank's medium of the HSV strains. Virus quantities per mouse given were for i.c. infection 10¹–10⁵ PFUs for HSV-1 and 10²–10⁴ PFUs for HSV-2, and for

Table 1: Patient data for CNS-derived HSV-1 encephalitis strains.

| Patient no. | Strain no. | Sex | Age | Outcome | Source of strain | Status of infection ^a |
|-------------|------------|-----|-----|-----------------|------------------|----------------------------------|
| 1 | 7682 | M | 8 | death | biopsy | primary |
| 2 | 3355 | F | 19 | death | biopsy | primary |
| 3 | 25 | M | 27 | mild sequelae | biopsy | primary |
| 4 | 47 | M | 48 | mild sequelae | biopsy | reactivated |
| 5 | 164 | M | 52 | death | biopsy | reactivated |
| 6 | 2762 | M | 58 | death | biopsy | ? ^b |
| 7 | 1666 | M | 59 | severe sequelae | biopsy | reactivated |
| 8 | E4 | F | 61 | death | autopsy | reactivated |
| 9 | 274 | F | 62 | death | biopsy | reactivated |
| 10 | E5 | M | 76 | death | autopsy | reactivated |

^aEncephalitis caused by probable primary or reactivated HSV-1 infection according to serological data.

^bSerological data not available.

nose infection 10^2 – 10^6 PFUs for both HSV-1 and HSV-2. As negative control, medium alone was used. The groups of mice were coded and observed daily for two weeks after i.c. and for three weeks after nose infection for signs of neurological involvement and the number of deaths were recorded. The LD_{50} was calculated for each strain.

HSV Replication in Nose Tissue, Trigeminal Ganglia and Brainstem of Mice. Groups of 40 mice were nose inoculated (see above) with HSV-1 strains F (dilution 10^7 PFUs/ml), McIntyre and KJ 502 (dilutions 10^5 , 10^6 and 10^7 PFUs/ml). On day 1 to 7, five mice from each group were sacrificed and, after perfusion through the left heart chamber with Eagle's medium, nose tissues, trigeminal ganglia and brainstems were collected and frozen at -70°C . The samples were thawed simultaneously, homogenized in 1.5 ml of Eagle's medium containing 1% penicillin-streptomycin, and centrifuged for 10 minutes at 2500 rpm. Supernatants were inoculated on plastic dishes with GMK AH-1 cells for plaque titration as described above.

Replication Assays. Twenty-four hour replication of all HSV strains was assayed by inoculating strains on GMK AH-1 cells in plastic dishes (inoculum 0.1 PFU/cell), and on C1300 cells, a mouse neuroblastoma cell line (inoculum 1.0 PFU/cell). After adsorption for 1 h, cell cultures were rinsed four times with Eagle's medium and 2 ml Eagle's medium was added to each culture. After incubation at 37° for 24 h, the cultures were frozen for later quantification in 10 log dilutions by plaque titration on GMK AH-1 cells as described above.

Statistical Methods. The nonparametric Wilcoxon's rank sum test was used throughout for comparisons between the CNS-derived HSV strains and the oral or genital HSV strains except for the analysis of time to death after nose inoculation of the HSV-1 strains, and HSV replication in cell cultures, where Student's t-test was used.

Results

Neurovirulence in Mice after CNS and Peripheral Infection. The HSV-infected mice died of encephalitis with seizures and/or paresis after both i.c. and nose inoculation of the HSV strains and the log PFU/ LD_{50} could be calculated for each strain. By definition, death after nose inoculation was characterized as neuroinvasiveness, since this route constitutes replication and transport along the trigeminal pathway before entrance to the CNS. Neurovirulence for each strain was defined based on the combination of results of neuroinvasiveness and LD_{50} after i.c. inoculation (see below). The results from mice infected with HSV-1 strains are shown in Figure 1. After i.c. inoculation, mean log PFU/ LD_{50} for HSV-1 ENC strains did not differ significantly from mean log PFU/ LD_{50} for HSV-1 REF strains. The HSV-1 BAN strain isolated from a patient with multiple sclerosis showed a higher PFU/ LD_{50} ratio than did any of the HSV-1 ENC or REF strains.

After nose inoculation, one HSV-1 ENC strain and four HSV-1 REF strains were non-lethal with the inocula used (Figure 1). Seven of the




Figure 1: Log PFU/ LD_{50} in Swiss albino mice i.c. or nose inoculated with 10-fold dilutions of 10 HSV-1 strains isolated from brains of patients with encephalitis (Enc.), 10 HSV-1 clinical reference strains (Ref.) obtained from patients with oral lesions only, the HSV-1 BAN strain isolated from the cerebrospinal fluid of a patient with multiple sclerosis (\blacktriangle), or HSV-2 laboratory reference strains F (\otimes), McIntyre (O) and KJ 502 (\bullet). Wilcoxon's rank sum test was used for statistical comparisons. Horizontal bars indicate median values.

HSV-1 ENC strains showed a lower log PFU/ LD_{50} than did any of the HSV-1 REF strains and, as a group, the HSV-1 ENC strains were significantly ($p < 0.02$) more neuroinvasive when compared with the HSV-1 REF strains. Also, in the mice who died, the time to death after nose inoculation of 10^6 , 10^5 and 10^4 PFUs of HSV-1 ENC strains was significantly ($p < 0.01$) shorter: 7.54 ± 1.41 , 8.14 ± 1.88 and 8.78 ± 2.11 days (mean \pm SD), respectively, as compared with 8.77 ± 1.44 , 10.5 ± 1.67 and 10.33 ± 0.58 days (mean \pm SD) for the same inocula of HSV-1 REF strains.

Data from i.c. and nose inoculation in mice were combined to classify all HSV-1 strains into three classes of neurovirulence (Table 2, see legend for limit values). Of the HSV-1 ENC strains, six were classified as highly neurovirulent (Class I), three as intermediately neurovirulent (Class II), and one as weakly neurovirulent (Class III). Of the HSV-1 REF strains, none qualified as Class I, seven belonged to Class II and three to Class III, as did the isolate HSV-1 BAN derived from a patient with multiple sclerosis.

The highly neurovirulent (Class I) HSV-1 LABREF strain KJ 502 and the Class II HSV-1 LABREF strain McIntyre differed in their neuroinvasiveness (a_3 versus a_1 , Table 2). The nature of this difference was further investigated by monitoring virus replication at three levels (skin, trigeminal ganglia and brainstem) after nose inoculation with these two strains and the Class III (a_0) HSV-1 LABREF strain F. As shown in Figure 2, HSV-1 replication reached its maximum after 2 to 4 days in the periphery, after 3 to 5 days in ganglia, and after 4 to 7 days in the brainstem with all three strains. However,

Table 2: Classes of neurovirulence in mice of HSV-1 strains.

| Class | HSV-1 strains ^a | | | | | |
|-------|----------------------------|-------------------------------|------------|-------------------------------|-----------|-------------------------------|
| | Encephalitis | | Laboratory | | Reference | |
| I | 7682 ^b | a ₃ b ₃ | | | | |
| | 3355 ^b | a ₃ b ₃ | | | | |
| | E4 | a ₃ b ₃ | | | | |
| | 25 ^b | a ₃ b ₃ | | | | |
| | 2762 | a ₃ b ₂ | KJ 502 | a ₃ b ₂ | | |
| | E5 | a ₂ b ₃ | | | | |
| II | 1666 | a ₃ b ₁ | | | 90395 | a ₂ b ₂ |
| | 47 | a ₂ b ₂ | | | 90355 | a ₂ b ₂ |
| | 274 | a ₂ b ₂ | | | 90132 | a ₂ b ₂ |
| | | | | | 90147 | a ₂ b ₂ |
| | | | | | 90579 | a ₂ b ₂ |
| | | | McIntyre | a ₁ b ₂ | 90602 | a ₁ b ₂ |
| III | | | | | 90270 | a ₀ b ₃ |
| | | | | | 90238 | a ₀ b ₂ |
| | | | | | 94783 | a ₀ b ₂ |
| | | | | | HSV-1 BAN | a ₁ b ₁ |
| | 164 | a ₀ b ₁ | F | a ₀ b ₁ | 90237 | a ₀ b ₁ |

^aHSV-1 strain characteristics expressed as a: neuroinvasiveness after nose inoculation (a₀ indicates > 10⁶; 10⁶ ≥ a₁ > 2 × 10⁵; 2 × 10⁵ ≥ a₂ ≥ 2 × 10⁴; and a₃ < 2 × 10⁴ PFU/LD₅₀) and b: neurovirulence after intracerebral inoculation (b₁ indicates > 20; 20 ≥ b₂ ≥ 2; and b₃ < 2 PFU/LD₅₀). Class I (highly neurovirulent) is defined as: a + b = 5 or 6; Class II (moderately neurovirulent) as: a + b = 3 or 4; and Class III (weakly neurovirulent) as: a + b = 1 or 2.

^bSerological data indicating a probable primary HSV-1 infection as the cause of the encephalitis.

two different patterns of progression of HSV-1 infection were found: strain F (Class III) replicated well at the periphery but to low titers at both the ganglionic and brainstem levels, whereas strains McIntyre (Class II) and KJ 502 (Class I) both replicated to relatively high titers in ganglia and brainstems after high-dose inoculation peripherally. When the infectious dose was lowered, although KJ 502 replicated better at all levels, strain McIntyre barely replicated in the brainstem, while the highly neuroinvasive strain KJ 502 readily progressed to the brainstem level.

In Figure 3, results from the assessment of virulence after i.c. or nose inoculation of HSV-2 strains are depicted. Generally, log PFU/LD₅₀ values were lower for the HSV-2 strains than for the HSV-1 strains. After i.c. inoculation, HSV-2 meningitis strains were significantly ($p < 0.01$) more virulent as compared with the HSV-2 reference strains, but no difference was found after nose inoculation.

Replication Assays. The replication yields in GMK-AH1 (mean log PFU/ml ± SD) and C1300

neuroblastoma cells (mean log PFU/ml ± SD) of HSV-1 ENC strains (7.68 ± 0.36 [5.04 ± 0.86], respectively) did not differ significantly from that of the HSV-1 REF strains (7.66 ± 0.44 [5.18 ± 0.66]). When these strains were grouped according to neurovirulence, replication yields were for Class I: 7.89 ± 0.27 [5.40 ± 0.61], for Class II: 7.58 ± 0.47 [5.18 ± 0.67], and for Class III: 7.58 ± 0.21 [4.53 ± 0.70]. Strain HSV-1 BAN, grouped as Class III in neurovirulence, displayed a markedly low replication capacity in both cell lines tested (6.79 ± 0.11 [4.41 ± 0.20]). Results for HSV-1 LABREF strains were for strain F: 7.61 ± 0.04 [4.96 ± 0.26], for strain McIntyre 8.45 ± 0.02 [6.32 ± 0.12] and for strain KJ502 8.15 ± 0.02 [not tested].

Likewise, the HSV-2 MEN strains gave mean replication yields (6.26 ± 0.19 [5.37 ± 0.25]) similar to that of the HSV-2 REF strains in both cell types used (6.31 ± 0.11 [5.17 ± 0.46]). Results for HSV-2 LABREF strains were for strain B4327UR: 6.47 ± 0.07 [5.07 ± 0.09] and for strain 7875: 6.59 ± 0.07 [5.50 ± 0.11].




Figure 2: Replication yields in PFU of HSV-1 laboratory reference strains consisting of the highly neurovirulent (Class I, see Table 2) strain KJ 502 (●—●), moderately neurovirulent (Class II) strain McIntyre (○—○), and weakly neurovirulent (Class III) strain F (×—×) in A) nose tissue, B) trigeminal ganglia, and C) brainstem of mice 1 to 7 days after nose inoculation of 0.1 ml of the HSV strains diluted to 10⁵ and 10⁶ PFU/ml (KJ 502 and McIntyre), and 10⁷ PFU/ml (KJ 502, McIntyre and F). Values are given as intervals, within which mean values ± SD of each group of five mice are found.




Figure 3: Log PFU/LD₅₀ in Swiss albino mice i.c. or nose inoculated with 10-fold dilutions of 14 HSV-2 strains isolated from the cerebrospinal fluids of patients with meningitis (Men.), 10 HSV-2 clinical reference strains isolated from patients with genital lesions only (Ref.), or HSV-2 laboratory reference strains B4327UR (■) and 7875 (□). Wilcoxon's rank sum test was used for statistical comparisons. Horizontal bars indicate median values.

Discussion

The pathogenesis of encephalitis caused by HSV-1 is largely unknown. The sporadic occurrence and scarcity of underlying immunocompromising conditions found in patients (26) may suggest pathogenetic relevance of viral alteration by recombination (27) or mutation. Although the existence and genomic locations of HSV neurovirulence have been well documented in experimental studies (5, 10, 12, 13), the link between neurovirulence in humans in the form of encephalitis and neurovirulence in animal models has hitherto not been established. The HSV-1 brain isolates used in this study were all proven to be neuropathogenic to man by causing encephalitis with either a fatal outcome or sequelae in spite of antiviral treatment.

The peripheral route of nose infection, where virus particles reach the brain via the trigeminal pathway by neuron to neuron transmission (28), was chosen to imitate the probable route that HSV-1 travels when this virus causes encephalitis in humans (29). The virulence to mice displayed by encephalitis-inducing HSV-1 strains by this route of infection suggests that invasiveness is a property of relevance for HSV-1 neurovirulence. Furthermore, we have shown that the same encephalitis-inducing strains also were significantly more invasive as compared with the HSV-1 REF strains in an in vitro model using cultured rat sensory neurons in a dual-chamber system permitting infection of neuritic extensions only (30).

In experimentally infected animals, spread along sensory neurons has been found to be the main route by which HSV enters the CNS (31, 32). HSV strain variability in capacity to invade the CNS has been described (4, 6), and non-neuroinvasive strains have been found to be unable to progress from sensory ganglia to the CNS (33, 34). In our study, results of viral replication at peripheral, ganglionic and brainstem levels after nose inoculation of mice indicate that the neuroinvasive HSV-1 LABREF strain KJ502 differs from the non-neuroinvasive strains McIntyre and F by its greater ability to reach and to replicate in the brainstem.

One of the HSV-2 isolates in this study was derived from a patient who was part of an epidemiological chain of three subjects who all suffered from meningitis after genital herpes infection. This might suggest the existence of HSV-2 strains that are more prone to induce meningitis, but studies on virulence characteristics of isolates from patients with meningitis are lacking. The finding in this study that CNS-derived meningitis-inducing HSV-2 strains showed enhanced neurovirulence after i.c. inoculation in mice might indicate the existence of such a

strain characteristic. This characteristic was not revealed by the peripheral route of nose inoculation used here. The clinical picture of HSV-2 induced meningitis described earlier (16, 17) with local neurological signs in the lumbosacral region but without spread to the brain or higher regions of the spinal cord may suggest that HSV-2 is a weakly neuroinvasive virus in adults. However, studies on HSV-2 invasiveness after genital inoculation in animals with meningitis-inducing strains are needed to further address this question, since a difference exists in tropism between HSV-1 and HSV-2 (1).

A connection between HSV-1 neurovirulence in vivo and replicative capacity has been suggested (11), but conflicting data have been presented. In later studies, neither an avirulent strain (34) nor a virulent strain (10) showed any correlation between virulence and replication in different cell lines. The indications of HSV-1 and HSV-2 neurovirulence properties of CNS-derived isolates found in vivo in our study were not paralleled by higher replication yields in either a highly permissive non-neuronal (GMK-AH-1) or a mouse neuroblastoma (C1300) cell line giving low replication yields. On the other hand, a tendency towards lower replication yields in C1300 cells was seen for the Class III non-neurovirulent strains in comparison with the Class I strains ($t = 2.15$, Student's t -test). Likewise, the strain HSV-1 BAN from a patient with multiple sclerosis, which displayed weak neurovirulence in vivo, gave a markedly low replication yield in both cell types.

Restricted viral replication as a pathogenetic factor in demyelination was found in a study of mice infected with mutants of the coronavirus mouse hepatitis virus (35), and has earlier been suggested for HSV in humans in a hypothesis on multiple sclerosis (36). However, we have not been able to test this hypothesis properly, due to the lack of additional viral isolates from the CNS of patients with multiple sclerosis.

In man, HSV-1 induced CNS infections such as encephalitis are rare events although this virus is neurotropic and commonly recovered from sensory ganglia at autopsy (37). The suggestion derived from our data that the HSV-1 strain characteristic of neuroinvasiveness is linked to encephalitis indicates the usefulness of well-defined clinical isolates in the further study of HSV neuropathogenesis.

Acknowledgements

This work was financially supported by the Gothenburg Medical Society, the Medical Faculty, University of Göteborg, the Swedish Medical Society and the Swedish Medical Research Council. We thank Miss Ann-Sofie Tylö

for skillful technical assistance, and Drs Marianne Forsgren, Elisabeth Olding-Stenquist, Monica Grandien, Jonas Blomberg, and GunBritt Löwhagen for clinical HSV isolates.

References

1. **Corey L, Spear PG:** Infections with herpes simplex virus. *New England Journal of Medicine* 1986, 314: 749-757.
2. **Nahmias AJ, Whitley RJ, Visintine AN, Takei Y, Alford CA:** Herpes simplex virus encephalitis: laboratory evaluations and their diagnostic significance. *Journal of Infectious Diseases* 1982, 145: 829-836.
3. **Sköldenberg B, Forsgren M, Alestig K, Bergström T, Burman L, Dahlquist E, Forkam A, Fryden A, Lövgren K, Norlin K, Norrby R, Olding-Stenquist E, Stiernstedt G, Uhnoo I, de Vahl K:** Acyclovir versus Vidarabine in herpes simplex encephalitis. *Lancet* 1984, ii: 707-711.
4. **Dix RD, McKendall RR, Baringer JR:** Comparative neurovirulence of herpes simplex virus type 1 strains after peripheral or intracerebral inoculation of BALB/c mice. *Infection and Immunity* 1983, 40: 103-112.
5. **Kaerner HC, Schröder CH, Ott-Hartmann A, Kümel G, Kirchner H:** Genetic variability of herpes simplex virus: development of a pathogenic variant during passaging of a nonpathogenic herpes simplex virus type 1 virus strain in mouse brain. *Journal of Virology* 1983, 46: 83-93.
6. **McKendall RR:** Comparative neurovirulence and latency of HSV-1 and HSV-2 following footpad inoculation of mice. *Journal of Medical Virology* 1980, 5: 25-32.
7. **Schröder CH, Kümel G, Glorioso J, Kirchner H, Kaerner HC:** Neuropathogenicity of herpes simplex virus in mice: protection against lethal encephalitis by co-infection with a non-encephalitogenic strain. *Journal of General Virology* 1983, 64: 1973-1982.
8. **Stroop WG, Schaefer DC:** Production of encephalitis restricted to the temporal lobes by experimental reactivation of herpes simplex virus. *Journal of Infectious Diseases* 1986, 153: 721-731.
9. **Day SP, Lausch RN, Oakes JE:** Evidence that the gene for herpes simplex virus type 1 DNA polymerase accounts for the capacity of an intertypic recombinant to spread from eye to central nervous system. *Virology* 1988, 163: 166-173.
10. **Javier RT, Izumi KM, Stevens JG:** Localization of a herpes simplex virus neurovirulence gene dissociated from high-titer virus replication in the brain. *Journal of Virology* 1988, 62: 1381-1387.
11. **Thompson RL, Stevens JG:** Biological characterization of a herpes simplex virus intertypic recombinant which is completely and specifically non-neurovirulent. *Virology* 1983, 131: 171-179.
12. **Thompson RL, Wagner EK, Stevens JG:** Physical location of a herpes simplex virus type-1 gene function(s) specifically associated with a 10 million-fold increase in HSV neurovirulence. *Virology* 1983, 131: 180-192.
13. **Meignier B, Longnecker R, Mavromara-Nazos P, Sears AE, Roizman B:** Virulence and establishment of latency by genetically engineered deletion mutants of herpes simplex virus type 1. *Virology* 1988, 162: 251-254.

14. **Dix RD, Waitzmann DM, Follansbee S, Pearson BS, Mendelson T, Smith P, Davis RL, Mills J:** Herpes simplex virus type 2 encephalitis in two homosexual men with persistent lymphadenopathy. *Annals of Neurology* 1985, 17: 203-206.
15. **Linneman CC, First MR, Alvira MM, Alexander JW, Schiff GM:** Herpesvirus type 2 meningoencephalitis following renal transplantation. *American Journal of Medicine* 1976, 61: 703-708.
16. **Craig CP, Nahmias AJ:** Different patterns of neurological involvement with herpes simplex virus types 1 and 2: isolation of herpes simplex virus type 2 from the buffy coat of two adults with meningitis. *Journal of Infectious Diseases* 1973, 127: 365-372.
17. **Sköldenberg B, Jeansson S, Wolontis S:** Herpes simplex virus type 2 and acute aseptic meningitis. *Scandinavian Journal of Infectious Diseases* 1975, 7: 227-232.
18. **Corey L, Adams HG, Brown ZA, Holmes KK:** Genital herpes simplex virus infections: clinical manifestations, course and complications. *Annals of Internal Medicine* 1983, 98: 958-972.
19. **Kristensson K, Vahlne A, Persson LA, Lycke E:** Neural spread of herpes simplex virus types 1 and 2 in mice after corneal or subcutaneous (footpad) inoculation. *Journal of Neurological Sciences* 1978, 35: 331-340.
20. **La Monica N, Almond JW, Racaniello VR:** A mouse model for poliovirus neurovirulence identifies mutations that attenuate the virus for humans. *Journal of Virology* 1987, 61: 2917-2920.
21. **Nilheden E, Jeansson S, Vahlne A:** Typing of herpes simplex virus by an enzyme-linked immunosorbent assay with monoclonal antibodies. *Journal of Clinical Microbiology* 1983, 17: 677-680.
22. **Olofsson S, Lundström M, Marsden H, Jeansson S, Vahlne A:** Characterization of a herpes simplex virus type 2-specified glycoprotein with affinity for N-acetylgalactosamine-specific lectins and its identification as g92K or gG. *Journal of General Virology* 1986, 67: 737-744.
23. **Bergström T, Andersen O, Vahlne A:** Isolation of herpes simplex virus type 1 during first attack of multiple sclerosis. *Annals of Neurology* 1989, 26: 283-285.
24. **Vahlne A, Blomberg J, Olofsson S, Lycke E:** Subtyping of herpes simplex virus. *Acta Pathologica Microbiologica Scandinavica (B)* 1975, 83: 506-512.
25. **Lopez C:** Genetics of natural resistance to herpes virus infection in mice. *Nature* 1975, 258: 152-153.
26. **Whitley RJ, Lakeman AD, Nahmias AJ, Roizman B:** DNA restriction-enzyme analysis of herpes simplex virus isolates obtained from patients with encephalitis. *New England Journal of Medicine* 1982, 307: 1060-1062.
27. **Javier RT, Sedarati F, Stevens JG:** Two avirulent herpes simplex viruses generate lethal recombinants in vivo. *Science* 1986, 234: 746-748.
28. **Kristensson K, Nennesmo I, Persson L, Lycke E:** Neuron to neuron transmission of herpes simplex virus. *Journal of Neurological Sciences* 1982, 54: 149-156.
29. **Damasio A, Van Hoesen GW:** The limbic system and the localisation of herpes simplex encephalitis. *Journal of Neurology, Neurosurgery and Psychiatry* 1985, 48: 297-301.
30. **Bergström T, Lycke E:** Neuroinvasion of herpes simplex virus. An in vitro model for characterization of neurovirulent strains. *Journal of General Virology* 1990, 71: 405-410.
31. **Cook ML, Stevens G:** Pathogenesis of herpes neuritis and ganglionitis in mice: evidence of intraaxonal transport of infection. *Infection and Immunity* 1973, 7: 272-288.
32. **Kristensson K, Lycke E, Sjöstrand J:** Spread of herpes simplex virus in peripheral nerves. *Acta Neuropathologica* 1971, 17: 44-53.
33. **Oakes JR, Gray WL, Lausch RN:** Herpes simplex virus type 1 DNA sequences which direct spread of virus from cornea to central nervous system. *Virology* 1986, 150: 513-517.
34. **Thompson RL, Cook ML, Devi-Rao GB, Wagner EK, Stevens JG:** Functional and molecular analyses of the avirulent wild-type herpes simplex virus type 1 strain KOS. *Journal of Virology* 1986, 58: 203-211.
35. **Koolen MJM, Love S, Wouda W, Calafat J, Horzinek MC, van der Zeijst BAM:** Induction of demyelination by a temperature-sensitive mutant of the coronavirus MHV-A59 is associated with restriction of viral replication in the brain. *Journal of General Virology* 1987, 68: 703-714.
36. **Koprowski H, Warren KG:** Can a defective herpes simplex virus cause multiple sclerosis? *Perspectives in Biology and Medicine* 1978, 22: 10-18.
37. **Baringer JR, Swoveland P:** Recovery of herpes-simplex virus from human trigeminal ganglions. *New England Journal of Medicine* 1973, 288: 648-650.