

Cellular membrane cholesterol is required for porcine reproductive and respiratory syndrome virus entry and release in MARC-145 cells

SUN Ying^{1,2}, XIAO ShaoBo^{1,2}, WANG Dang^{1,2}, LUO Rui^{1,2}, LI Bin^{1,2},
CHEN HuanChun^{1,2} & FANG LiuRong^{1,2*}

¹State Key Laboratory of Agricultural Microbiology, Huazhong Agricultural University, Wuhan 430070, China;

²Laboratory of Animal Virology, College of Veterinary Medicine, Huazhong Agricultural University, Wuhan 430070, China

Received December 10, 2010; accepted May 27, 2011

Cholesterol represents one of the key constituents of small, dynamic, sterol- and sphingolipid-enriched domains on the plasma membrane. It has been reported that many viruses depend on plasma membrane cholesterol for efficient infection. In this study, the role of the plasma membrane cholesterol in porcine reproductive and respiratory syndrome virus (PRRSV) infection of MARC-145 cells was investigated. Pretreatment of MARC-145 cells with methyl- β -cyclodextrin (M β CD), a drug used to deplete cholesterol from cellular membrane, significantly reduced PRRSV infection in a dose-dependent manner. This inhibition was partially reversed by supplementing exogenous cholesterol following M β CD treatment, suggesting that the inhibition of PRRSV infection was specifically mediated by removal of cellular cholesterol. Further detailed studies showed that depletion of cellular membrane cholesterol significantly inhibited virus entry, especially virus attachment and release. These results indicate that the presence of cholesterol in the cellular membrane is a key component of PRRSV infection.

porcine reproductive and respiratory syndrome virus (PRRSV), cholesterol, virus entry, release, membrane fusion

Citation: Sun Y, Xiao S B, Wang D, *et al.* Cellular membrane cholesterol is required for porcine reproductive and respiratory syndrome virus entry and release in MARC-145 cells. *Sci China Life Sci*, 2011, 54: 1011–1018, doi: 10.1007/s11427-011-4236-0

Porcine reproductive and respiratory syndrome (PRRS), characterized by severe reproductive failure in sows and respiratory distress in piglets and growing pigs, is now considered one of the most economically important diseases in countries with intensive swine industries [1–3]. PRRS virus (PRRSV), the etiology of PRRS, is a positive-strand RNA virus that belongs to the genus *Arterivirus*, family Arteriviridae, along with *Lactate-dehydrogenase-elevating virus* (LDV) of mice, *Equine arteritis virus* (EAV) and *Simian hemorrhagic fever virus* (SHFV) [4]. It has been demonstrated that PRRSV has a strong tropism for cells of the monocyte/macrophage lineage *in vivo*, and for porcine al-

veolar macrophages (PAM) and African green monkey kidney cells (MARC-145) *in vitro* [5–8]. Accumulating evidence has revealed that the replication of PRRSV in cultured cells partially depends on the composition of the host cellular plasma membrane [9,10]. Cellular surface molecules, such as heparan sulfate and sialoadhesin, play an important role in PRRSV attachment and internalization [11–13]. Moreover, simian vimentin and CD151 were considered to play a role in PRRSV infection of MARC-145 cells [14,15]. However, some links within the detailed mechanism involved in PRRSV replication cycle remain poorly understood.

The infectivity of enveloped viruses often depends on the lipid composition of the host cell membrane. For instance,

*Corresponding author (email: fanglr@mail.hzau.edu.cn)

infection by *Semliki Forest virus*, a species of the *Alpha-virus* genus, requires both cholesterol and sphingolipids [16,17]. Additionally, virus entry and assembly of certain retroviruses and filoviruses depend on cholesterol-rich membrane microdomains [18,19]. Cholesterol is an abundant, key component of the eukaryotic cell membrane that forms liquid-ordered microenvironments in the plasma membrane due to its rigidity and hydrophobicity [20]. Previous studies demonstrated that plasma membrane cholesterol plays an important role in the entry and infection processes of many viruses, especially in some enveloped viruses [21], including *Human immunodeficiency virus type 1* (HIV-1) [22], *Poliovirus* [23], *Murine coronavirus* [24], *Vaccinia virus* [25], *Herpes simplex virus* [26], *Foot-and-mouth disease virus* [27], and *Severe acute respiratory syndrome-related coronavirus* (SARS-CoV) [28]. Moreover, the plasma membrane cholesterol content also affects the release of *Influenza A virus* from the infected cell [29]. However, few reports on the potential relationship between cholesterol and the replication of viruses in the *Arteriviridae* exist, although it was recently reported that cholesterol is a determinant of *Equine arteritis virus* entry [30].

In this study, we investigated whether plasma membrane cholesterol plays a role during PRRSV infection and at which step(s) it functions. Our study clearly showed that depletion of cholesterol from host cells inhibited virus infection, and primarily affected virus entry and release processes.

1 Materials and methods

1.1 Viruses, cells and reagents

PRRSV strain CH-1a, the first field isolate from China, was kindly provided by Dr. Tong G. Z. (Shanghai Veterinary Research Institute, Shanghai, China). PRRSV strain WUH3, a highly pathogenic North American-type PRRSV, was isolated in 2007 [31]. MARC-145 cells were cultured and maintained in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal bovine serum (FBS) at 37°C in a humidified 5% CO₂ incubator. Methyl- β -cyclodextrin (M β CD) and water-soluble cholesterol were purchased from Sigma (St Louis, USA).

1.2 Virus purification and biotinylation

To biotinylate PRRSV, the virus was first purified following the protocols described by Nauwynck *et al.* [11]. The titer of the purified virus was approximately 5×10^6 TCID₅₀ mL⁻¹ (50% tissue culture infectious doses mL⁻¹), as determined by a cytopathic effect (CPE) assay in MARC-145 cells. The protein concentration of the virus preparation was 2 mg mL⁻¹, as determined by the Bradford assay with bovine serum albumin (BSA) used as a protein quantitation

standard. Biotinylation was performed with a protein biotinylation kit (GE Healthcare, Uppsala, Sweden), according to manufacturer's instructions. The biotinylated viruses were collected after purification on a Sephadex G-25 column and diluted in phosphate-buffered saline (PBS; pH 7.4) at a concentration of 0.2 mg mL⁻¹ and stored at -80°C. Virus infectivity and the ability to enter cells were not significantly decreased after being biotinylated.

1.3 M β CD treatment of cells

1.3.1 M β CD toxicity assay

To determine the toxicity of M β CD, confluent MARC-145 cells in 96-well plates were treated with serum-free DMEM with or without M β CD. After incubation at 37°C for 1 h, M β CD-treated cells and mock-treated cells were harvested for the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay to detect cell viability.

1.3.2 Measurement of cellular cholesterol content

To determine if treatment with M β CD reduces the cellular cholesterol content, MARC-145 cells in 24-well plates were washed three times with cold PBS, and then incubated for 1 h with serum-free DMEM (control cells) or various concentrations of M β CD (treated cells). After three additional washes with PBS, cells were harvested and the cholesterol content was measured using the Amplex Red Cholesterol Assay Kit (Molecular Probes, Oregon, USA), as directed by the manufacturer.

1.3.3 Cholesterol depletion

To deplete plasma membrane cholesterol, confluent MARC-145 cells in 24-well plates were washed three times with cold PBS, and then incubated for 1 h with serum-free DMEM (control cells) or DMEM containing 10 mmol L⁻¹ M β CD (treated cells).

1.3.4 Cholesterol replenishment

To replenish plasma membrane cholesterol, the protocols established by Popik *et al.* [32] were used. In brief, confluent MARC-145 cells in 24-well plates were treated with serum-free DMEM (control cells) or DMEM containing 10 mmol L⁻¹ M β CD (treated cells) for 1 h. Cells were then washed three times with PBS and incubated with serum-free DMEM alone or DMEM containing 400 μ g mL⁻¹ water-soluble cholesterol for 1 h. After three washes with PBS, cells were harvested for cholesterol level detection or infected with PRRSV.

1.4 RNA extraction and real-time RT-PCR

Total RNA was extracted from cell suspensions or supernatant with RNAprep Micro Kit (Tiangen, Beijing, China), according to the manufacturer's instructions. RNA was re-

versetranscribed into cDNA using ReverTra Ace- α (Toyobo, Osaka, Japan) with Oligo (dT)₂₀. Quantitative real-time PCR was performed according to protocols described by Egli *et al.* [33]. In brief, the generated cDNA was amplified by quantitative real-time PCR using primers for the PRRSV ORF7 gene, reverse primer (5'-AAATGGGGCTTCTCCG-GGTTTT-3') and forward primer (5'-TCAGCTGTGCCA-AATGCTGG-3'), and specific fluorescent probe (5'-TCC-CGGTCCCTTGCCTCTGGA-3'; sense orientation), which was 5'-labeled with 6-carboxyfluorescein (FAM; reporter) and tetramethylrhodamine (TAMRA) at its 3'-end. Real-time PCR was performed in a single tube in an ABI Prism 7500 Real-Time PCR System (Applied Biosystems, Foster City, CA). For each run, four no-template controls containing water instead of DNA, as well as five PRRSV ORF7 gene standards corresponding to 10¹², 10¹¹, 10¹⁰, 10⁹, 10⁸ copies per tube were included. Samples of interest were run in triplicate. Data from the TaqMan[®] run were analyzed using the SDS software (Version 1.7; Applied Biosystems).

1.5 Virion penetration analysis using confocal microscopy

Confluent MARC-145 cells in 24-well plates were incubated with 10 μ g of biotinylated PRRSV in PBS per well at 4°C for 1 h to achieve the maximal virus-cell attachment. Supernatants were removed and the cells were washed three times with cold PBS. Cells were next left untreated, or treated with M β CD, or treated with M β CD and cholesterol replenishment, as described above. During the treatment, viruses complete the penetration process. After removal of the supernatant, cells were incubated in DMEM containing 2% FBS for 1 h, and then washed three times with cold PBSA and incubated with 1:50-diluted streptavidin-fluorescein isothiocyanate (FITC) for 1 h on ice. Subsequently, cells were washed once with PBS and resuspended in cold PBS containing 4% paraformaldehyde and fixed for 10 min at room temperature. After removal of paraformaldehyde, cells were washed twice with PBS containing 2% FBS, and the biotinylated virions were visualized on a confocal laser scanning microscope (LSM 510 Meta; Carl Zeiss, Gottingen, Germany).

1.6 Statistical analysis

Student's *t*-test was used to evaluate differences in results and *P*<0.05 was considered statistically significant.

2 Results

2.1 M β CD treatment of MARC-145 reduces the cellular cholesterol content and has little effect on cell viability

To investigate the effect of M β CD on the cholesterol level

of MARC-145 cells, MARC-145 cells were mock-treated or treated with various concentrations of M β CD. As shown in Figure 1A, M β CD treatment significantly reduced cellular cholesterol levels in a dose-dependent manner. When the concentration of M β CD was up to 10 mmol L⁻¹, cellular cholesterol level was reduced by approximately 75% compared with that in mock-treated cells. As the concentration of M β CD was further increased (>10 mmol L⁻¹), cellular cholesterol level was no longer reduced, indicating that the remaining 25% cholesterol represented threshold levels of cholesterol which could not be extracted by higher doses of M β CD.

After confirming the effect of M β CD on MARC-145 cellular cholesterol, analyses were performed to exclude the possibility that M β CD causes cell toxicity. To this end, cell viabilities were detected after treatment with various concentration of M β CD by MTT. As shown in Figure 1B, M β CD had little cytotoxicity on MARC-145 cells when the concentration of M β CD was <10 mmol L⁻¹ (Figure 1B). Accordingly, 10 mmol L⁻¹ M β CD was used for all the subsequent experiments.

2.2 Cholesterol depletion inhibits PRRSV infection

It is known that the plasma membrane lipid composition plays a key role in the infection by many enveloped viruses




Figure 1 The effect of M β CD on cholesterol depletion and MARC-145 cell viability. A, Reduction of cholesterol level in MARC-145 cells. B, Toxicity of M β CD to MARC-145 cells. Error bars indicate the standard deviations of three independent experiments.

[24–28]. To investigate the role of plasma membrane cholesterol in the infection of PRRSV, MARC-145 cells were mock-treated or treated with various concentrations of M β CD and infected with PRRSV strain CH-1a at $10^{4.5}$ TCID₅₀ per well. After incubation for 36 h, cell suspensions were harvested and virus production was determined by TCID₅₀ assay on MARC-145 cells. As shown in Figure 2A, M β CD treatment inhibited the infection of PRRSV strain CH-1a in a dose-dependent manner, suggesting that cholesterol is necessary for PRRSV replication in MARC-145 cells. Moreover, treatment with 10 mmol L^{-1} M β CD efficiently reduced virus titer to $10^{2.75}$ TCID₅₀ mL⁻¹ compared with that of $10^{6.5}$ TCID₅₀ mL⁻¹ in mock-treated cells.

To further elucidate the effects of M β CD on the replication cycle of PRRSV, mock-treated and M β CD-treated cells were infected with PRRSV strain CH-1a at $10^{4.5}$ TCID₅₀ per well. Cell suspensions were harvested at various times post-infection (p.i.) and the titers of virus production in this

cell suspension were determined. As shown in Figure 2B, virus reproduction was significantly inhibited in M β CD-treated cells, and virus titers at all stages of the replication cycle were notably reduced compared with those in mock-treated cells. Specifically, at 36 h p.i., virus titers in M β CD-treated cells were 7.9×10^3 -fold lower than that in mock-treated cells. We also investigated the role of plasma membrane cholesterol in the infection of PRRSV strain WUH3, a highly pathogenic PRRSV that has been emerging in China in recent years. Significant inhibitory effects were observed (Figure 2C), indicating that the inhibitory effects of plasma membrane cholesterol on PRRSV replication are not strain-specific.

If the inhibition of PRRSV replication was specifically due to the cholesterol depletion by M β CD, then cholesterol replenishment in M β CD-treated cells should restore PRRSV replication. To clarify this hypothesis, M β CD-treated cells were incubated with DMEM containing water-soluble cho-




Figure 2 Cholesterol depletion inhibits PRRSV infection. A, Virus titers determined by TCID₅₀ (50% tissue culture infective dose) assay at 36 h p.i. after cholesterol depletion. B and C, Virus yields of PRRSV strains CH-1a (B) and WUH3 (C) from mock-treated (■) and M β CD-treated (10 mmol L^{-1}) (▲) cells at various times after infection. D, Cholesterol levels measured from equal number of mock-treated, M β CD-treated or M β CD-cholesterol-treated cells. E, Virus titer at 36 h p.i. after cholesterol depletion or cholesterol replenishment. Error bars indicate the standard deviations of three independent experiments.

lesterol and then cholesterol levels were measured. As shown in Figure 2D, the cellular cholesterol level of M β CD-cholesterol-treated cells was partially restored to 85% of mock-treated cell cholesterol levels. Subsequently, virus yields from the infected M β CD-cholesterol-treated cells were assayed and compared with that from M β CD-treated cells. The titer of PRRSV from the infected M β CD-cholesterol-treated cells was increased to approximately 3.35×10^3 -fold of that from M β CD-treated cells (Figure 2E). These results definitively demonstrated that decreasing cellular cholesterol levels resulted in the reduced ability of PRRSV replication and with the replenishment of cellular cholesterol, virus replication was restored to a certain extent.

2.3 Cholesterol depletion inhibits PRRSV entry

To further illuminate at which step(s) cholesterol depletion reduces PRRSV replication, we first investigated the effect of cholesterol depletion on virus entry. Mock-treated, M β CD-treated and M β CD-cholesterol-treated MARC-145 cells in 24-well plates were infected with PRRSV at $10^{4.5}$ TCID₅₀ per well. After incubation for 1 h, cells were harvested and intracellular PRRSV RNA levels were measured by real-time RT-PCR. As shown in Figure 3A, compared with the number of viral genomes in mock-treated cells, a significant decrease (80%) in viral genomes in M β CD-treated cells was detected. Furthermore, cholesterol replenishment partially restored intracellular viral genomes and there was only a 17% reduction after adding exogenous cholesterol. These results suggest that entry of PRRSV is dependent on the cellular membrane cholesterol.

The entry phases of PRRSV infection can be divided into two stages: the first stage requires PRRSV attachment to cells, and the second stage occurs when PRRSV penetrates the cellular membrane. To determine whether M β CD reduces PRRSV attachment to MARC-145 cells, mock-treated, M β CD-treated or M β CD-cholesterol-treated MARC-145 cells were infected with PRRSV at 4°C for 1 h. All cells were collected and virus genomes were detected by quantitative real-time PCR assay. As shown in Figure 3B, virus genomes presented in M β CD-treated cells were only 23.8% of that presented in mock-treated cells. Furthermore, cholesterol replenishment partially restored virus attachment to approximately 75% of which presented in mock-treated cells, suggesting that the effect was specifically due to the cholesterol depletion (Figure 3B).

The effect of cholesterol depletion on virus attachment was defined, but next we determined whether penetration of PRRSV into MARC-145 cell was also affected by cholesterol depletion. To this end, MARC-145 cells in 24-well plates were infected with 10 μ g biotinylated-PRRSV per well and were mock-treated or treated with M β CD or




Figure 3 Cholesterol depletion inhibits PRRSV entry. A, Quantitative RT-PCR analysis of PRRSV entry. B, Quantitative RT-PCR analysis of PRRSV attachment. Error bars indicate the standard deviations of three independent experiments.

M β CD-cholesterol. During the treatment, virions attached on these cells completed the penetration process and were detected by confocal microscopy. No significant differences were observed between various groups (data not shown), suggesting that M β CD treatment has little effect on the second stage of virus entry. Altogether, these data indicated that cellular membrane cholesterol was required for PRRSV entry into MARC-145 cells, and its depletion by M β CD mainly affects virus attachment, rather than penetration.

2.4 Cholesterol depletion results in an enhanced release of virus particles with reduced infectivity

To investigate the effect of M β CD on virus release from MARC-145 cells, cells were infected with PRRSV strain CH-1a at $10^{4.5}$ TCID₅₀ per well. At 24 h p.i., cells were left untreated or treated with M β CD or M β CD-cholesterol. At the indicated time points post-treatment, the supernatants were harvested and the virus was detected by real-time RT-PCR. The data indicated that the number of viral genomes present in M β CD-treated cells was notably higher than those in mock-treated cells at each time point, and this increase was partially neutralized by cholesterol replenishment (Figure 4A). This suggests that the ability of virus release from MARC-145 is inversely related to cellular cholesterol level.

To further determine the effect of cholesterol depletion




Figure 4 Cholesterol depletion results in an enhanced release of virus particles with reduced infectivity. A, Quantitative RT-PCR measurement of PRRSV genomes released in cholesterol depleted or cholesterol replenished cell culture supernatant at various times after the treatment. B, Virus infectivity of virions released from mock-treated, MβCD-treated and MβCD-cholesterol-treated cells. Error bars indicate the standard deviations of three independent experiments.

on virus release, the infectivity of released virus was examined. MARC-145 cells were infected with harvested supernatant at the same amount according to the results of real-time RT-PCR. At 36 h p.i., the replication of PRRSV in MARC-145 cells was analyzed by detecting viral genomes in infected cell suspensions by real-time RT-PCR. As shown in Figure 4B, the amount of PRRSV genomes expressed in cells infected with released virus from MβCD-treated cells was approximately 10.6% of that in cells infected with released virus from mock-treated cells, and cholesterol replenishment partially counteracted this effect. These findings indicate that the infectivity of virus released from MβCD-treated cells was decreased compared with those released from MβCD-cholesterol-treated and mock-treated cells. According to these data, we can conclude that MβCD treatment caused an enhancement of virus particle released from infected cells but the infectivity of released viruses was decreased by this treatment.

3 Discussion

Many studies have reported that cholesterol is involved in the replication of some enveloped virus, such as *Herpes simplex virus* [26], *Vaccinia virus* [25], *Murine coronavirus* [24], SARS-CoV [28], *Varicella-zoster virus* [34], and

Pseudorabies virus [35]. The present study showed that the inhibition of PRRSV infection *in vitro* occurred as host cell membrane cholesterol decreased due to MβCD treatment. This effect could be counteracted by the addition of exogenous cholesterol and demonstrated the importance of membrane cholesterol for PRRSV infection.

Being an essential component of cell membrane, cholesterol could potentially inhibit virus entry by several different mechanisms. First, cholesterol may affect virus entry by modifying the interaction of the virus particle with host cell membrane. Cholesterol level is important for maintaining biological membrane fluidity [36], and its removal could reduce lateral diffusion within the membrane. This reduction in fluidity could probably influence entry of PRRSV within the membrane. Second, the lipid environment, including cholesterol level, is known to affect the charge properties of some ion channels [37], and the ion channels formed on cellular membrane require the presence of cholesterol for effective cytoplasmic delivery of the viral genome during virus entry. Cholesterol and sphingolipids are components of lipid rafts, and lipid recognition by certain proteins of viruses might be imperative for virus entry [38]. Moreover, there are also other studies indicating that cholesterol removal resulted in an inhibition of cellular signaling pathways [39].

A previous study demonstrated that the entry of *Equine arteritis virus*, another member of Arteriviridae, could also be inhibited by cholesterol depletion, and virus entry requires plasma membrane cholesterol [30]. In the current study, we examined the effect of plasma membrane cholesterol on PRRSV attachment and penetration, the two stages of virus entry. We found that cholesterol mainly affects virus attachment, rather than penetration. It was reported that PRRSV enters porcine alveolar macrophages (PAMs) via receptor-mediated endocytic vesicles and the low pH-dependent endocytic pathway [11]. Thus, it is possible that the effect of plasma membrane cholesterol on PRRSV entry into MARC-145 cell is also due to its influence on the endocytosis process.

A significant finding of the present study is that cholesterol depletion facilitated the release process of PRRSV but also resulted in decreased infectivity of released virus particles. This finding is consistent with reports showing that cholesterol depletion caused an enhancement of *Influenza A virus* particle release from infected cells and a decreased infectivity of virus particles [29]. In addition, the infectivity of *Newcastle disease virus* could be reduced by cholesterol depletion and possibly due to virus particles released from cholesterol-depleted cells having heterogeneous densities and structural abnormalities [40]. Teissier and Pecheur [38] had demonstrated that the molecular shape of lipids, including cholesterol and sphingolipid, could affect membrane curvature and further influence membrane defor-

mation and fluidity. Thus, structural abnormalities of virus particles due to cholesterol depletion and the physical function of cholesterol in the plasma membrane may be responsible for the findings reported herein.

In conclusion, the current study demonstrated that both efficient entry of PRRSV into MARC-145 cells and the release of PRRSV are dependent on the presence of plasma membrane cholesterol. These data suggest that further investigation of the prevention and control of PRRSV, in particular, prevention of viral entry, is warranted.

This work was supported by the National Natural Science Foundation of China (Grant No. 30770082) and National Basic Research Program of China (Grant No. 2005CB523200).

- 1 Rossow K D. Porcine reproductive and respiratory syndrome. *Vet Pathol*, 1998, 35: 1–20
- 2 Meulenbergh J J. PRRSV, the virus. *Vet Res*, 2000, 31: 11–21
- 3 Mateu E, Diaz I. The challenge of PRRS immunology. *Vet J*, 2008, 177: 345–351
- 4 Plagemann P G, Moennig V. Lactate dehydrogenase-elevating virus, equine arteritis virus, and simian hemorrhagic fever virus: a new group of positive-strand RNA viruses. *Adv Virus Res*, 1992, 41: 99–192
- 5 Benfield D A, Nelson E, Collins J E, et al. Characterisation of swine infertility and respiratory syndrome (SIRS) virus (isolate ATCC VR-2332). *J Vet Diagn Invest*, 1992, 4: 127–133
- 6 Duan X, Nauwynck H J, Pensaert M B. Effects of origin and state of differentiation and activation of monocytes/macrophages on their susceptibility to porcine reproductive and respiratory syndrome virus (PRRSV). *Arch Virol*, 1997, 142: 2483–2497
- 7 Kim H S, Kwang J, Yoon I J, et al. Enhanced replication of porcine reproductive and respiratory syndrome (PRRS) virus in a homogeneous subpopulation of MA-104 cell line. *Arch Virol*, 1993, 133: 477–483
- 8 Wensvoort G, de Kluyver E P, Pol J M, et al. Lelystad virus, the cause of porcine epidemic abortion and respiratory syndrome: a review of mystery swine disease research at Lelystad. *Vet Microbiol*, 1992, 33: 185–193
- 9 Kreutz L C. Cellular membrane factors are the major determinants of porcine reproductive and respiratory syndrome virus tropism. *Virus Res*, 1998, 53: 121–128
- 10 Van Gorp H, Van Breedam W, Delputte P L, et al. The porcine reproductive and respiratory syndrome virus requires trafficking through CD163-positive early endosomes, but not late endosomes, for productive infection. *Arch Virol*, 2009, 154: 1939–1943
- 11 Nauwynck H J, Duan X, Favoreel H W, et al. Entry of porcine reproductive and respiratory syndrome virus into porcine alveolar macrophages via receptor-mediated endocytosis. *J Gen Virol*, 1999, 80: 297–305
- 12 Delputte P L, Costers S, Nauwynck H J. Analysis of porcine reproductive and respiratory syndrome virus attachment and internalization: distinctive roles for heparan sulphate and sialoadhesin. *J Gen Virol*, 2005, 86: 1441–1445
- 13 Van Breedam W, Delputte P L, Van Gorp H, et al. Porcine reproductive and respiratory syndrome virus entry into the porcine macrophage. *J Gen Virol*, 2010, 91: 1659–1667
- 14 Kim J K, Fahad A M, Shanmukhappa K, et al. Defining the cellular target(s) of porcine reproductive and respiratory syndrome virus blocking monoclonal antibody 7G10. *J Virol*, 2006, 80: 689–696
- 15 Shanmukhappa K, Kim J K, Kapil S. Role of CD151, a tetraspanin, in porcine reproductive and respiratory syndrome virus infection. *J Virol*, 2007, 4: 62
- 16 Kielian M C, Helenius A. Role of cholesterol in fusion of Semliki forest virus with membranes. *J Virol*, 1984, 52: 281–283
- 17 Nieva J L, Bron R, Corver J, et al. Membrane fusion of Semliki forest virus requires sphingolipids in the target membrane. *EMBO J*, 1994, 13: 2797–2804
- 18 Bavari S, Bosio C M, Wiegand E, et al. Lipid raft microdomains: a gateway for compartmentalized trafficking of Ebola and Marburg viruses. *J Exp Med*, 2002, 195: 593–602
- 19 Del Real G, Jimenez-Baranda S, Lacalle R A, et al. Blocking of HIV-1 infection by targeting CD4 to nonraft membrane domains. *J Exp Med*, 2002, 196: 293–301
- 20 Simons K, Ikonen E. Functional rafts in cell membranes. *Nature*, 1997, 387: 569–572
- 21 Chazal N, Gerlier D. Virus entry, assembly, budding, and membrane rafts. *Microbiol Mol Biol Rev*, 2003, 67: 226–237
- 22 Viard M, Parolini I, Sargiacomo M, et al. Role of cholesterol in human immunodeficiency virus type 1 envelope protein-mediated fusion with host cells. *J Virol*, 2002, 76: 11584–11595
- 23 Danthi P, Chow M. Cholesterol removal by methyl-beta-cyclodextrin inhibits poliovirus entry. *J Virol*, 2004, 78: 33–41
- 24 Thorp E B, Gallagher T M. Requirements for CEACAMs and cholesterol during murine coronavirus cell entry. *J Virol*, 2004, 78: 2682–2692
- 25 Chung C S, Huang C Y, Chang W. Vaccinia virus penetration requires cholesterol and results in specific viral envelope proteins associated with lipid rafts. *J Virol*, 2005, 79: 1623–1634
- 26 Bender F C, Whitbeck J C, Ponce de Leon M, et al. Specific association of glycoprotein B with lipid rafts during herpes simplex virus entry. *J Virol*, 2003, 77: 9542–9552
- 27 Martin-Acebes M A, Gonzalez-Magaldi M, Sandvig K, et al. Productive entry of type C foot-and-mouth disease virus into susceptible cultured cells requires clathrin and is dependent on the presence of plasma membrane cholesterol. *Virology*, 2007, 369: 105–118
- 28 Li G M, Li Y G, Yamate M, et al. Lipid rafts play an important role in the early stage of severe acute respiratory syndrome-coronavirus life cycle. *Microbes Infect*, 2007, 9: 96–102
- 29 Barman S, Nayak D P. Lipid raft disruption by cholesterol depletion enhances influenza A virus budding from MDCK cells. *J Virol*, 2007, 81: 12169–12178
- 30 Nitschke M, Korte T, Tievesch C, et al. Equine arteritis virus is delivered to an acidic compartment of host cells via clathrin-dependent endocytosis. *Virology*, 2008, 377: 248–254
- 31 Li B, Xiao S, Wang Y, et al. Immunogenicity of the highly pathogenic porcine reproductive and respiratory syndrome virus GP5 protein encoded by a synthetic ORF5 gene. *Vaccine*, 2009, 27: 1957–1963
- 32 Popik W, Alce T M, Au W C. Human immunodeficiency virus type 1 uses lipid raft-colocalized CD4 and chemokine receptors for productive entry into CD4⁺ T cells. *J Virol*, 2002, 76: 4709–4722
- 33 Egli C, Thur B, Liu L, et al. Quantitative Taqman RT-PCR for the detection and differentiation of European and North American strains of porcine reproductive and respiratory syndrome virus. *J Virol Methods*, 2001, 98: 63–75
- 34 Hambleton S, Steinberg S P, Gershon M D, et al. Cholesterol dependence of varicella-zoster virion entry into target cells. *J Virol*, 2007, 81: 7548–7558
- 35 Desplanques A S, Nauwynck H J, Vercauteren D, et al. Plasma membrane cholesterol is required for efficient pseudorabies virus entry. *Virology*, 2008, 376: 339–345
- 36 Burger K, Gimpl G, Fahrenholz F. Regulation of receptor function by cholesterol. *Cell Mol Life Sci*, 2000, 57: 1577–1592
- 37 Chang H M, Reistetter R, Mason R P, et al. Attenuation of channel kinetics and conductance by cholesterol: an interpretation using

- structural stress as a unifying concept. *J Membr Biol*, 1995, 143: 51–63
- 38 Teissier E, Pecheur E I. Lipids as modulators of membrane fusion mediated by viral fusion proteins. *Eur Biophys J*, 2007, 36: 887–899
- 39 Stulnig T M, Berger M, Sigmund T, *et al.* Signal transduction via glycosyl phosphatidylinositol-anchored proteins in T cells is inhibited by lowering cellular cholesterol. *J Biol Chem*, 1997, 272: 19242–19247
- 40 Laliberte J P, McGinnes L W, Peebles M E, *et al.* Integrity of membrane lipid rafts is necessary for the ordered assembly and release of infectious Newcastle disease virus particles. *J Virol*, 2006, 80: 10652–10662

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.