



Review article

The dual role of alpha7 nicotinic acetylcholine receptor in inflammation-associated gastrointestinal cancers



Khalil Hajiasgharzadeh^a, Mohammad Hossein Somi^b, Saeed Sadigh-Eteghad^c, Ahad Mokhtarzadeh^a, Dariush Shafehbandi^a, Behzad Mansoori^a, Ali Mohammadi^d, Mohammad Amin Doustvandi^a, Behzad Baradaran^{a,*}

^a Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

^b Liver and Gastrointestinal Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

^c Neurosciences Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

^d Department of Cancer and Inflammation Research, Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark

ARTICLE INFO

Keywords:
 Clinical research
 Gastrointestinal system
 Immunology
 Oncology
 Pharmacology
 Physiology
 Alpha7 nicotinic acetylcholine receptor
 Cholinergic anti-inflammatory pathway
 Gastrointestinal cancers
 Inflammation
 Cancer regulation

ABSTRACT

Alpha7 nicotinic acetylcholine receptor ($\alpha 7\text{nAChR}$) is one of the main subtypes of nAChRs that modulates various cancer-related properties including proliferative, anti-apoptotic, pro-angiogenic and pro-metastatic activities in most of the cancers. It also plays a crucial role in inflammation control through the cholinergic anti-inflammatory pathway in numerous pathophysiological contexts. Such diverse physiological and pathological functions that initiate from this receptor may have significant impacts in determining the outcome of different cancers. Various tissues of gastrointestinal (GI) cancers such as gastric, colorectal, pancreatic and liver cancers have shown the up-regulated expression of $\alpha 7\text{nAChR}$ as compared to normal adjacent tissues. According to the well-established connection between inflammation and tumorigenesis in the digestive system, there are mounting studies demonstrated either stimulatory or inhibitory effects of $\alpha 7\text{nAChR}$ signaling in the development of GI cancers. To date, the precise underlying mechanisms related to this receptor in patients with GI cancers have not been fully elucidated. Regarding the paradoxical modulatory effects of this receptor in carcinogenesis, in this review, we aim to summarize the accumulated evidence about the involvement of $\alpha 7\text{nAChR}$ in inflammation-associated GI cancers. It seems that the complex influences of $\alpha 7\text{nAChR}$ may be a promising target in designing novel strategies in the treatment of such pathologic conditions.

1. Introduction

Gastrointestinal (GI) cancers is a common term that generally refers to the different types of cancers that influence the digestive system. These cancers originate from cells in the esophagus, stomach, exocrine pancreas, liver, gallbladder, biliary tract, small intestine, colon, rectum, and anus [1]. All of them are among the most frequent types of malignancies and represent an important public health problem with an immense burden on patients and societies worldwide [2]. Of note, gastric cancer, colorectal cancer and hepatocellular carcinoma (i.e. common type of liver cancer) are the preeminent causes of cancer-related deaths, so that they constitute ~25% of all cancer-related mortalities [2]. GI cancers are classified among multifactorial diseases caused by many different reasons, such as chronic inflammation and infection,

environmental carcinogenic risk factors, and genetic predisposition in individuals [3].

In recent decades, mounting studies in humans, as well as several experimental studies in animals, points to crucial impacts of nicotinic acetylcholine receptors (nAChRs) in various cancers initiation, progression, and metastasis [4]. Among different subtypes of nAChRs, homo-pentameric alpha7-subtype of nAChR ($\alpha 7\text{nAChR}$) appears to be of particular significance in cancer research and is one of the major regulators of a wide variety of human cancers [5]. These receptors are named based on their activation by acetylcholine (ACh) as their predominant endogenous ligand and nicotine, which is an exogenous substance. They are functionally expressed by a variety of human cancer cell and tissues such as GI cancers. Many different GI tumor cells such as gastric [6], colorectal [7], pancreas [8], and liver cell types [9], as well as GI

* Corresponding author.

E-mail address: baradarban@tbzmed.ac.ir (B. Baradaran).

tumor-infiltrating immune cells including lymphocytes [10], monocytes [11], and macrophages [12] express $\alpha 7nAChR$ subunits.

Here, we highlighted an important and of growing interest topic of the role of $\alpha 7nAChR$ in digestive carcinogenesis. Accumulating evidence has shown that increased activity of this receptor has been associated with cancer cell growth, survival, proliferation, angiogenesis, and metastasis [13, 14, 15]. Also, it has been reported that $\alpha 7nAChR$ can participate in the regulation of tumor stem cells, tumor microenvironment, and epithelial-mesenchymal transition, thus promoting the progress of tumors [15, 16]. As another side of the same coin, $\alpha 7nAChR$ is an essential regulator of inflammation and its anti-inflammatory function emerged as a novel therapeutic approach for inflammation-based diseases in recent years [17, 18]. This dual role of $\alpha 7nAChR$ in the modulation of GI inflammation and cancers is the main subject of the current review (Figure 1). It seems that the diverse physiological actions and associated disorders have made of $\alpha 7nAChR$ a promising target for effective drug development in GI cancers prevention and treatment.

2. GI inflammation and cancers

About two hundred years ago, Rudolph Virchow noticed that inflammatory cells have existed in tumor tissues proposing that persistent inflammation performs a central role in carcinogenesis [19]. Since then it has been identified that approximately twenty-five percent of all cancers are associated with chronic inflammation and mounting evidence implies that this inflammatory condition is one of the main precursors of some cancers [19]. Among these cancers, GI tract cancers are well-known cancer types that frequently arise as a result of chronic inflammation [20]. The inflammatory microenvironment has been considered as an intrinsic niche for these tumors development and progression [21]. In recent years, the causal relationship between activation of some inflammatory transcriptional factors such as nuclear factor-kappaB (NF- κ B) and production of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF- α) with cancers pathogenesis has been a major focus [22, 23]. The inflammatory responses promote both the proliferation and

survival of tumor cells and also enhance their angiogenesis and metastasis [24]. On another hand, epidemiological studies suggest that anti-inflammatory drugs reduce the incidence rate and mortality from GI cancers [25]. Therefore, it seems that inhibition of inflammatory transcription factors and downregulation of pro-inflammatory cytokines may be valuable anticancer drug candidates [26]. Considering these facts, it is important to mention that the inflammation surely plays a role in carcinogenesis, but it is not the only phenomenon, and in some cancers, it is not the most important; but here our main intention is the inflammation-associated GI cancers and modulatory effects of $\alpha 7nAChR$ in these types of cancers.

There are numerous findings indicate that the activation of $\alpha 7nAChR$ and downstream signaling pathways could significantly attenuate inflammatory responses and suggests that this receptor could be an effective therapeutic target for inflammatory diseases [18, 27]. In these studies, $\alpha 7nAChR$ activation via vagus nerve stimulation or other ACh synthesizing cells (e.g. T cells) [28], as well as exogenous agonists application possesses beneficial anti-inflammatory properties [29]. For the initial time, Tracey and colleagues exhibit that the “cholinergic anti-inflammatory pathway” (CIP) modulates the immune system through mechanisms that act on $\alpha 7nAChR$ expressed on immune cells [17]. In their proposed model, ACh released from vagal terminals or from other ACh synthesizing cells interacts with $\alpha 7nAChRs$ and suppressed pro-inflammatory cytokine production and inflammation [30]. The molecular mechanisms and signaling pathways that lead to the anti-inflammatory effects following induction of $\alpha 7nAChR$ and inflammation-suppressing effects of vagus nerve have been reviewed in several articles and are not the subject of the current review [30]. To date, different studies have been conducted to investigate the anti-inflammatory actions of CIP and $\alpha 7nAChRs$ in a variety of tissues including the GI system [31]. The outcome of such experiments has indicated that activation of $\alpha 7nAChR$ initiates intracellular signal transduction cascades that significantly downregulates the nuclear translocation of NF- κ B and attenuates the production of inflammatory cytokines such as TNF- α in numerous pathological conditions [32, 33, 34].




Figure 1. Schematic representation of the cancer-promoting and anti-inflammatory effects of $\alpha 7nAChR$ in the gastrointestinal (GI) tumor microenvironment. Both of the GI tumor cells, as well as the GI tumor-infiltrating immune cells, express $\alpha 7nAChRs$ which activated by several endogenous or exogenous ligands. These receptors exert either stimulatory or inhibitory effects in the GI-tumor processes. Such paradoxical effects should be carefully considered in the designing of novel pharmacological or surgical therapeutic strategies for the patients suffering from GI cancers.

35, 36]. Therefore the therapeutic potential of $\alpha 7$ nAChR in the inflammatory microenvironment of tumors and the protective role of the vagal nerve activity in carcinogenesis should be carefully considered [37]. Further to this immunomodulatory effect of $\alpha 7$ nAChR, previous studies have shown that this receptor is one of the key molecules that mediate GI cancers initiation, progression, metastasis, and therapy responses. Table 1 summarizes the modulatory effects of $\alpha 7$ nAChR in GI cancers regulation that have been investigated in previous studies. In the following sections, through a large number of such findings, we have detailed the most notable experiments that aimed to identify the role of gene expression and activation or blocking of $\alpha 7$ nAChR in the modulation of inflammatory and carcinogenic responses in different parts of the digestive system. In addition, we discuss the implication of this receptor in the interplay between the cholinergic neural pathway and immune-related activities that regulate different pathophysiological events during GI carcinogenesis.

3. $\alpha 7$ nAChR and colorectal cancer

Colorectal cancer (CRC) is one of the most serious worldwide health problems. This cancer causes at least seven-hundred thousand deaths

every year, which makes it one of the most deadly cancers in the world [38]. Treatment of CRC consists of surgery, chemotherapy, and radiotherapy, but due to the poor effectiveness of such available treatments, the need to develop novel strategies for this cancer therapy (such as gene therapy) is urgent [38, 39]. The participation of the $\alpha 7$ nAChR in this cancer pathogenesis has been investigated in previous studies. Activation of this receptor by nicotine contributes to CRC cell proliferation and inhibition of apoptosis [7, 40]. In these experiments, co-pretreatment of nicotine with $\alpha 7$ nAChR antagonist or reduced expression of this receptor by transfection of specific siRNA abolished nicotine-stimulated cell proliferation [7]. Thus, it is postulated that this receptor plays an important modulatory function in mediating the proliferative effect of nicotine on colon cancer cells. In addition to cancerous cells proliferation, blocking and struggle against the metastatic character of this cancer remain a serious obstacle, while the majority of patients that suffer from CRC eventually die due to the metastatic disease [41]. Related to these leading causes of death, it has been shown that nicotine and NNK through $\alpha 7$ nAChR-related signaling could enhance colon cancer cell migration and metastasis [42, 43]. Logically, both of the $\alpha 7$ nAChR antagonist and $\alpha 7$ nAChR siRNA methods have used to block the induced migration of CRC cells and the attenuated results confirmed that $\alpha 7$ nAChR played the

Table 1. Summary of some experimental studies indicating the modulatory effects of alpha7 nicotinic acetylcholine receptor ($\alpha 7$ nAChR) in gastrointestinal cancers regulation.

| Cancer type | Function | Commentary | References |
|-------------|---------------------------|---|------------|
| GC | Enhanced metastasis | Nicotine and its derived nitrosamine compound (i.e. NNK) through activation of $\alpha 7$ nAChR-related signaling pathways significantly enhanced gastric cancer (GC) metastasis. | [56, 57] |
| GC | Induced proliferation | Exposure to nicotine and NNK via $\alpha 7$ nAChR mechanisms significantly induced cell proliferation in the AGS cell line. | [58] |
| GC | Enhanced chemosensitivity | $\alpha 7$ nAChR enhanced the sensitivity of cancerous cells to chemotherapeutic reagents including taxanes and ixabepilone. | [6, 59] |
| GC | Enhanced chemoresistance | Targeted delivery of siRNA against $\alpha 7$ nAChR makes gastric tumor cells became more resistant to 5-FU treatment. | [73] |
| CRC | Induced proliferation | Nicotine promotes cell proliferation via $\alpha 7$ nAChR in human colorectal cancer (CRC) cell line. | [7] |
| CRC | Enhanced migration | Nicotine and tobacco-specific carcinogen (NNK) enhanced CRC cells migration through $\alpha 7$ nAChR-mediated mechanisms. | [42, 43] |
| CRC | Enhanced tumor growth | ACh itself via $\alpha 7$ nAChR serves as an important autocrine/paracrine growth factor in the human colon adenocarcinoma cell line HT-29. | [44] |
| CRC | Suppressed tumorigenesis | Nicotine suppresses acute colitis and CRC associated with chronic colitis in mice an effect that attenuated by the antagonist of $\alpha 7$ nAChR. | [49] |
| CRC | Suppressed metastasis | $\alpha 7$ nAChR in tumor-associated macrophages inhibits CRC metastasis in both animal model and LoVo human CRC cell line. | [52, 53] |
| PC | Induced proliferation | Nicotine and its nitrosated carcinogenic derivatives, promote cell proliferation of pancreatic cancer (PC) through activation of $\alpha 7$ nAChR. | [8] |
| PC | Induced CSCs renewal | Nicotine induces self-renewal of pancreatic cancer stem cells (CSCs) through $\alpha 7$ nAChR dependent mechanisms. | [92] |
| PC | Enhanced metastasis | Nicotine and cigarette-smoke promote the metastasis of PC via $\alpha 7$ nAChR downstream signaling cascades. | [93] |
| LC | Induced proliferation | Nicotine acts through $\alpha 7$ nAChR to stimulate the cholangiocyte proliferation in a xenograft mice model of liver cancer (LC). | [78] |
| LC | Increased carcinogenesis | NNK through increased expression of $\alpha 7$ nAChR caused hepatic damage and LC progression in an experimental animal model. | [79] |
| LC | Enhanced tumor growth | Nicotine-triggered $\alpha 7$ nAChR activation promotes both in vitro and in vivo tumor growth of HCC cells. | [81] |

important role in nicotine and NNK stimulated cell migration in CRC [42, 43]. Further to the nicotine and nitrosamines, Pettersson and coworkers have shown that ACh itself via the $\alpha 7$ nAChR serves as an important autocrine/paracrine growth factor in the human colon adenocarcinoma cell line HT-29 that positively contributes to the large intestine cancer progression [44].

In opposition to these tumor stimulatory effects, it also elucidated that, blockade of $\alpha 7$ nAChR may disrupt the homeostasis and negatively influence the normal physiologic functions such as intestine immunology. The host inflammatory responses that recruit distinct immune cells, cytokines, and other immune mediators have a consistent positive association with all steps of CRC, including initiation, promotion, progression, and metastasis [45]. This reciprocal interaction highlights the need for identification of intrinsic immunomodulatory mechanisms such as nicotinic anti-inflammatory pathways in the CRC microenvironment regulation. It has been proposed that vagal signaling downregulates GI inflammation, and the mechanism is thought to involve ACh binding to the $\alpha 7$ nAChR, as a modulator of CIP output [46]. In an animal model of colitis which induced by a chemical reagent (i.e. dextran sulfate sodium), vagotomy resulted in the increased level of NF- κ B activation and aggravates colitis in animals [47]. Also, some other investigations have shown in cancer subjects that high vagal activity, reported by heart rate variability (HRV) indexes correlates with advantageous prognosis in the patients and supports the beneficial role of vagal activity in tumor modulation [48]. In addition, it has been shown that nicotine suppresses acute colitis and colon cancer which associated with chronic colitis in mice [49]. In this study, a specific antagonist of $\alpha 7$ nAChR attenuated the inhibitory influences of nicotine on acute colitis [49]. These types of studies highlight the impact of anti-inflammatory mediators such as vagal anti-inflammatory signaling that mediated by $\alpha 7$ nAChR on the outcome of patients with CRC [50, 51]. In another experiment, $\alpha 7$ nAChR in tumor-associated macrophages inhibits CRC metastasis through its downstream intracellular signaling pathways which suggests that the expression of this receptor could be a valuable prognostic marker in CRC [52]. Similarly, Xiang and colleagues indicated that $\alpha 7$ nAChR inhibits cell invasion and metastasis of LoVo human CRC cells through downstream signaling pathways that initiated from this receptor [53]. It is important to mention that, for CRC, we have to make a clear distinction between inflammation-associated cancers in patients with inflammatory conditions, such as inflammatory bowel disease, and cancers developed following the adenoma-adenocarcinoma sequence, since the role of $\alpha 7$ nAChR in these two types of cancers may be different. Therefore, the anti-inflammatory and anti-cancer effects of $\alpha 7$ nAChR should be carefully considered in developing novel CRC therapeutic strategies.

4. $\alpha 7$ nAChR and gastric cancer

Gastric cancer (GC) is one of the top-ranked causes of cancer-related deaths. According to the global cancer statistics, approximately 950,000 new GC cases were reported, and due to poor effectiveness of available treatments, nearly 750,000 deaths occurred in the world during 2012 [2]. The incidence rate of this cancer demonstrates a strong etiologic association with smoking [54]. This habit emerged as a serious worldwide health problem and identified as an established environmental risk factor for GC development [54]. Smoking Tobacco contains various components that many of them have carcinogenic properties. Among these carcinogenic constituents of tobacco smoke, nicotine and nicotine-derived nitrosamines including 4-(methylnitrosamino)-1-(3-pyridyl)-1-butane (NNK), N-nitrosonornicotine (NNN) and N-nitroso diethylamine (DEN), have a great binding affinity to $\alpha 7$ nAChR and partly by binding to this receptor exert their carcinogenic functions [55]. These agents activate numerous cellular signaling pathways downstream of $\alpha 7$ nAChR, resulting in the stimulation of cancer-related properties.

Related to this fact, Wang and colleagues demonstrated that NNK through activation of $\alpha 7$ nAChR significantly enhanced human gastric

adenocarcinoma cells (AGS) migration [56]. In this study to investigate the mechanism of NNK-enhanced migration and verify that this effect is specifically mediated by $\alpha 7$ nAChR, the selective $\alpha 7$ nAChR antagonist, methyllycaconitine, was used in GC cells. In addition to this antagonist, $\alpha 7$ nAChR expression level by small interfering RNA (siRNA) technology was silenced [56]. The results determined that both $\alpha 7$ nAChR antagonist and $\alpha 7$ nAChR-siRNA inhibited NNK-induced migration [56]. Similarly, Lien et al. have shown that GC cells exposure to nicotine itself caused enhanced gastric cancer metastasis specifically by $\alpha 7$ nAChR, an effect suppressed by $\alpha 7$ nAChR-siRNA transfection [57]. In another experiment, both nicotine and NNK significantly induced cell proliferation in AGS cells that expressed $\alpha 7$ nAChR [58]. Treatment of these cells with $\alpha 7$ nAChR antagonist α -bungarotoxin blocked these proliferative effects [58]. In addition, the silencing of the $\alpha 7$ receptors expression level in AGS gastric tumor cells by siRNA enhances the sensitivity of these cancerous cells to taxanes treatment [59]. Similarly, $\alpha 7$ nAChR reduced the sensitivity of GC cells to ixabepilone as another chemotherapeutic reagent [6]. Therefore, the $\alpha 7$ nAChR expression level in patients with GC may be a good indicator of chemotherapy drugs sensitivity. Altogether, it can be concluded that $\alpha 7$ nAChR is an important regulator of gastric tumor cells proliferation, metastasis, and chemoresistance.

These studies highlighted the unfavorable consequences of exogenous activators of $\alpha 7$ nAChR in GC regulation. On another hand, vagus nerve activation and subsequently, ACh release from its terminals as major endogenous sources of ACh, by binding to $\alpha 7$ nAChR may play an additional stimulatory role in GC tumorigenesis [60]. It is delineated that the vagus nerve activation is necessary processes for gastric carcinogenesis [61]. Zhao and colleagues have shown that the vagus denervation of the stomach suppresses gastric tumorigenesis of the denervated portion in a mouse model of GC [62]. In this study, surgical removing of the vagus nerve (i.e. vagotomy) or pharmacological inhibition of vagus nerve signaling markedly reduced GC tumor incidence and progression [62]. This finding suggests that vagal denervation via reducing $\alpha 7$ nAChR activity might represent a feasible strategy for the control of gastric tumorigenesis.

In addition to exogenous (e.g. nicotine and NNK) or endogenous (e.g. ACh from the vagal terminal) activators of $\alpha 7$ nAChR and their carcinogenic functions, the GC etiology is closely linked to inflammation. This cancer is one of the best examples of inflammation-associated cancers in humans that is characterized by a slow stepwise evolution from superficial gastritis (inflammation of the lining of the stomach) to the final malignancy [63]. This sequence provides an excellent opportunity for the early detection and prevention of the underlying events preceding the development of the neoplasm [64]. However, the molecular mechanisms that regulate the transition from one step to the subsequent step are not still completely explained. In this cancer, chronic infection with *Helicobacter pylori* (*H. pylori*) is the strongest risk factor and ~90% of new cases of non-cardia GC worldwide attributed to this infection [65]. Infection with these bacteria causes an inflammatory response that enhances the infiltration of immune cells into the gastric mucosa [66]. Several studies clearly demonstrated the positive association between inflammation and pathogenesis of GC [67]. Recently, by RNA sequencing of gastritis and tumor tissues, it was indicated that the gene expression patterns of gastric tumors were similar to the gene expression of gastritis, indicating that these changes in the expression of tumors were induced by inflammation-dependent mechanisms [68]. Therefore, understanding the role of underlying molecular mechanisms involved in the modulation of inflammatory responses in GC pathogenesis is of particular importance.

Despite some studies claiming that the reduced activity of $\alpha 7$ nAChR may successfully block GC growth and progression; but at the same time, it might also block intrinsic mechanisms that are essential for inflammatory reaction control. Relating to this concept, in opposition to the benefits of vagal inhibition in GC progression, some others suggest that $\alpha 7$ nAChR signaling as well as high vagal activity predicts better cancer prognosis and also determined the increased risk of cancer mortality after

vagotomy [69, 70]. Fujita et al. have shown the enhancement of gastric carcinogenesis following vagotomy in an experimental animal model [71]. Similarly, promotion by vagotomy and increased incidence of GC has been shown in another experiment in rats [72]. As another evidence that serves the beneficial effects of $\alpha 7$ nAChR in GC, Chen et al. have shown that $\alpha 7$ nAChR enhanced the sensitivity of GC cells to 5-fluorouracil (5-FU) as a well-known chemotherapeutic drug [73]. In this study, when the expression of this receptor was knock-downed by a specific siRNA targeting $\alpha 7$ nAChR in AGS cells, these cells became more resistant to 5-FU treatment compared with the negative control cells [73]. Therefore, in opposition to the detrimental effects of $\alpha 7$ nAChR in GC development and therapy responses, due to its other beneficial biological effects, the clinical application of $\alpha 7$ nAChR antagonists in GC treatment needs to be taken into serious consideration.

5. $\alpha 7$ nAChR and liver cancer

Liver cancer (LC) is the second most common cancer in males, the seventh in females and the third largest cause of cancer-related deaths worldwide [74]. The primary LC is initiated in the liver cells and is divided into different types, among them, hepatocellular carcinoma (HCC) is the most frequent malignant liver tumor and accounts for around ninety percent of the primary LCs [75]. However, this cancer is one of the common causes of cancer-associated deaths, the existing therapeutic options are not fully effective and the major approach is preventing LC at high-risk patients [76]. In an important study, Sakata et al. have shown that in humans, the highest accumulation of $\alpha 7$ nAChR observed in the liver and this data demonstrated the importance of $\alpha 7$ receptor-related functions in this organ [77]. Similarly, it has been reported that $\alpha 7$ nAChR is one of the predominantly expressed nicotinic receptors in both LC cell lines and primary hepatoma cells, demonstrating that these receptors may play crucial roles in the regulation of LC development and progression [9]. This receptor is more highly expressed in LC cells compared with normal cells and serves as a predominant protein responsible for nicotine-mediated LC progression [78]. Previous studies have shown the involvement of $\alpha 7$ nAChR in liver pathological processes such as promoting carcinogenesis pathways in the liver. Aizawa et al. have shown that NNK as a high-affinity ligand of the $\alpha 7$ nAChR through increased expression and activation of this receptor caused to hepatic damage and HCC progression in an animal model [79]. Similarly, Martínez and colleagues study suggest that nicotine acts through $\alpha 7$ nAChR to stimulate the cholangiocyte proliferation in both *in vitro* and in xenograft mice model of LC [78]. In this context, the anti-tumor activity of acetylcholinesterase in HCC cells can also be justified by considering the fact that ACh is an important endogenous activator of $\alpha 7$ nAChR [80]. Also, Wan et al. revealed that nicotine-triggered $\alpha 7$ nAChR activation through signaling pathways associated with this receptor and promotes both *in vitro* and *in vivo* tumor growth of HCC cells [81].

There is numerous other evidence in the literature that displays the cancer-promoting effects of $\alpha 7$ nAChR, but on the other hand, there are also some controversial finding about its detrimental effects in the liver. The liver reciprocally communicates with the brain partly via the hepatic vagus branch (HVB) and $\alpha 7$ nAChR represents a unique molecular link between this parasympathetic system and immune- and cancer-related activities in the liver [82]. It is well-known that inflammatory processes are key players in LC development and progression [83]. Thus understanding the role of intrinsic inflammation controlling processes is of significant importance. In some animal models of experimental liver injury, cholinergic fibers are known to increase in the damaged area [84]. Also, the positive correlation between vagal activity that can be measured by different parameters of HRV and liver diseases has been well established in previous studies [85, 86]. These interactions may display the importance of CIP that exerts its protective effects through

$\alpha 7$ nAChR-dependent mechanisms in life-threatening conditions such as LC pathogenesis. In a most relevant study to this hypothesis which carried out by Hiramoto and coworkers, the protective role of the HVB activity against liver metastasis was founded in an experimental model in mice [87]. In addition, numerous findings demonstrated other beneficial effects of vagal firing in liver damage progression that exerts its protective effects via $\alpha 7$ nAChR based signaling pathways [88, 89].

6. $\alpha 7$ nAChR and pancreatic cancer

Pancreatic cancer (PC) is another leading cause of cancer-related deaths in the world, especially in developed countries [8]. Previous works have already established that PC cells express high levels of the $\alpha 7$ nAChR subunit [90]. Similar to other GI cancers, nicotine is a major risk factor for this cancer but the underlying mechanisms are inadequately explained [91]. It has been shown that nicotine and its nitrosated carcinogenic derivatives, promote cell proliferation of PC through $\alpha 7$ nAChR-mediated mechanisms [8]. In addition, PC stem cells stimulate tumor growth and metastasis, and it has been shown that nicotine induces self-renewal of these stem cells through $\alpha 7$ nAChR dependent mechanisms [92]. In this study chronic exposure to nicotine increased the protein expression of the $\alpha 7$ receptor thereby promoting the malignant potential of human PC cells [92]. Momi and colleagues determined that nicotine promotes metastasis of PC via $\alpha 7$ nAChR stimulation and subsequent activation of its downstream signaling cascades [93]. Also, chronic nicotine exposure inhibits the therapeutic effects of chemotherapy medications on PC in both *in vitro* and *in vivo* experiments [94]. By considering these findings, antagonizing the $\alpha 7$ nAChR emerged a new therapeutic strategy for combating tobacco-related cancers [95]. For example, it was shown that human secreted Ly-6/uPAR related protein-1 (SLURP-1), an endogenous peptide antagonist of $\alpha 7$ nAChR, is reported to have antiproliferative outcomes and significantly suppressed the development and progression of cancer-related characteristics in several cancer cell lines [96, 97]. Throm et al. indicated that SLURP-1 is a potential anti-malignant molecule that serves as an important tumor suppressor in PC [97]. In this study, it has been shown that patients with higher expression of $\alpha 7$ nAChR and its endogenous ligand SLURP-1 in resected pancreatic tumors had a better survival rate compared to those with a lower level of these proteins expression [97].

In addition, it has been shown that some inflammatory regulators such as NF- κ B activation and inflammatory cytokines production have a positive stimulatory role in PC associated properties such as proliferation and metastasis, thus inhibited activity of these inflammatory pathways by $\alpha 7$ nAChR-based pathways may represent a promising anti-cancer approach in PC therapy [98, 99]. Regarding the well-established anti-inflammatory functions of the nicotinic neural pathway, it has been determined that blocked activity of this pathway by vagotomy or pretreatment with pharmacological inhibitors of $\alpha 7$ nAChR resulted in enhanced severity of experimental pancreatitis and subsequent carcinogenesis [100]. The promoting effects of vagotomy on pancreatic carcinogenesis in animal models were demonstrated in previous studies [101, 102]. Recently, Partecke et al. demonstrated that sub-diaphragmatic vagotomy caused increased tumor growth and worsened survival rate in an animal PC model [103]. They concluded that these cancer-promoting effects of vagal denervation occur due to the removed anti-inflammatory processes and subsequently, over-production of inflammatory cytokines such as TNF- α in pancreatic tissue [103]. In normal physiological conditions, this excessive inflammatory process is restricted by CIP and $\alpha 7$ nAChR-based signaling pathway. It is important to note that unbalanced activity of this receptor following surgical or pharmacological blocking may cause undesired inflammation and cancer-promoting effects. Thus, the anti-inflammatory function of vagal signaling should carefully considered in some operative procedures especially in the upper GI tract that

in which partial or complete transection of the vagus nerve fibers usually will occur during the surgery.

7. Conclusion

This review highlighted the paradoxical effects of $\alpha 7$ nAChR-related signaling pathways on GI tract cancers development and progression. As described in the text, several effects that depended on the presence and correct function of $\alpha 7$ nAChR occurred in the regulation of GI cancers. Accumulated evidence demonstrated the cancer-promoting influences of this receptor in GI cancers, but, some other studies demonstrated beneficial effects for them in both cancer and non-cancer conditions. This evidence represents the excellent future perspectives of $\alpha 7$ nAChR-based strategies in cancer therapy. We consider the GI cancers as well-known inflammation-associated cancers and don't address just the damaging effects of $\alpha 7$ nAChR signaling in GI carcinogenesis, but somewhat concentrate on the anti-inflammatory functions of this receptor. This receptor exerts anti-inflammatory effects of vagal signaling and may serve the possible beneficial influences on GI cancers. There is an interesting hypothesis that supports the idea that the vagus nerve targets different organs and the brain via this neural pathway become informed about the preclinical neoplasms and suppressed them [104]. In accordance with this hypothesis, it should be mentioned that in some other non-cancer experiments the therapeutic benefits of $\alpha 7$ nAChR agonists were used to improve wound healing processes as well as in the treatment of neurodegenerative illnesses such as Parkinson's and Alzheimer's diseases [105]. However, the impacts of other regulatory mechanisms such as single nucleotide polymorphism and regulatory microRNAs affecting $\alpha 7$ functions throughout the body should be investigated in future studies [106, 107]. Further insights regarding the therapeutic potential of $\alpha 7$ nAChR that implicated in pathological conditions are discussed in a recent review and are beyond the scope of this manuscript to mention all of them [108]. The pathogenesis of GI cancers are complex disorders and not mediated by a single factor. Regarding this fact, the $\alpha 7$ nAChR inhibitors have not been widely used for cancer prevention and treatment because of their possible complications. Hence, successful treatment may require comprehensive strategies to modulate several, rather than just a single underlying factor. On another hand, it is important to investigate intracellular mechanisms by which $\alpha 7$ nAChR regulates GI tumor cells properties. Altogether, $\alpha 7$ nAChRs are expressed in a variety of tissues including GI and hepatic cells and there is evidence to suggest that activation of these receptors has a paradoxical effect on the progression of cancer in the GI tract. On one hand, it suppresses inflammation and makes cellular microenvironment less favorable for tumorigenesis and on the other hand, activation of these receptors in cancerous cells promotes cell proliferation and migration. Therefore, in contrast to the benefits of their blockade in cancer therapy, the unique anti-inflammatory properties of them should be carefully considered in the designing of novel pharmacological or surgical therapeutic strategies.

Declarations

Author contribution statement

All authors listed have significantly contributed to the development and the writing of this article.

Funding statement

This work was supported by the National Institute for Medical Research Development (NIMAD), Iran (project no. 972536) and Tabriz University of Medical Sciences, Tabriz, Iran (project no. 59256).

Competing interest statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

References

- [1] P. Grierson, K.H. Lim, M. Amin, Immunotherapy in gastrointestinal cancers, *J. Gastrointest. Oncol.* 8 (2017) 474–484.
- [2] L.A. Torre, F. Bray, R.L. Siegel, J. Ferlay, J. Lortet-tieulent, A. Jemal, Global cancer statistics, 2012, *CA A Cancer J. Clin.* 65 (2015) 87–108.
- [3] A. Mohammadi, B. Mansoori, P. Savadi, V. Khaze, M. Minouei, N.A.J. McMillan, S. Hallaj-Nezhadi, B. Baradarani, Targeting of high mobility group A2 by small interfering RNA-loaded nanoliposome-induced apoptosis and migration inhibition in gastrointestinal cancer cells, *J. Cell. Biochem.* (2018).
- [4] Y. Zhao, The oncogenic functions of nicotinic acetylcholine receptors, *J. Oncol.* 2016 (2016).
- [5] H.M. Schuller, Regulatory role of the alpha7nAChR in cancer, *Curr. Drug Targets* 13 (2012) 680–687.
- [6] C.C. Tu, C.Y. Huang, W.L. Cheng, C.S. Hung, Y.J. Chang, P.L. Wei, Silencing A7-nAChR levels increases the sensitivity of gastric cancer cells to ixabepilone treatment, *Tumor Biol.* 37 (2016) 9493–9501.
- [7] H.P.S. Wong, L. Yu, E.K.Y. Lam, E.K.K. Tai, W.K.K. Wu, C.H. Cho, Nicotine promotes cell proliferation via $\alpha 7$ -nicotinic acetylcholine receptor and catecholamine-synthesizing enzymes-mediated pathway in human colon adenocarcinoma HT-29 cells, *Toxicol. Appl. Pharmacol.* 221 (2007) 261–267.
- [8] M.H. Al-Wadei, H.A.N. Al-Wadei, H.M. Schuller, Pancreatic cancer cells and normal pancreatic duct epithelial cells express an autocrine catecholamine loop that is activated by nicotinic acetylcholine receptors 3, 5, and 7, *Mol. Canc. Res.* 10 (2012) 239–249.
- [9] L. Zhang, L.L. Wu, H.B. Huan, X.J. Chen, X.D. Wen, D.P. Yang, F. Xia, Sympathetic and parasympathetic innervation in hepatocellular carcinoma, *Neoplasma* 64 (2017) 840–846.
- [10] M.J. De Rosa, M.D.C. Esandi, A. Garelli, D. Rayes, C. Bouzat, Relationship between $\alpha 7$ nAChR and apoptosis in human lymphocytes, *J. Neuroimmunol.* 160 (2005) 154–161.
- [11] H. Yoshikawa, M. Kurokawa, N. Ozaki, K. Nara, K. Atou, E. Takada, H. Kamochi, N. Suzuki, Nicotine inhibits the production of proinflammatory mediators in human monocytes by suppression of I-kB phosphorylation and nuclear factor-kB transcriptional activity through nicotinic acetylcholine receptor $\alpha 7$, *Clin. Exp. Immunol.* 146 (2006) 116–123.
- [12] W.J. de Jonge, E.P. van der Zanden, F.O. The, M.F. Bijlsma, D.J. van Westerloo, R.J. Bennink, H. Berthoud, S. Uematsu, S. Akira, R.M. van den Wijngaard, G.E. Boeckxstaens, Stimulation of the vagus nerve attenuates macrophage activation by activating the Jak2-STAT3 signaling pathway, *Nat. Immunol.* 6 (2005) 844–851.
- [13] R.D. Eglington, K.C. Brown, P. Dasgupta, Nicotinic acetylcholine receptors in cancer: multiple roles in proliferation and inhibition of apoptosis, *Trends Pharmacol. Sci.* 29 (2008) 151–158.
- [14] S. Singh, S. Pillai, S. Chellappan, Nicotinic acetylcholine receptor signaling in tumor growth and metastasis, *J. Oncol.* 2011 (2011) 1–11.
- [15] S. Pillai, S. Chellappan, Alpha7 nicotinic acetylcholine receptor subunit in angiogenesis and epithelial to mesenchymal transition, *Curr. Drug Targets* 13 (2012) 671–679.
- [16] P.H. Nguyen, Y. Toucheuf, T. Durand, P. Aubert, E. Duchalais, S. Bruley des Varannes, C. Veron, M. Neunlist, T. Matysiak-Budnik, Acetylcholine induces stem cell properties of gastric cancer cells of diffuse type, *Tumor Biol.* 40 (2018) 1–9.
- [17] H. Wang, M. Yu, M. Ochani, C.A. Amella, M. Tanovic, S. Susarla, J.H. Li, H. Wang, H. Yang, L. Ulloa, Y. Al-Abed, C.J. Czura, K.J. Tracey, Nicotinic acetylcholine receptor alpha7 subunit is an essential regulator of inflammation, *Nature* 421 (2003) 384–388.
- [18] M. Bencherif, P.M. Lippiello, R. Lucas, M.B. Marrero, Alpha7 nicotinic receptors as novel therapeutic targets for inflammation-based diseases, *Cell. Mol. Life Sci.* 68 (2011) 931–949.
- [19] B.B. Aggarwal, B. Sung, S.C. Gupta, *Inflammation and Cancer*, Springer Basel, Basel, 2014.
- [20] K. Echizen, H. Oshima, M. Nakayama, M. Oshima, The inflammatory microenvironment that promotes gastrointestinal cancer development and invasion, *Adv. Biol. Regul.* 68 (2018) 39–45.
- [21] K.A.F. Pennel, J.H. Park, D.C. McMillan, A.K. Roseweir, J. Edwards, Signal interaction between the tumour and inflammatory cells in patients with gastrointestinal cancer: implications for treatment, *Cell. Signal.* 54 (2019) 81–90.
- [22] H. Wang, C.H. Cho, Effect of NF-kappaB signaling on apoptosis in chronic inflammation-associated carcinogenesis, *Curr. Cancer Drug Targets* 10 (2010) 593–599.
- [23] J.P. Waters, J.S. Poher, J.R. Bradley, Tumour necrosis factor and cancer, *J. Pathol.* 230 (2013) 241–248.
- [24] J. Candido, T. Hagemann, Cancer-related inflammation, *J. Clin. Immunol.* 33 (2013).
- [25] S.S. Husain, I.L. Szabo, A.S. Tamawski, NSAID inhibition of GI cancer growth: clinical implications and molecular mechanisms of action, *Am. J. Gastroenterol.* 97 (2002) 542–553.
- [26] M. Letícia de Castro Barbosa, R. Alves da Conceicao, A. Guerra Manssour Fraga, B. Dias Camarinha, G. Cristina de Carvalho Silva, A. Gilcer Ferreira Lima, E. Azevedo Cardoso, V. de Oliveira Freitas Lione, NF- κ B signaling pathway

- inhibitors as anticancer drug candidates, anticancer, *Agents Med. Chem.* 17 (2017) 483–490.
- [27] W.J. de Jonge, L. Ulloa, The alpha7 nicotinic acetylcholine receptor as a pharmacological target for inflammation, *Br. J. Pharmacol.* 151 (2007) 915–929.
- [28] M. Rosas-Ballina, P.S. Olofsson, M. Ochani, S.I. Valdés-Ferrer, Y.A. Levine, C. Reardon, M.W. Tusche, V.A. Pavlov, U. Andersson, S. Chavan, T.W. Mak, K.J. Tracey, Acetylcholine-synthesizing T cells relay neural signals in a vagus nerve circuit, *Science* 334 (2011) 98–101.
- [29] L. V Borovikova, S. Ivanova, M. Zhang, H. Yang, G.I. Botchkina, L.R. Watkins, H. Wang, N. Abumrad, J.W. Eaton, K.J. Tracey, Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin, *Nature* 405 (2000) 458–462.
- [30] V.A. Pavlov, K.J. Tracey, Neural regulation of immunity: molecular mechanisms and clinical translation, *Nat. Neurosci.* 20 (2017) 156–166.
- [31] E. Meroni, N. Stakenborg, M.F. Viola, G.E. Boeckxstaens, Intestinal macrophages and their interaction with the enteric nervous system in health and inflammatory bowel disease, *Acta Physiol.* (2018) 1–11.
- [32] A. Kiss, Y. Tratsiakovich, A. Mahdi, J. Yang, A.T. Gonon, B.K. Podesser, J. Pernow, Vagal nerve stimulation reduces infarct size via a mechanism involving the alpha-7 nicotinic acetylcholine receptor and downregulation of cardiac and vascular arginase, *Acta Physiol.* 221 (2017) 174–181.
- [33] W.R. Parrish, M. Rosas-Ballina, M. Gallowitsch-Puerta, M. Ochani, K. Ochani, L.-H. Yang, L. Hudson, X. Lin, N. Patel, S.M. Johnson, S. Chavan, R.S. Goldstein, C.J. Czura, E.J. Miller, Y. Al-Abed, K.J. Tracey, V.A. Pavlov, Modulation of TNF release by choline requires alpha7 subunit nicotinic acetylcholine receptor-mediated signaling, *Mol. Med.* 14 (2008) 567–574.
- [34] K. Chen, Y. Sun, Y. Diao, L. Ji, D. Song, T. Zhang, α7 nicotinic acetylcholine receptor agonist inhibits the damage of rat hippocampal neurons by TLR4/Myd88/NF-κB signaling pathway during cardiopulmonary bypass, *Mol. Med. Rep.* 16 (2017) 4770–4776.
- [35] K. Hajiasgharzadeh, S. Sadigh-Eteghad, B. Mansoori, A. Mokhtarzadeh, D. Shafebandi, M.A. Doustvandi, Z. Asadzadeh, B. Baradaran, Alpha7 nicotinic acetylcholine receptors in lung inflammation and carcinogenesis: friends or foes? *J. Cell. Physiol.* (2019).
- [36] Z. Zhou, Y. Liu, X. Chen, F. Li, X. Tong, Y. Ding, C. Tang, Treatment of experimental non-alcoholic steatohepatitis by targeting α7 nicotinic acetylcholine receptor-mediated inflammatory responses in mice, *Mol. Med. Rep.* 12 (2015) 6925–6931.
- [37] E. Reijmen, L. Vannucci, M. De Couck, J. De Grève, Y. Gidron, Therapeutic potential of the vagus nerve in cancer, *Immunol. Lett.* 202 (2018) 38–43.
- [38] H. Brody, Colorectal cancer, *Nature* 521 (2015). S1–S1.
- [39] S. Davudian, N. Shahari, T. Kazemi, B. Mansoori, S. Salehi, A. Mohammadi, D. Shafebandi, V.K. Shahgoli, M. Asadi, B. Baradaran, BACH1 silencing by siRNA inhibits migration of HT-29 colon cancer cells through reduction of metastasis-related genes, *Biomed. Pharmacother.* 84 (2016) 191–198.
- [40] S. Dinicola, V. Morini, P. Coluccia, S. Proietti, F. D'Anselmi, A. Pasqualato, M.G. Masiello, A. Palombo, G. De Toma, M. Bizzarri, A. Cucina, Nicotine increases survival in human colon cancer cells treated with chemotherapeutic drugs, *Toxicol. Vitro* 27 (2013) 2256–2263.
- [41] A. Mohammadi, B. Mansoori, P.C. Baradaran, S.C. Baradaran, B. Baradaran, Anacyclus pyrethrum extract exerts anticancer activities on the human colorectal cancer cell line (HCT) by targeting apoptosis, metastasis and cell cycle arrest, *J. Gastrointest. Canc.* 48 (2017) 333–340.
- [42] P.L. Wei, L.J. Kuo, M. Te Huang, W.C. Ting, Y.S. Ho, W. Wang, J. An, Y.J. Chang, Nicotine enhances colon cancer cell migration by induction of fibronectin, *Ann. Surg. Oncol.* 18 (2011) 1782–1790.
- [43] P.-L. Wei, Y.-J. Chang, Y.-S. Ho, C.-H. Lee, Y.-Y. Yang, J. An, S.-Y. Lin, Tobacco-specific carcinogen enhances colon cancer cell migration through α7-nicotinic acetylcholine receptor, *Ann. Surg.* 249 (2009) 978–985.
- [44] A. Pettersson, L. Nilsson, G. Nylund, A. Khorram-Manesh, S. Nordgren, D.S. Delbro, Is acetylcholine an autocrine/paracrine growth factor via the nicotinic α7-receptor subtype in the human colon cancer cell line HT-29? *Eur. J. Pharmacol.* 609 (2009) 27–33.
- [45] J. Terzić, S. Grivennikov, E. Karin, M. Karin, Inflammation and colon cancer, *Gastroenterology* 138 (2010) 2101–2114.
- [46] V.A. Pavlov, K.J. Tracey, Neural circuitry and immunity, *Immunol. Res.* 63 (2015) 38–57.
- [47] C. O'Mahony, H. van der Kleij, J. Bienenstock, F. Shanahan, L. O'Mahony, Loss of vagal anti-inflammatory effect: in vivo visualization and adoptive transfer, *AJP Regul. Integr. Comp. Physiol.* 297 (2009) R1118–R1126.
- [48] C. Mouton, A. Ronson, D. Razavi, F. Delhaye, N. Kupper, M. Paesmans, M. Moreau, J.M. Nogaret, A. Hendlisz, Y. Gidron, The relationship between heart rate variability and time-course of carcinoembryonic antigen in colorectal cancer, *Auton. Neurosci. Basic Clin.* 166 (2012) 96–99.
- [49] S. Hayashi, T. Hamada, S.F. Zaidi, M. Oshiro, J. Lee, T. Yamamoto, Y. Ishii, M. Sasahara, M. Kadokawa, Nicotine suppresses acute colitis and colonic tumorigenesis associated with chronic colitis in mice, *Am. J. Physiol. Gastrointest. Liver Physiol.* 307 (2014) G968–G978.
- [50] J.H. Park, D.C. McMillan, P.G. Horgan, C.S. Roxburgh, The impact of anti-inflammatory agents on the outcome of patients with colorectal cancer, *Cancer Treat. Rev.* 40 (2014) 68–77.
- [51] M. Pelissier-Rota, M. Lainé, B. Ducarouge, B. Bonaz, M. Jacquier-Sarlin, Role of cholinergic receptors in colorectal Cancer : potential therapeutic implications of vagus nerve Stimulation ? *J. Canc. Ther.* 2013 (2013) 1116–1131.
- [52] R. Fei, Y. Zhang, S. Wang, T. Xiang, W. Chen, α7 nicotinic acetylcholine receptor in tumor-associated macrophages inhibits colorectal cancer metastasis through the JAK2/STAT3 signaling pathway, *Oncol. Rep.* 38 (2017) 2619–2628.
- [53] T. Xiang, F. Yu, R. Fei, J. Qian, W. Chen, CHRNA7 inhibits cell invasion and metastasis of LoVo human colorectal cancer cells through PI3K/Akt signaling, *Oncol. Rep.* 35 (2016) 999–1005.
- [54] V.Y. Shin, C.-H. Cho, Nicotine and gastric cancer, *Alcohol* 35 (2005) 259–264.
- [55] H.M. Schuller, Nitrosamines as nicotinic receptor ligands, *Life Sci.* 80 (2007) 2274–2280.
- [56] W. Wang, H. Chin-Sheng, L.J. Kuo, P.L. Wei, Y.C. Lien, F.Y. Lin, H.H. Liu, Y.S. Ho, C.H. Wu, Y.J. Chang, NNK enhances cell migration through α7-nicotinic acetylcholine receptor accompanied by increased of fibronectin expression in gastric cancer, *Ann. Surg. Oncol.* 19 (2012) 580–588.
- [57] Y.C. Lien, W. Wang, L.J. Kuo, J.J. Liu, P.L. Wei, Y.S. Ho, W.C. Ting, C.H. Wu, Y.J. Chang, Nicotine promotes cell migration through alpha7 nicotinic acetylcholine receptor in gastric cancer cells, *Ann. Surg. Oncol.* 18 (2011) 2671–2679.
- [58] V.Y. Shin, H.C. Jin, E.K.O. Ng, J. Yu, W.K. Leung, C.H. Cho, J.J.Y. Sung, Nicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone induce cyclooxygenase-2 activity in human gastric cancer cells: involvement of nicotinic acetylcholine receptor (nAChR) and β-adrenergic receptor signaling pathways, *Toxicol. Appl. Pharmacol.* 233 (2008) 254–261.
- [59] C.C. Tu, C.Y. Huang, W.L. Cheng, C.S. Hung, B. Uyanga, P.L. Wei, Y.J. Chang, The α7-nicotinic acetylcholine receptor mediates the sensitivity of gastric cancer cells to taxanes, *Tumor Biol.* 37 (2016) 4421–4428.
- [60] H.-L. Rabben, C.-M. Zhao, Y. Hayakawa, T.C. Wang, D. Chen, Vagotomy and gastric tumorigenesis, *Curr. Neuropharmacol.* 14 (2016) 967–972.
- [61] Vagal innervation is necessary for gastric tumorigenesis, *Canc. Discov.* 4 (2014) OF15–OF15.
- [62] C.-M. Zhao, Y. Hayakawa, Y. Kodama, S. Muthupalan, C.B. Westphalen, G.T. Andersen, A. Flatberg, H. Johannessen, R.A. Friedman, B.W. Renz, A.K. Sandvik, V. Beisvag, H. Tomita, A. Hara, M. Quante, Z. Li, M.D. Gershon, K. Kaneko, J.G. Fox, T.C. Wang, D. Chen, Denervation suppresses gastric tumorigenesis, *Sci. Transl. Med.* 6 (2014) 250ra115–250ra115.
- [63] K. Oguma, H. Oshima, M. Oshima, Inflammation, tumor necrosis factor and Wnt promotion in gastric cancer development, *Future Oncol.* 6 (2010) 515–526.
- [64] C. Luigiano, F. Ferrara, S. Fagoonee, C. Fabbri, V. Cennamo, C. Virgilio, M. Bassi, C. Favara, C. Morace, P. Consolo, R. Pellicano, Is Helicobacter pylori the infectious target to prevent gastric cancer? an interdisciplinary point of view, *Infect. Disord. - Drug Targets* 12 (2012) 340–345.
- [65] C. De Martel, J. Ferlay, S. Franceschi, J. Vignat, F. Bray, D. Forman, M. Plummer, Global burden of cancers attributable to infections in 2008: a review and synthetic analysis, *Lancet Oncol.* 13 (2012) 607–615.
- [66] S.L. Servetas, D.R. Bridge, D.S. Merrell, Molecular Mechanisms of Gastric Cancer Initiation and Progression by Helicobacter pylori, *Curr. Opin. Infect. Dis.* 29 (2016) 304–310.
- [67] P. Ernst, Review article: the role of inflammation in the pathogenesis of gastric cancer, *Aliment. Pharmacol. Ther.* 13 (1999) 13–18.
- [68] K. Echizen, O. Hirose, Y. Maeda, M. Oshima, Inflammation in gastric cancer: interplay of the COX-2/prostaglandin E2 and Toll-like receptor/MyD88 pathways, *Canc. Sci.* 107 (2016) 391–397.
- [69] M. De Couck, R. Caers, D. Spiegel, Y. Gidron, The role of the vagus nerve in cancer prognosis: a systematic and a comprehensive review, *J. Oncol.* 2018 (2018).
- [70] C.P. Caygill, R.L. Knowles, R. Hall, Increased risk of cancer mortality after vagotomy for peptic ulcer: a preliminary analysis, *Eur. J. Canc. Prev.* 1 (1991) 35–37.
- [71] M. Fujita, M. Takami, M. Usugane, S. NamDei, T. Taguchi, Enhancement of gastric carcinogenesis in dogs given N-methyl-N'-nitro-N-nitrosoguanidine following vagotomy, *Cancer Res.* 39 (1979) 811–816.
- [72] M. Tatsuta, H. Yamamura, H. Iishi, M. Ichii, S. Noguchi, M. Baba, H. Taniguchi, Promotion by vagotomy of gastric carcinogenesis induced by N-methyl-N'-nitro-N-nitrosoguanidine in Wistar rats, *Cancer Res.* 45 (1985) 194–197.
- [73] W.Y. Chen, C.Y. Huang, W.L. Cheng, C.S. Hung, M. Te Huang, C.J. Tai, Y.N. Liu, C.L. Chen, Y.J. Chang, Alpha 7-nicotinic acetylcholine receptor mediates the sensitivity of gastric cancer cells to 5-flourouracil, *Tumor Biol.* 36 (2015) 9537–9544.
- [74] K. Hajiasgharzadeh, M.H. Somi, D. Shafebandi, A. Mokhtarzadeh, B. Baradaran, Small interfering RNA-mediated gene suppression as a therapeutic intervention in hepatocellular carcinoma, *J. Cell. Physiol.* 234 (2019) 3263–3276.
- [75] E.U. Cidon, Systemic treatment of hepatocellular carcinoma: past, present and future, *World J. Hepatol.* 9 (2017) 797–807.
- [76] B. Yousefi, M. Darabi, B. Baradaran, M.S. Khaniani, M. Rahbani, M. Darabi, S. Fayezi, A. Mehdizadeh, N. Salianni, M. Shaaker, Inhibition of MEK/ERK1/2 signaling affects the fatty acid composition of HepG2 human hepatic cell line, *Bioimpacts* 2 (2012) 145–150.
- [77] M. Sakata, J. Wu, J. Toyohara, K. Oda, M. Ishikawa, K. Ishii, K. Hashimoto, K. Ishiwata, Biodistribution and radiation dosimetry of the (alpha)7 nicotinic acetylcholine receptor ligand [11C]CHIBA-1001 in humans, *Nucl. Med. Biol.* 38 (2011) 443–448.
- [78] A.K. Martinez, K. Jensen, C. Hall, A. O'Brien, L. Ehrlich, T. White, F. Meng, T. Zhou, J. Greene, F. Bernuzzi, P. Invernizzi, D.E. Dostal, T. Lairmore, G. Alpini, S.S. Glaser, Nicotine promotes cholangiocarcinoma growth in xenograft mice, *Am. J. Pathol.* 187 (2017) 1093–1105.
- [79] K. Aizawa, C. Liu, S. Tang, S. Veeramachaneni, K.-Q. Hu, D.E. Smith, X.-D. Wang, Tobacco carcinogen induces both lung cancer and non-alcoholic steatohepatitis

- and hepatocellular carcinomas in ferrets which can be attenuated by lycopene supplementation, *Int. J. Canc.* 139 (2016) 1171–1181.
- [80] B. Pérez-Aguilar, C.J. Vidal, G. Palomec, F. García-Dolores, M.C. Gutiérrez-Ruiz, L. Bucio, J.L. Gómez-Olivares, L.E. Gómez-Quiroz, Acetylcholinesterase is associated with a decrease in cell proliferation of hepatocellular carcinoma cells, *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1852 (2015) 1380–1387.
- [81] C. Wan, M. Wu, S. Zhang, Y. Chen, C. Lu, α 7nAChR-mediated recruitment of PP1 γ promotes TRAF6/NF- κ B cascade to facilitate the progression of Hepatocellular Carcinoma, *Mol. Carcinog.* 57 (2018) 1626–1639.
- [82] K. Hajiasgharzadeh, S.M. Tavangar, M. Javan, A.R. Dehpour, A.R. Mani, Does hepatic vagus nerve modulate the progression of biliary fibrosis in rats? *Auton. Neurosci.* 185 (2014) 67–75.
- [83] B. Sun, M. Karin, Inflammation and liver tumorigenesis, *Front. Med.* 7 (2013) 242–254.
- [84] H. Akiyoshi, T. Terada, Mast cell, myofibroblast and nerve terminal complexes in carbon tetrachloride-induced cirrhotic rat livers, *J. Hepatol.* 29 (1998) 112–119.
- [85] A.R. Mani, S. Montagnese, C.D. Jackson, C.W. Jenkins, I.M. Head, R.C. Stephens, K.P. Moore, M.Y. Morgan, Decreased heart rate variability in patients with cirrhosis relates to the presence and degree of hepatic encephalopathy, *Am. J. Physiol. Gastrointest. Liver Physiol.* 296 (2009) G330–G338.
- [86] A.S. Bhogal, M. De Rui, D. Pavanello, I. El-Azizi, S. Rowshan, P. Amadio, S. Montagnese, A.R. Mani, Which heart rate variability index is an independent predictor of mortality in cirrhosis, *Dig. Liver Dis.* (2018).
- [87] T. Hiramoto, K. Yoshihara, Y. Asano, N. Sudo, Protective role of the hepatic vagus nerve against liver metastasis in mice, *Neuroimmunomodulation* 24 (2018) 341–347.
- [88] T. Nishio, K. Taura, K. Iwaisako, Y. Koyama, K. Tanabe, G. Yamamoto, Y. Okuda, Y. Ikeno, K. Yoshino, Y. Kasai, M. Okuno, S. Seo, T. Sakurai, M. Asagiri, E. Hatano, S. Uemoto, Hepatic vagus nerve regulates Kupffer cell activation via α 7 nicotinic acetylcholine receptor in nonalcoholic steatohepatitis, *J. Gastroenterol.* 52 (2017) 965–976.
- [89] M. Ni, H. Fu, F. Huang, T. Zhao, J. Chen, D. Li, F. Shen, D. Ph, J. Chen, D. Li, D. Ph, F. Shen, D. Ph, Vagus nerve attenuates hepatocyte apoptosis upon Ischemia-Reperfusion via α 7 nicotinic acetylcholine receptor on Kupffer cells in mice, *Anesthesiology* 125 (2016) 1005–1016.
- [90] J.G. Trevino, S. Pillai, S. Kunigal, S. Singh, W.J. Fulp, B.A. Centeno, S.P. Chellappan, Nicotine induces inhibitor of differentiation-1 in a src-dependent pathway promoting metastasis and chemoresistance in pancreatic adenocarcinoma, *Neoplasia* 14 (2012) 1102–1114.
- [91] R. Lochan, H.L. Reeves, A.K. Daly, R.M. Charnley, The role of tobacco-derived carcinogens in pancreas cancer, *ISRN Oncol.* 2011 (2011) 1–9.
- [92] M.H. Al-Wadei, J. Banerjee, H.A.N. Al-Wadei, H.M. Schuller, Nicotine induces self-renewal of pancreatic cancer stem cells via neurotransmitter-driven activation of sonic hedgehog signalling, *Eur. J. Canc.* 52 (2016) 188–196.
- [93] N. Momi, M.P. PonnuSamy, S. Kaur, S. Rachagani, S.S. Kunigal, S. Chellappan, M.M. Ouellette, S.K. Batra, Nicotine/cigarette smoke promotes metastasis of pancreatic cancer through α 7nAChR-mediated MUC4 upregulation, *Oncogene* 32 (2013) 1384–1395.
- [94] J. Banerjee, H.A.N. Al-Wadei, H.M. Schuller, Chronic nicotine inhibits the therapeutic effects of gemcitabine on pancreatic cancer in vitro and in mouse xenografts, *Eur. J. Canc.* 49 (2013) 1152–1158.
- [95] C.-H. Wu, C.-H. Lee, Y.-S. Ho, Nicotinic acetylcholine receptor-based blockade: applications of molecular targets for cancer therapy, *Clin. Canc. Res.* 17 (2011) 3533–3541.
- [96] E.N. Lyukmanova, M.A. Shulepko, M.L. Bychkov, Z.O. Shenkarev, A.S. Paramonov, A.O. Chugunov, A.S. Arseniev, D.A. Dolgikh, M.P. Kirpichnikov, Human SLURP-1 and SLURP-2 proteins Acting on nicotinic acetylcholine receptors reduce proliferation of human colorectal adenocarcinoma HT-29 cells, *Acta Nat.* 6 (2014) 60–66.
- [97] V.M. Throm, D. Männle, T. Giese, A.S. Bauer, M.M. Gaida, J. Kopitz, T. Bruckner, K. Plaschke, S.P. Grekova, K. Felix, T. Hackert, N.A. Giese, O. Strobel, V.M. Throm, D. Männle, T. Giese, A.S. Bauer, M.M. Gaida, J. Kopitz, T. Bruckner, K. Plaschke, S.P. Grekova, K. Felix, T. Hackert, N.A. Giese, O. Strobel, Endogenous CHRNA7-ligand SLURP1 as a potential tumor suppressor and anti-nicotinic factor in pancreatic cancer, *Oncotarget* 9 (2018) 11734–11751.
- [98] Q. Li, G. Yang, M. Feng, S. Zheng, Z. Cao, J. Qiu, L. You, L. Zheng, Y. Hu, T. Zhang, Y. Zhao, NF- κ B in pancreatic cancer: its key role in chemoresistance, *Canc. Lett.* 421 (2018) 127–134.
- [99] S. Farajzadeh Valilou, M. Keshavarz-Fathi, N. Silvestris, A. Argentiero, N. Rezaei, The role of inflammatory cytokines and tumor associated macrophages (TAMs) in microenvironment of pancreatic cancer, *Cytokine Growth Factor Rev* 39 (2018) 46–61.
- [100] D.J. van Westerloo, I.A. Giebelan, S. Florquin, M.J. Bruno, G.J. LaRosa, L. Ulloa, K.J. Tracey, T. van der Poll, The vagus nerve and nicotinic receptors modulate experimental pancreatitis severity in mice, *Gastroenterology* 130 (2006) 1822–1830.
- [101] A. Rivenson, D. Hoffmann, B. Prokopczyk, S. Amin, S.S. Hecht, Induction of lung and exocrine pancreas tumors in F344 rats by tobacco-specific and areca- derived N-nitrosamines, *Cancer Res.* 48 (1988) 6912–6917.
- [102] T. Ogawa, T. Makino, K. Mizumoto, F. Nakayama, Promoting effect of truncal vagotomy on pancreatic carcinogenesis initiated with N-nitrosobis(2-oxopropyl) amine in syrian golden hamsters, *Carcinogenesis* 12 (1991) 1227–1230.
- [103] L. Ivo Partecke, A. Käding, D. Nguyen Trung, S. Diedrich, M. Senderl, F. Weiss, J.-P. Kühn, J. Mayerle, K. Beyer, W. von Bernstorff, C.-D. Heidecke, W. Kefler, Subdiaphragmatic vagotomy promotes tumor growth and reduces survival via TNF α in a murine pancreatic cancer model, *Oncotarget* 8 (2017) 22501–22512.
- [104] Y. Gidron, H. Perry, M. Glennie, Does the vagus nerve inform the brain about preclinical tumours and modulate them? *Lancet Oncol.* 6 (2005) 245–248.
- [105] S. Sadigh-Eteghad, J. Mahmoudi, S. Babri, M. Talebi, Effect of alpha-7 nicotinic acetylcholine receptor activation on beta-amyloid induced recognition memory impairment. Possible role of neurovascular function, *Acta Cir. Bras.* 30 (2015) 736–742.
- [106] M. Asadi, D. Shafehbandi, A. Zarintan, N. Pedram, B. Baradaran, V. Zafari, M. Shirmohamadi, S. Hashemzadeh, TP53 Gene Pro72Arg (rs1042522) single nucleotide polymorphism as not a risk factor for colorectal cancer in the Iranian Azari population, *Asian Pac. J. Cancer Prev.* APJCP 18 (2017) 3423–3427.
- [107] N. HosseiniAhli, M. Aghapour, P.H.G. Duijf, B. Baradaran, Treating cancer with microRNA replacement therapy: a literature review, *J. Cell. Physiol.* (2018) 1–15.
- [108] D. Bertrand, C.-H.L. Lee, D. Flood, F. Marger, D. Donnelly-Roberts, Therapeutic potential of 7 nicotinic acetylcholine receptors, *Pharmacol. Rev.* 67 (2015) 1025–1073.