


WHAT'S NEW IN INTENSIVE CARE



Adjuvant therapies in critical care: steroids in community-acquired pneumonia

Antoni Torres^{1,2,3*} , Miquel Ferrer^{1,2,3} and Michael S. Niederman⁴

© 2017 Springer-Verlag GmbH Germany and ESICM

Severe community-acquired pneumonia (CAP) is present in up to 19% of hospitalized patients with CAP and still carries a high morbidity and mortality. A French multicentre study of severe pneumococcal CAP patients admitted to intensive care units (ICU) reported an overall mortality rate of 29% [1]. The high mortality of severe CAP occurs despite the fact that the majority of patients receive an early and adequate antibiotic treatment. This is probably due, in part, to an imbalanced and disproportionate local and systemic inflammatory response that contributes to impairment of gas exchange, sepsis and end-organ dysfunction.

In humans, intravenous corticosteroids attenuate the local and systemic inflammatory response [2]. In a model of *Pseudomonas aeruginosa* pneumonia in mechanically ventilated piglets, we observed a lower lung bacterial burden and less severe histological pneumonia in animals treated with corticosteroids plus ciprofloxacin than those who did not get corticosteroids [3].

Since 1955, multiple randomised clinical trials (RCTs) have assessed the efficacy of adjuvant treatment with systemic corticosteroids in patients with CAP. Among these, treatment with hydrocortisone resulted in a striking decrease in mortality in patients with severe CAP [4].

Until now, ten meta-analyses about corticosteroids in CAP have been published, with assessment of mortality [5–14]. Four focused on severe CAP [5, 7, 12, 13] and four others [6, 9–11] analysed severe CAP as a subgroup. In seven of these eight meta-analyses that evaluated patients with severe CAP, mortality was significantly decreased, with odds ratios ranging from 0.21 to 0.64. The length of stay was significantly reduced in the nine meta-analyses

that assessed this outcome (in one it was not searched for [6]). Time to clinical stability was reduced by about 1 day in four meta-analyses [8–10, 14], while this reduction was not estimated quantitatively in another one [11], and this information was not assessed in the remaining five studies. In five out of seven studies that reported the data, hyperglycemia was significantly more common in the treatment arm. Gastrointestinal bleeding was not increased by the use of corticosteroids in the six meta-analyses in which it was specifically investigated. Table 1 summarizes the characteristics and findings of these meta-analyses.

Despite the benefits observed in the meta-analyses and a reasonable frequency of side effects, the use of corticosteroids in CAP is still controversial and not routine. There are pitfalls in some of the RCTs included in these meta-analyses. First, the inclusion of patients with low severity may not be informative, since these patients have a low mortality and consequently it is very difficult to demonstrate differences in important outcomes such as treatment failure and mortality. Second, while critical illness-related corticosteroid insufficiency was not systematically assessed in these studies [15], few limited inclusion to those with high initial degrees of systemic inflammation. According to the rationale of using steroids in CAP, a high inflammatory response is necessary to benefit from this treatment. Patients with a high inflammatory response, i.e. high levels of C-reactive protein (CRP), have higher rates of treatment failure and mortality [16]. Third, the dosages, the type and the length of treatment are very different among RCTs, which makes it very difficult to establish comparisons among them. Fourth, the primary endpoints are different between studies and some of them, such as length of stay or even time to clinical stability, could be considered “soft”, since the former depends on other variables and the latter is driven by the persistence of fever, which in

*Correspondence: atorres@ub.edu

¹ Department of Pneumology, Respiratory Institute, Hospital Clinic of Barcelona, University of Barcelona (UB), Villarroel 170, 08036 Barcelona, Spain

Full author information is available at the end of the article

Table 1 Summary of the most relevant characteristics of meta-analyses published that have assessed the effects of corticosteroids in patients hospitalized with community-acquired pneumonia with the main outcomes reported

Author	Year	N RCTs	N patients	Mortality ^a	Mortality (SCAP) ^a	LOS ^b	TTCS ^b	Hyperglycaemia ^a	GI bleeding	Other side effects ^a
Siempos	2008	8	189 SCAP	–	0.21 (0.05–0.83)	Reduced (NE)	–	–	–	–
Nie	2012	9	1001	0.62 (0.32–1.04)	0.26 (0.11–0.64)	–	–	2.64 (1.68–4.15)	Similar	Similar superinfection
Cheng	2014	4	264 SCAP	–	0.39 (0.17–0.90)	Decreased in ICU	–	–	Similar	–
Siemieniuk	2015	12	1974	0.67 (0.45–1.01)	–	–1.00 (–1.79 to –0.21)	–1.22 (–2.08 to –0.35)	1.49 (1.01–2.19)	Similar	MV: 0.45 (0.26–0.49) ARDS: 0.24 (0.10–0.56)
Horita	2015	10	1780	0.80 (0.53–1.21)	0.41 (0.19–0.90)	–0.98 (–1.26 to –0.71)	–1.16 (–1.73 to –0.58)	Increased (NE)	–	–
Marti	2015	14	2077	0.84 (0.55–1.29)	0.47 (0.23–0.96)	0.89 ^c (0.70–0.89)	0.89 ^c (0.84–0.94)	1.59 (1.06–2.38)	Similar	MV: 0.41 (0.29–0.60)
Wan ^d	2016	9	1667	0.72 (0.43–1.21)	0.64 (0.32–1.29)	Reduced (NE)	Reduced (NE)	Similar	Similar	ARDS: 0.21 (0.08–0.59)
Bi	2016	8	538 SCAP	–	0.46 (0.28–0.77)	–4.76 (–8.13 to 1.40)	–	Similar	Similar	MV: 0.50 (0.27–0.91) ARDS: 0.23 (0.07–0.80)
Wu	2017	10	729 SCAP	–	0.49 (0.29–0.85)	–4.11 (–6.61 to 1.81)	–	–	–	–
Briel	2017	6	1506 (individual)	0.75 (0.46–1.21)	–	–1.15 (–1.75 to –0.55)	–1.63 (–1.62 to –0.43)	2.15 (1.60–2.90)	–	CAP-related re-admission 1.85 (1.03–3.32)

RCT randomised clinical trials, SCAP severe community-acquired pneumonia, NE not estimated, LOS length of stay in hospital, unless otherwise stated, TTCS time to clinical stability, GI gastrointestinal, ICU intensive care unit, MV mechanical ventilation, ARDS acute respiratory distress syndrome

^a Values correspond to odds ratios or risk ratios, with 95% confidence intervals in parentheses

^b Values correspond to weighted mean differences in days, with 95% confidence intervals in parentheses

^c Values correspond to geometrical mean ratios, with 95% confidence intervals in parentheses

^d This study also included six cohort studies with 4095 cases, but estimates are reported only from the nine RCTs

fact is downregulated by corticosteroids. Finally, it should be noted that the small sample size of some individual studies included in these meta-analyses is a limitation.

We performed a RCT [17] comparing methylprednisolone (0.5 mg/kg b.i.d. during 5 days) vs. placebo; but unlike other studies, we only included severe CAP patients and those with a high systemic inflammatory response, defined by CRP > 15 mg/dL. Also, rather than using mortality as a primary endpoint, we chose treatment failure, since it is associated with higher mortality of CAP. We also monitored the systemic inflammatory response, using different biomarkers, until day 7 after the inclusion of patients in the trial.

Our results showed a decrease in the treatment failure rate. The reduction in treatment failure was more evident in late treatment failure, defined as radiographic progression or persistence of respiratory failure, development of shock, need for invasive mechanical ventilation not present at baseline or death between 72 and 120 h after treatment initiation, and was especially clear for radiographic progression. Mortality did not differ significantly between groups, but the study was not powered for this endpoint. Despite the fact that we did not observe a rebound inflammation 2 days after corticosteroid discontinuation in CAP, we cannot exclude such a complication after longer periods of treatment without tapering.

After reviewing all the meta-analyses published and considering our own findings, we believe that it is now time to start introducing treatment with corticosteroids to selected ICU patients with severe CAP. We recommend selecting severe CAP patients with a high inflammatory response, as measured by CRP, at similar dosage and duration as we used in our RCT [17]. Methylprednisolone, prednisone or hydrocortisone at equivalent dosage appear similarly effective. Despite the fact that we excluded patients with influenza pneumonia in our trial because of data suggesting that corticosteroids are associated with increased mortality in influenza pneumonia, this association may not be present when adequate adjustments are made for time-dependent differences, and therefore these patients probably should not be excluded from corticosteroid treatment. In the future, we may find that other biomarkers are more specific in identifying patients who most benefit from this therapy, and we may have new therapies that are more selective regarding their anti-inflammatory impact on pneumonia.

We also need more data on the possible effect of corticosteroids on other viral pneumonias caused by adenovirus, rhinovirus, respiratory syncytial virus or others such as MERS. On the basis of the available data, we believe that corticosteroids are useful in specific patients with severe CAP and can help to decrease length of stay,

time to clinical stability, treatment failure and possibly mortality.

Author details

¹ Department of Pneumology, Respiratory Institute, Hospital Clinic of Barcelona, University of Barcelona (UB), Villarroel 170, 08036 Barcelona, Spain. ² Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain. ³ Centro de Investigación Biomedica En Red-Enfermedades Respiratorias (CibeRes, CB06/06/0028), Barcelona, Spain. ⁴ New York Presbyterian/Weill Cornell Medical Center, New York, NY, USA.

Acknowledgements

This study was supported by Grants SEPAR, FUCAP and SOCAP, 2009-SGR-661, IDIBAPS, ICREA academia 2013, Instituto de Salud Carlos III (PI15/00506), CibeRes (CB06/06/0028)-ISCIII.

Received: 6 September 2017 Accepted: 11 October 2017

Published online: 25 October 2017

References

- Mongardon N, Max A, Bougle A, Pene F, Lemiale V, Charpentier J, Cariou A, Chiche JD, Bedos JP, Mira JP (2012) Epidemiology and outcome of severe pneumococcal pneumonia admitted to intensive care unit: a multicenter study. *Crit Care* 16:R155
- Monton C, Ewig S, Torres A, El Ebiary M, Filella X, Rano A, Xaubet A (1999) Role of glucocorticoids on inflammatory response in nonimmunosuppressed patients with pneumonia: a pilot study. *Eur Respir J* 14:218–220
- Sibila O, Luna CM, Agusti C, Baquero S, Gando S, Patron JR, Morato JG, Absi R, Bassi N, Torres A (2008) Effects of glucocorticoids in ventilated piglets with severe pneumonia. *Eur Respir J* 32:1037–1046
- Confalonieri M, Urbino R, Potena A, Piattella M, Parigi P, Puccio G, Della PR, Giorgio C, Blasi F, Umberger R, Meduri GU (2005) Hydrocortisone infusion for severe community-acquired pneumonia: a preliminary randomized study. *Am J Respir Crit Care Med* 171:242–248
- Siempos II, Vardakas KZ, Kopterides P, Falagas ME (2008) Adjunctive therapies for community-acquired pneumonia: a systematic review. *J Antimicrob Chemother* 62:661–668
- Nie W, Zhang Y, Cheng J, Xiu Q (2012) Corticosteroids in the treatment of community-acquired pneumonia in adults: a meta-analysis. *PLoS One* 7:e47926
- Cheng M, Pan ZY, Yang J, Gao YD (2014) Corticosteroid therapy for severe community-acquired pneumonia: a meta-analysis. *Respir Care* 59:557–563
- Siemieniuk RA, Meade MO, Alonso-Coello P, Briel M, Evaniew N, Prasad M, Alexander PE, Fei Y, Vandvik PO, Loeb M, Guyatt GH (2015) Corticosteroid therapy for patients hospitalized with community-acquired pneumonia: a systematic review and meta-analysis. *Ann Intern Med* 163(7):519–528
- Horita N, Otsuka T, Haranaga S, Namkoong H, Miki M, Miyashita N, Higa F, Takahashi H, Yoshida M, Kohno S, Kaneko T (2015) Adjunctive systemic corticosteroids for hospitalized community-acquired pneumonia: systematic review and meta-analysis 2015 update. *Sci Rep* 5:14061
- Marti C, Groscurin O, Harbarth S, Combescure C, Abbas M, Rutschmann O, Perrier A, Garin N (2015) Adjunctive corticotherapy for community acquired pneumonia: a systematic review and meta-analysis. *PLoS One* 10:e0144032
- Wan YD, Sun TW, Liu ZQ, Zhang SG, Wang LX, Kan QC (2016) Efficacy and safety of corticosteroids for community-acquired pneumonia: a systematic review and meta-analysis. *Chest* 149:209–219
- Bi J, Yang J, Wang Y, Yao C, Mei J, Liu Y, Cao J, Lu Y (2016) Efficacy and safety of adjunctive corticosteroids therapy for severe community-acquired pneumonia in adults: an updated systematic review and meta-analysis. *PLoS One* 11:e0165942
- Wu WF, Fang Q, He GJ (2017) Efficacy of corticosteroid treatment for severe community-acquired pneumonia: a meta-analysis. *Am J Emerg Med*. doi:10.1016/j.ajem.2017.07.050

-
14. Briel M, Spoorenberg SMC, Snijders D, Torres A, Fernandez-Serrano S, Meduri GU, Gabarrús A, Blum CA, Confalonieri M, Kasenda B, Siemieniuk RAC, Boersma W, Bos WJW, Christ-Crain M, Ovidius study group, Capisce study group, STEP study group (2017) Corticosteroids in patients hospitalized with community-acquired pneumonia: systematic review and individual patient data meta-analysis. *Clin Infect Dis*. doi:[10.1093/cid/cix801](https://doi.org/10.1093/cid/cix801)
 15. Annane D, Pastores SM, Rochweg B, Arlt W, Balk RA, Beishuizen A, Briegel J, Carcillo J, Christ-Crain M, Cooper MS, Marik PE, Umberto MG, Olsen KM, Rodgers S, Russell JA, van den BG (2017) Guidelines for the diagnosis and management of critical illness-related corticosteroid insufficiency (CIRCI) in critically ill patients (Part I): Society of Critical Care Medicine (SCCM) and European Society of Intensive Care Medicine (ESICM) 2017. *Intensive Care Med*. doi:[10.1097/CCM.0000000000002737](https://doi.org/10.1097/CCM.0000000000002737)
 16. Menendez R, Martinez R, Reyes S, Mensa J, Filella X, Marcos MA, Martinez A, Esquinas C, Ramirez P, Torres A (2009) Biomarkers improve mortality prediction by prognostic scales in community-acquired pneumonia. *Thorax* 64:587–591
 17. Torres A, Sibila O, Ferrer M, Polverino E, Menendez R, Mensa J, Gabarrús A, Sellares J, Restrepo MI, Anzueto A, Niederman MS, Agusti C (2015) Effect of corticosteroids on treatment failure among hospitalized patients with severe community-acquired pneumonia and high inflammatory response: a randomized clinical trial. *JAMA* 313:677–686