




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Original research

Potentially preventable urinary tract infection in patients with type 2 diabetes – A hospital-based study

Maria Cristina Carrondo^{a,*}, Joaquim Jorge Moita^b^a Department of Clinical Physiology, Polytechnic Institute of Castelo Branco, High School of Health Dr. Lopes Dias, Castelo Branco, Portugal^b Hospital Center and University of Coimbra, Coimbra, Portugal

ARTICLE INFO

Keywords:

Type 2 diabetes mellitus
 Urinary tract infections
 In-hospital
 Prediction model
 Cross-sectional study

ABSTRACT

Aim: To investigate the prevalence of urinary tract infections in hospitalized patients with type 2 diabetes mellitus and identify corresponding risk factors.

Methods: We conducted a cross-sectional study on 7.347 patients with type 2 diabetes mellitus as the principal diagnosis, using hospitalization discharge summary data from January 1 to December 31, 2015. Disease stages were classified as stages 1, 2, and 3.

Results: Of 7.347 patients, 16.2% had urinary tract infections. The urinary tract infection prevalence was 24.4% in 428 patients in stage 1 and 4.8% in 2.840 patients in stage 2; it was higher among patients who underwent medical procedures than among those who underwent surgery (24.4% vs 4.8%). In multivariate regression analysis, age (OR = 1.031; 95% CI = 1.02–1.04), length of hospitalization (OR = 1.018; 95% CI = 1.013–1.024), sex (woman) (OR = 2.248; 95% CI = 1.778–2.842), comorbidity of stage 3 cerebrovascular disease (OR = 1.737; 95% CI = 1.111–2.714), and comorbidity of stage 1 colorectal cancer (OR = 2.417; 95% CI = 1.152–5.074) were found to be the risk factors of urinary tract infection in the ten hospitals considered. **Conclusions:** Our findings suggest that urinary tract infection prevalence was higher in women without evidence of organ injury and those receiving medical treatment. Comorbidities (cerebrovascular disease and colorectal cancer) were identified as risk factors.

1. Introduction

The prevalence of type 2 diabetes mellitus (T2DM) has increased in recent years (Zheng et al., 2018; *isk Factor Collabora*, 2016). In 2015, the International Diabetes Federation reported that approximately 415 million adults (aged 20–79 years) had T2DM; this number is expected to increase to 642 million by 2040 (Zheng et al., 2018). Globally, the prevalence of age-standardized adult diabetes increased from 4.3% in 1980 to 9.0% in 2014 in men and from 5.0% to 7.9% in women in the same time frame (*isk Factor Collabora*, 2016). By 2015, T2DM had occurred in 12.1% of men and 7.7% of women in Portugal (*Sociedade Portuguesa de Diabetologia*, 2015); only 7.3% of these patients received primary care (*Sociedade Portuguesa de Diabetologia*, 2015).

T2DM and its complications have contributed to an increase in the rate of admissions (Zheng et al., 2018). The admission rates for patient with T2DM are two to six times higher than those for patients without diabetes (Lara-Rojas et al., 2019). In Portugal, 88–870 adults had T2DM

in 2006, and hospital admission rates increased by 82.7% between 2006 and 2015, with acute myocardial infarction being the primary cause of hospitalization (*Sociedade Portuguesa de Diabetologia*, 2015). Microangiopathic disease and stroke were identified as predictors of hospitalization within Portugal's National Health System (*Sociedade Portuguesa de Diabetologia*, 2015). In contrast, cardiovascular disease is the primary cause of hospitalization in Spain (Lara-Rojas et al., 2019).

In Portugal, urinary tract infections (UTIs) are frequent adverse events during hospitalization (<https://apurologia.pt/>, A, 2008), for which T2DM is a major risk factor (Wilke et al., 2015; Hirji et al., 2012). However, the cause of UTI is considered to be multifactorial (<https://apurologia.pt/>, A, 2008). Studies (Wilke et al., 2015; Hirji et al., 2012) have reported that UTI is prevalent in those with diabetes, particularly in older women. Wilke et al. (2015) showed that 3.86% of UTI events were associated with inpatient hospitalization and 35.6% of UTI events were caused by more than one reported bacterium. It is difficult to

Abbreviations: T2DM, type 2 diabetes mellitus; UTI, urinary tract infection; COC, complications-of-care

* Corresponding author. Polytechnic Institute of Castelo Branco, High School of Health Dr. Lopes Dias, Department of Clinical Physiology, Avenida do Empresário, 6000-767, Castelo Branco, Portugal.

E-mail address: cristina.carrondo@ipcb.pt (M.C. Carrondo).

<https://doi.org/10.1016/j.obmed.2020.100190>

Received 14 January 2020; Accepted 25 January 2020

2451-8476/ © 2020 Elsevier Ltd. All rights reserved.

determine the actual incidence of UTIs in Portugal, but approximately 15% of women aged 70 years have asymptomatic bacteriuria; this rate increases to 30–40% in elderly women admitted to geriatric institutions and practically to 100% in patients with permanent urinary tubes (<https://apurologia.pt/>, A, 2008).

Mazzi et al. (2017) reported that other risk factors, such as severe hypoglycemic events, advanced age, and the presence of comorbidities, may increase the occurrence of infection in diabetes patients. Another study (de Decker et al., 2017) found that a high burden of comorbidities (stroke, heart failure, myocardial infarction, and functional cognitive impairment) increased the risk of hypoglycemic events in elderly patients (aged over 80 years) with diabetes. Mendez-Bailon et al. (2017) studied 715 patients with diabetes admitted for cardiovascular diseases who underwent invasive cardiac procedures, and found that obesity (18%), arterial hypertension (67.3%), lipid abnormalities (45.8%), renal failure (14.07%), and intermittent claudication (1.65%) were risk factors for adverse events during hospitalization.

Recently, a meta-analysis (Rodríguez-Acelas et al., 2017) revealed that the duration of surgery, duration of central venous catheterization, number of re-operations, type of hospitalization, admission to the intensive care unit, use of immunosuppressants, and use of cephalosporins were risk factors for adverse events during hospitalization in patients with diabetes. Akirov et al. (2017) reported that variability in glycemic levels (glucose \geq 180 mg/dl) during hospitalization increases the length of stay, rate of infections and complications, and mortality risk.

This study aimed to investigate the prevalence of UTI events during hospitalization and corresponding risk factors in those with diabetes that were hospitalized.

1.1. Subjects

The study was performed in 32 public hospitals in Portugal. The medical data of 7,347 patients with T2DM between January 1 and December 31, 2015 were reviewed.

Inclusion criteria were hospitalized patients with T2DM as the principal diagnosis, older than 18 years. Exclusion criteria included patients with type 1 of diabetes mellitus and patients with T2DM admitted to hospitals providing oncology services and mental health.

2. Materials and methods

This retrospective study used medical data from the National Hospital Morbidity Database provided by the Central Administration of the Health System. We used (1) the Diagnosis Related Group (DRG) to identify episodes of T2DM (codes DX 250.00) as principal diagnoses during hospitalization and (2) *Staging* [13] version 5.26 to identify the severity of T2DM, its comorbidities and stages, and UTI events.

Staging uses a specific methodology – *MEDSTAT-COC* [13] – that identifies potential UTI events (designated as complications-of-care [COC] Urinary) through principal and secondary diagnostic codes, *ICD-9-CM*, as well as *ICD-9-CM* procedure codes, age, sex, and the number of days of hospitalization. The parameters for COC Urinary were as follows: code 59010 – acute pyelonephritis not specified; code 5909 – infection of the kidney not specified; code 5950 – acute cystitis; code 5953 – trigonitis; code 5959 – cystitis not specified; and code 5990 – urinary tract infection not specified. UTI events that started from day 1 of admission until discharge from the hospital were investigated in this study.

2.1. Statistical analysis

Bivariate analysis was used to determine the prevalence of UTIs. The distribution and the rate of COC Urinary were calculated using the following formula14:

- COC Urinary infection rate = total number of patients with infection/sum of patients at risk without infection and at risk with infection * 100.

Through the *Staging* software (Thomson Reuters, 2009), COC risk was explicitly assessed according to the relationship between the principal and secondary diagnosis codes, *ICD-9-CM*, and defined as (i) a patient at risk for a particular infection and (ii) a patient with evidence of potential infection of care during hospitalization.

To determine the risk factors of UTI, we used multivariate logistic regression analysis. The risk adjustment model was used to estimate the correlation between the occurrence of COC Urinary (dependent variable) and the various risk factors (independent variables). The forward conditional method was used for stepwise selection of the variables, whereas the Hosmer and Lemeshow (H-L) test was used to adjust the model with the independent variables. For the validation of the model, its discriminatory capability, sensitivity, and specificity were analyzed using the area under the curve of the receiver operating characteristic curve (AUROC) (c). COC Urinary was set as the adverse outcome and dependent variable that assumed the values: 0 (without infection) and 1 (with infection). Independent variables were selected from variables previously reported to be associated with less favorable health outcomes (Iezzoni and Iezzoni, 2003) as follows:

- (1) Length of hospitalization: the length of stay causes adverse outcomes because patients are more exposed to hospital infections. The variable was constructed based on the total number of days in-hospital.
- (2) Severity of primary disease: the severity of disease increases the risk of death or organ failure and weakens the immune system. We considered 3 stages (Thomson Reuters, 2009): stage 1, initial phase of the disease without evidence of organ injury; and stages 2 and 3, more advanced severity levels of the disease with evidence of organ injury and micro- and macrovascular complications, respectively.
- (3) The comorbidities and stage: comorbidities increase the disease burden. We focused on the most relevant comorbidities (defined according to the number of in- and out-of-hospital deaths previously reported to represent a significant burden in those with diabetes. The following comorbidities were included: lipid abnormalities (NUT82); essential hypertension (CVS13); obesity (NUT02); coronary artery disease without prior coronary revascularization (CVS11); cerebrovascular disease (NEU04); congestive heart failure (CVS09); pneumonia: bacterial (RES15); renal failure (GUS08); other disorders of the respiratory system (RES83); rhino, adeno, and corona virus infections (RES24); neoplasm, malignant: colon and rectum (GIS27); neoplasm, malignant: lungs, bronchi, or mediastinum (RES13); and neoplasm, malignant: stomach (GIS30). For all comorbidities, we considered three stages (Thomson Reuters, 2009): stage 1, known diagnosis, without local or systemic complications; stage 2, disease limited to an organ or system, increased risk of adverse outcomes; and stage 3, generalized involvement of the system, with systemic complications and with even higher risk of adverse outcomes.
- (4) Treatment type: the treatment type may increase the risk of bacterial infections, particularly if preventive procedures are not taken. This was categorized as surgical (surgical treatments) and medical (other invasive diagnostic and therapeutic procedures) and was constructed based on the admission episode and was classified according to the DRG criteria.
- (5) Hospitals: some hospitals do not apply infection control programs. This variable was constructed based on the volume of hospitalization (episodes/population) in descending order: hospital (1) represents the highest-volume hospital, and hospital (32) represents the lowest-volume hospital. Age and sex.

OR values were presented with their respective 95% confidence intervals (CI) or p-values. The p-value set to indicate statistical significance was $p < 0.005$. All statistical analyses were performed using the IBM SPSS Statistics for Windows, Version 22.0. (Armonk, NY: IBM Corp.).

2.2. Ethical considerations

This study was approved by the Central Administration of the Health System. The need for written informed consent from all participants was waived because this was a retrospective study with anonymized patient data.

3. Results

3.1. Study population

Of the 7,347 patients with T2DM who were admitted, 3,757 (51.1%) patients were men, and 3,590 (48.9%) were women. The majority of patients were elderly, with 71.1% above 65 years. The median duration of hospitalization was 8 days (average 12 ± 18 days).

The total UTI rate was 16.2%. Patient characteristics are shown in Table 1.

Fig. 1 shows the proportion of T2DM patients by hospital. In hospital (1), 631 patients were admitted, corresponding to 7.9% of the population; meanwhile, in hospital (32), 101 patients were admitted, corresponding to 1.3% of the hospital production.

3.2. Urinary tract infection rate

The UTI infection rate was significantly higher in women than in men (23.6% vs 10.5%; OR = 2.65; $p < 0.0001$), in those aged above 85 years than in those aged 18–64 years (27.5% vs 9%; OR = 3.84; $p < 0.0001$), and in those with stage 1 disease than in those with stage 2 (24.4% vs 4.8%; OR = 0.16; $p < 0.0001$). It was also significantly higher in those who received medical treatment than those who received surgical treatment (24.6% vs 4.8%; OR = 6.52; $p < 0.0001$). Table 2 shows the values of UTI rate by sex, age (stratified), severity of T2DM, comorbidities, and treatment type.

3.3. Risk factors

Multivariate regression analysis with COC Urinary as the dependent variable showed that the following variables were significant predictors of UTIs: age of at least 18 years (OR = 1.031; 95% CI = 1.02–1.041); length of hospitalization (OR = 1.018; 95% IC = 1.013–1.024); sex (woman) (OR = 2.248; 95% IC = 1.778; 2.842); comorbidity of stage 3 cerebrovascular disease (OR = 1.737; 95% IC = 1.111–2.714); and comorbidity of stage 1 colorectal cancer (OR = 2.417; 95% IC = 1.152–5.074). Table 3 shows the result of the logistic regression analysis with COC Urinary as the dependent variable.

Table 1
Demographic characteristics.

Population		Total (%)	Age (years)				Mean \pm SD
			18–64	65–74	75–84	> 85	
Sex	Men	3757 (51.1)	1284 (60.7)	1053 (57.5)	1066 (45.8)	354 (33.0)	69 \pm 13
	Women	3590 (48.9)	832 (39.3)	779 (42.5)	1259 (54.2)	720 (67.0)	73 \pm 13
Severity of T2DM	Stage 1	428 (5.8)					
	Stage 2	2840 (38.7)					
	Stage 3	4079 (55.5)					
Treatment type	Surgical	1813 (24.7)					
	Medical	5534 (75.3)					

Abbreviations: T2DM, type 2 of diabetes mellitus; SD, standard deviation.




Fig. 1. Frequency of admitted T2DM patients by hospital
Abbreviations: T2DM, type 2 of diabetes mellitus.

Among hospitals, hospitals 19 and 32 showed the lowest and higher OR value of 2.167 and 4.741, respectively. Meanwhile, among the three stages of disease severity, stage 2 showed the highest OR of 0.604 (0.304,1.199).

The p-value found for the model was $p < 0.0001$ in Table 4. The H-L test revealed a p-value of 0.848 in Table 5, and the area under curve was 0.862, indicating that the model had a good sensitivity and specificity in Fig. 2.

Collectively, the results of this model show that (1) the risk of UTIs was higher in hospitals 5, 8, 11, 12, 13, 19, 20, 27, 28, and 32; (2) the comorbidity of stage 3 cerebrovascular disease and comorbidity of stage 1 colorectal cancer are risk factors for UTIs; and (3) the severity of T2DM is not an explanatory factor of UTIs. Our study showed that UTI rate is higher in women without evidence of organ injury and in those receiving medical treatment.

4. Discussion

A recent epidemiological study (Tandogdu and Wagenlehner, 2016) showed significant geographic variations in the frequency of UTIs in those with diabetes, with the rate being the highest in developing countries (24%) compared with that in the US (12.9%) and Europe (19.6%) (Tandogdu and Wagenlehner, 2016). In our study, the rate of UTIs was 16.2%, indicating differences in healthcare delivery and systems between developing and developed countries.

We found a higher rate of UTIs in women than in men (UTI rate, 23.6% vs 10.5%). Further, it was also higher in those aged at least 85 years (27.5%). Popejoy et al. (2017) found that 48.5% of women with diabetes and aged above 65 years had UTIs. The discrepancy in the UTI rates may imply better healthcare in Portugal.

UTIs were also more common after medical treatment than after surgical treatment (24.6% vs 4.8%), consistent with the findings reported in a previous study (Shah et al., 2018). This was attributed to (i) the excessive use of certain medical devices involving urologic

Table 2
COC Urinary rate by sex, age, severity of T2DM, comorbidities, and treatment type.

COC Urinary					
Variables		Rate (%)	OR	P-value	
Sex	Men (ref.)	10.5	2.65	< 0.0001	
Age (years)	18–64 (ref.)	9.0			
	65–74	12.6	1.47	0.011	
	75–84	22.2	2.9	< 0.0001	
Severity of T2DM	Above 85	27.5	3.84	< 0.0001	
	Stage 1 (ref.)	24.4			
	Stage 2	4.8	0.16	< 0.0001	
Comorbidities	Stage 3	24.7	1.02	0.9445	
	NUT82 (ref.)	14.6			
	CVS13	21.5	1.6	0.0005	
	NUT02	16.8	1.18	0.2614	
	CVS11	14.8	1.02	0.9266	
	NEU04	22.7	1.72	0.0008	
	CVS09	20.7	1.52	0.0423	
	RES15	24.6	1.91	0.0006	
	GUS08	23.1	1.76	0.0034	
	RES83	25.5	2.01	0.0003	
	RES24	18.5	1.32	0.4	
	GIS27	28.2	2.29	0.0031	
	RES13	8.3	0.53	0.5468	
	GIS30	12.5	0.84	0.8133	
	Treatment type	Surgical (ref.)	4.8	6.52	< 0.0001
		Medical	24.6		

COC Urinary, urinary tract infection after procedure; NUT82, lipid abnormalities; CVS13, essential hypertension; NUT02, obesity; CVS11, coronary artery disease without prior coronary revascularization; NEU04, cerebrovascular disease; CVS09, congestive heart failure; RES15, pneumonia: bacterial; GUS08, renal failure; RES83, other disorders of the respiratory system; RES24, rhino, adeno, and corona virus infections; GIS27, neoplasm, malignant: colon and rectum; RES13, neoplasm, malignant: lungs, bronchi, or mediastinum; GIS30, neoplasm, malignant: stomach. Rate (%) = Total number of patients with infection/sum of patient at risk without infection and at risk with infection *100, Abbreviation: T2DM, type 2 of diabetes mellitus; OR, odds ratio; ref., reference group.

manipulation (e.g. catheters); (ii) failure to follow aseptic techniques; and (iii) procedures in patients with diabetes with poor glycemic control and/or with asymptomatic bacteriuria. The difference in the frequency of UTIs between the type of treatment may be due to the different therapeutic strategies employed; for example, antibiotic prophylaxis was administered in patients undergoing surgery.

With respect to the severity of diabetes, we found a higher UTI rate in patients with stage 2 diabetes than that of those with stage 1 diabetes (4.8% vs 24.4% respectively; OR = 0.16), contradicting previous studies (isk Factor Collabora, 2016; Ljungqvist et al., 2005). Increased hyperglycemia may occur after surgical stress (Ljungqvist et al., 2005), and, thus, our results may indicate a non-effective glycemic control in those with stage 1 severity of diabetes. Typically, patients with diabetes are only diagnosed with UTI during their first hospital visit in the health services of the Portuguese National Health System because of the gap in follow-up of these patients in primary care (Sociedade Portuguesa de Diabetologia, 2015). This may have caused a higher risk of UTIs in those with stage 1 severity T2DM than those with stage 2, because patients with stage 2 T2DM have previous glycemic control.

Logistic regression analysis showed the influence of hospitalization on the risk of UTI. This is consistent with the findings of some studies (https://apurologia.pt/. A, 2008; Wilke et al., 2015; Hirji et al., 2012; Mazzi et al., 2017; de Decker et al., 2017; Mendez-Bailon et al., 2017; Rodríguez-Acelas et al., 2017; Shwartz et al., 1994) that have reported a high number of unnecessary hospitalizations. One study (Rosenthal et al., 2014) showed that marked differences exist in the use of medical resources, hospital beds, diagnostic exams, and medical treatments putting the patient at risk of infection. The findings in the current study

Table 3
Logistic regression analysis for COC Urinary.

COC Urinary		
Independent variables	OR	(95% CI)
Age	1.031	(1.02–1.041)
Duration of hospitalization (days)	1.018	(1.013–1.024)
Sex		
Men (Ref.)	1	
Women	2.248	(1.778–2.842)
Stages NEU04		
Without (Ref.)	1	
Stage 1	1.12	(0.68–1.843)
Stage 3	1.737	(1.111–2.714)
Stages GIS27		
Without (Ref.)	1	
Stage 1	2.417	(1.152–5.074)
Stage 2	2.936	(0.782–11.024)
Stage 3	2.024	(0.171–24.013)
Hospital		
1 (Ref.)	1	
5	4.024	(2.319–6.983)
8	3.149	(1.643–6.035)
11	2.995	(1.337–6.712)
12	3.339	(1.589–7.015)
13	2.176	(1.039–4.558)
19	2.167	(1.088–4.315)
20	3.06	(1.23–7.617)
27	2.828	(1.05–7.618)
28	4.528	(1.949–10.518)
32	4.741	(1.988–11.309)
Severity of T2DM		
Stage 1 (Ref.)	1	
Stage 2	0.604	(0.304–1.199)
Constant	< 0.001	

NEU04, cerebrovascular disease; GIS27, neoplasm, malignant: colon and rectum.

Abbreviations: OR, odds ratio; CI, confidence interval; Ref., reference group; T2DM, type 2 of diabetes mellitus.

Table 4
Omnibus tests of model coefficients.

		Chi-square	Df	Sig.
Step 1	Step	540.183	50	.000
	Block	540.183	50	.000
	Model	540.183	50	.000
Step 2	Step	45.478	1	.000
	Block	585.662	51	.000
	Model	585.662	51	.000
Step 3	Step	54.315	1	.000
	Block	639.976	52	.000
	Model	639.976	52	.000
Step 4	Step	42.777	1	.000
	Block	682.753	53	.000
	Model	682.753	53	.000
Step 5	Step	69.087	30	.000
	Block	751.840	83	.000
	Model	751.840	83	.000
Step 6	Step	15.208	2	.000
	Block	767.048	85	.000
	Model	767.048	85	.000
Step 7	Step	9.890	1	.002
	Block	776.937	86	.000
	Model	776.937	86	.000
Step 8	Step	6.763	1	.009
	Block	783.700	87	.000
	Model	783.700	87	.000
Step 9	Step	11.956	3	.008
	Block	795.656	90	.000
	Model	795.656	90	.000
Step 10	Step	5.263	1	.022
	Block	800.919	91	.000
	Model	800.919	91	.000

Abbreviations: df, degrees of freedom; Sig., significance.

Table 5
Hosmer and Lemeshow test.

Step	Chi-square	Df	Sig.
1	.000	7	1.000
2	12.967	8	.113
3	7.956	8	.438
4	3.705	8	.883
5	5.943	8	.654
6	7.156	8	.520
7	8.076	8	.426
8	4.112	8	.847
9	5.147	8	.742
10	4.095	8	.848

Abbreviations: df, degrees of freedom; Sig., significance.




Fig. 2. Receiver operating characteristics curve.

may be due to (1) discrepancies in the use of hospital resources; (2) concentration of specific procedures to certain patient groups; and (3) differences in the clinical management of patients. Further, some studies (Galiczewski and Shurpin, 2017; Kummerow Broman et al., 2018; Fairfield et al., 2018) have reported unsafe treatments, inadequate human and technical resources, and, consequently, poor healthcare practices, leading to variations in the frequency of UTIs among geographic areas. Some hospitals do not observe safety protocols in the insertion, maintenance, and aseptic removal of devices and do not comply with guidelines for the prevention of UTIs, such as the assessment of the need for catheterization (based on individual risk); do not select the type of catheter according to the predicted duration of catheterization; do not implement prevention and control policies for asymptomatic bacteriuria; and do not follow clinical practice guidelines for managing hospitalized patients with diabetes who have UTIs.

With respect to sex, women were at a higher risk of UTIs than men (OR = 2.248), consistent with other studies (Popejoy et al., 2017; Costantini et al., 2017). This may be due to (i) hypotonic neurogenic bladder and (2) physiological changes in bladder associated with aging. This raises a question about the efficacy of treatment of urinary tract alterations.

With respect to diabetes severity, no difference was observed in the risk of UTIs between those with stage 2 and stage 1 T2DM, in contrast to other studies that reported that the severity of diabetes is a predictor of COC Urinary (isk Factor Collabora, 2016; Popejoy et al., 2017). These findings may be due to the differences in clinical coding between health

institutions and the definition of the severity of diabetes. This result raises a question about the use of the guidelines for T2DM diagnosis.

With respect to the comorbidities, we found that cerebrovascular disease and malignant neoplasm of the colon and rectum are predictors of UTIs, consistent with previous studies (Kwaan et al., 2017; Mi et al., 2018; Vermeij et al., 2018). Particularly, the association between stage 3 cerebrovascular disease and UTI may be due to the use of ineffective prophylactic procedures and differences in clinical coding. These findings indicate the need for more effective clinical safety protocols for nursing staff. Meanwhile, the comorbidity of malignant neoplasm of the colon and rectum may be due to (1) ineffective prophylactic measures, (2) differences in clinical practice, (3) differences in clinical coding, and (4) differences in the use of hospital resources, as previously reported (Kwaan et al., 2017; D'Hondt et al., 2017; Eveno et al., 2010). Some hospitals do not offer specialized care, and thus, we could assess what treatments are available in these hospitals.

Our study has several limitations. First, biochemical data, particularly, the glycemic values in pre- and post-treatment that may result in less favorable outcomes, were lacking. Second, inconsistencies may have been present in the clinical coding of medical data (codes ICD-9-CM) among hospitals. Therefore, random audits of medical processes have caused better handling of medical data, in Portugal. Third, we do not know if the cause of UTIs was preventable or not. Lastly, this study has all the limitations and risks of bias inherent to cross-sectional study design, thus limiting the generalizability of the conclusions to other populations. Given the increasing incidence of T2DM worldwide, particularly in Portugal, diabetes-specific healthcare guidelines during hospitalization are needed, and the number of physicians and nurses in primary care should be increased to meet the increased need for proper management of comorbidities, in order to avoid more hospitalizations.

In this study, we identified individuals with diabetes, who were at a higher risk of UTI, which is the most common adverse event during hospitalization in patients with diabetes. Our findings suggest that women in stage 1 of T2DM have the highest prevalence of UTIs and those who have received medical treatment with fewer comorbidities. More effective health policies should be implemented to manage T2DM, particularly in stage 1 disease. Home visits by primary care physicians and nurses are crucial for those with limited healthcare access for better diabetes management and for reducing admission rates. In hospitals, health professionals should control hyperglycemia before and after medical treatments and should adopt safe protocols to reduce UTIs. Further, health education on diet and nutrition should be provided to reduce comorbidities due to increased risk for UTIs. This study can be beneficial for risk stratification of these patients and for the early identification of those at higher risk of infection, and the findings can be used to implement optimal healthcare measures to prevent UTIs.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declarations of interest

None of the authors has any declarations of interest.

Acknowledgments

None.

References

- Akirov, A., Grossman, A., Shochat, T., et al., 2017. Hyperglycemia on admission and hospitalization outcomes in patients with atrial fibrillation. *Clin. Cardiol.* 40, 1123–1128.
- Costantini, E., Illiano, E., Giannitsas, K., et al., 2017. Urological dysfunction in young

- women: an inheritance of childhood? *BJU Int.* 121, 453–457.
- D Hondt, M., Yoshihara, E., Dedrye, L., et al., 2017. Transanal endoscopic operation for benign rectal lesions and T1 carcinoma. *J. Soc. Laparoendosc. Surg.* 21 pii: e2016.00093.
- de Decker, L., Hanon, O., Boureau, A.S., et al., 2017. Association between hypoglycemia and the burden of comorbidities in hospitalized vulnerable older diabetic patients: a cross-sectional, population-based study. *Diabetes Ther.* 8, 1405–1413.
- Eveno, C., Lamblin, A., Mariette, C., et al., 2010. Sexual and urinary dysfunction after proctectomy for rectal cancer. *J. Vis. Surg.* 147, e21–e30.
- Fairfield, K.M., Black, A.W., Lucas, F.L., et al., 2018. Behavioral risk factors and regional variation in cardiovascular health care and death. *Am. J. Prev. Med.* 54, 376–384.
- Galiczewski, J.M., Shurpin, K.M., 2017. An intervention to improve the catheter associated urinary tract infection rate in a medical intensive care unit: direct observation of catheter insertion procedure. *Intensive Crit. Care Nurs.* 40, 26–34.
- Hirji, I., Guo, Z., Andersson, S.W., et al., 2012. Incidence of urinary tract infection among patients with type 2 diabetes in the UK General Practice Research Database (GPRD). *J. Diabet. Complicat.* 26, 513–516.
- Associação Portuguesa de Urologia, 2008. *Guia de prática clínica: cistite não complicada na mulher.* [Cited 2019 August 6] Available from: <https://apurologia.pt/wp-content/uploads/2018/10/Guia-cistite.pdf>.
- Iezzoni, L.I., 2003. Reasons for risk adjustment. In: Iezzoni, L.I. (Ed.), *Lit. – Risk Adjustment for Measuring Health Care Outcomes*, third ed. Health Administration Press. American College of Healthcare Executives, Chicago, pp. 1–17.
- NCD Risk Factor Collaboration (NCD-RisC), 2016. Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* 387, 1513–1530s.
- Kummerow Broman, K., Ward, M.J., Poulouse, B.K., et al., 2018. Surgical transfer decision making: how regional resources are allocated in a regional transfer network. *Joint Comm. J. Qual. Patient Saf.* 44, 33–42.
- Kwaan, M.R., Fan, Y., Jarosek, S., et al., 2017. Long-term risk of urinary adverse events in curatively treated patients with rectal cancer: a population-based analysis. *Dis. Colon Rectum* 60, 682–690.
- Lara-Rojas, C.M., Pérez-Belmonte, L.M., López-Carmona, M.D., et al., 2019. National trends in diabetes mellitus hospitalization in Spain 1997–2010: analysis of over 5.4 millions of admissions. *Eur. J. Intern. Med.* 60, 83–89.
- Ljungqvist, O., Nygren, J., Soop, M., et al., 2005. Metabolic perioperative management: novel concepts. *Curr. Opin. Crit. Care* 11, 295–299.
- Mazzi, S., Ravasio, R., Forlani, G., et al., 2017. Estimating the risk of severe hypoglycemic event related to glucose-lowering treatment among Italian patients with diabetes: the HYPOTHESIS database. *Clinicoecon. Outcomes Res.* 9, 711–720.
- Mendez-Bailon, M., Lorenzo-Villalba, N., Muñoz-Rivas, N., et al., 2017. Transcatheter aortic valve implantation and surgical aortic valve replacement among hospitalized patients with and without type 2 diabetes mellitus in Spain (2014–2015). *Cardiovasc. Diabetol.* 16, 144.
- Mi, H., Li, S., Li, H., et al., 2018. The effects of infection on severe stroke patients in the neurological intensive care unit in China. *Int. J. Neurosci.* 128, 715–720.
- Popejoy, M.W., Long, J., Huntington, J.A., 2017. Analysis of patients with diabetes and complicated intra-abdominal infection or complicated urinary tract infection in phase 3 trials of ceftolozane/tazobactam. *BMC Infect. Dis.* 17, 316.
- Rodríguez-Acelas, A.L., de Abreu Almeida, M., Engelman, B., et al., 2017. Risk factors for health care-associated infection in hospitalized adults: systematic review and meta-analysis. *Am. J. Infect. Contr.* 45, e149–e156.
- Rosenthal, V.D., Maki, D.G., Mehta, Y., et al., 2014. International Nosocomial Infection Control Consortium (INICC) report, data summary of 43 countries for 2007–2012. Device-associated module. *Am. J. Infect. Contr.* 42, 942–956.
- Shah, K.J., Cherabuddi, K., Shultz, J., et al., 2018. Ampicillin for the treatment of complicated urinary tract infections caused by vancomycin resistant enterococcus spp (VRE): a single-center university hospital experience. *Int. J. Antimicrob. Agents* 51, 57–61.
- Shwartz, M., Ash, A.S., Anderson, J., et al., 1994. Small area variations in hospitalization rates: how much you see depends on how you look. *Med. Care* 32, 189–201.
- Sociedade Portuguesa de Diabetologia Relatório Anual do Observatório da Diabetes: “Diabetes Factos e Números – o ano 2015.” [Cited 2019 August 3] Available from: <http://spd.pt/images/OND/DFN2015.pdf>.
- Tandogdu, Z., Wagenlehner, F.M., 2016. Global epidemiology of urinary tract infections. *Curr. Opin. Infect. Dis.* 29, 73–79.
- Thomson Reuters, 2009. *Disease Staging: Clinical and Coded Criteria.* The MEDSTAT Group, Inc., Ann Arbor, Michigan [Cited 2019 May 6] Available from: www.hcup-us.ahrq.gov/db/nation/nis/DiseaseStagingV5.27ClinicalandCodedCriteria.pdf, Version 5.27.
- Vermeij, J.D., Westendorp, W.F., Dippel, D.W., et al., 2018. Antibiotic therapy for preventing infections in people with acute stroke. *Cochrane Database Syst. Rev.* 1, CD008530. <https://doi.org/10.1002/14651858.CD008530>.
- Wilke, T., Boettger, B., Berg, B., et al., 2015. Epidemiology of urinary tract infections in type 2 diabetes mellitus patients: an analysis based on a large sample of 456,586 German T2DM patients. *J. Diabet. Complicat.* 29, 1015–1023.
- Zheng, Y., Ley, S.H., Hu, F.B., 2018. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat. Rev. Endocrinol.* 14, 88–98.