



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

COVID-19 and the consequences of isolating the elderly

As countries are affected by coronavirus disease 2019 (COVID-19), the elderly population will soon be told to self-isolate for “a very long time” in the UK, and elsewhere.¹ This attempt to shield the over-70s, and thereby protect over-burdened health systems, comes as worldwide countries enforce lockdowns, curfews, and social isolation to mitigate the spread of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).

However, it is well known that social isolation among older adults is a “serious public health concern” because of their heightened risk of cardiovascular, autoimmune, neuro-cognitive, and mental health problems.² Santini and colleagues³ recently demonstrated that social disconnection puts older adults at greater risk of depression and anxiety.

If health ministers instruct elderly people to remain home, have groceries and vital medications delivered, and avoid social contact with family and friends, urgent action is needed to mitigate the mental and physical health consequences.

Self-isolation will disproportionately affect elderly individuals whose only social contact is out of the home, such as at daycare venues, community centres, and places of worship. Those who do not have close family or friends, and rely on the support of voluntary services or social care, could be placed at additional risk, along with those who are already lonely, isolated, or secluded.

Online technologies could be harnessed to provide social support networks and a sense of belonging,⁴ although there might be disparities in access to or literacy in digital resources. Interventions could simply involve more frequent telephone contact with significant others, close family and friends, voluntary organisations,

or health-care professionals, or community outreach projects providing peer support throughout the enforced isolation. Beyond this, cognitive behavioural therapies could be delivered online to decrease loneliness and improve mental wellbeing.⁵

Isolating the elderly might reduce transmission, which is most important to delay the peak in cases, and minimise the spread to high-risk groups. However, adherence to isolation strategies is likely to decrease over time. Such mitigation measures must be effectively timed to prevent transmission, but avoid increasing the morbidity of COVID-19 associated with affective disorders. This effect will be felt greatest in more disadvantaged and marginalised populations, which should be urgently targeted for the implementation of preventive strategies.

We declare no competing interests.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

*Richard Armitage, Laura B Nellums
msxra37@nottingham.ac.uk

Division of Epidemiology and Public Health,
University of Nottingham, Nottingham NG5 1PB, UK

- 1 BBC. Coronavirus: isolation for over-70s ‘within weeks’. March 15, 2020. <https://www.bbc.co.uk/news/uk-51895873> (accessed March 16, 2020).
- 2 Gerst-Emerson K, Jayawardhana J. Loneliness as a public health issue: the impact of loneliness on health care utilization among older adults. *Am J Public Health*; 2015; **105**: 1013–19.
- 3 Santini Z, Jose P, Cornwell E, et al. Social disconnectedness, perceived isolation, and symptoms of depression and anxiety among older Americans (NSHAP): a longitudinal mediation analysis. *Lancet Public Health*; 2020; **5**: e62–70.
- 4 Newman M, Zainal N. The value of maintaining social connections for mental health in older people. *Lancet Public Health*; 2020; **5**: e12–13.
- 5 Käll A, Jägholm S, Hesse H, et al. Internet-based cognitive behavior therapy for loneliness: a pilot randomized controlled trial. *Behav Ther* 2020; **51**: 54–68.



Published Online
March 19, 2020
[https://doi.org/10.1016/S2468-2667\(20\)30061-X](https://doi.org/10.1016/S2468-2667(20)30061-X)