

Published in final edited form as:

Tob Control. 2019 June 19; 29(4): 367–373. doi:10.1136/tobaccocontrol-2018-054891.

Socioeconomic inequalities in children's exposure to tobacco retailing based on individual-level GPS data in Scotland

Fiona M Caryl¹, Niamh K Shortt², Jamie Pearce², Garth Reid³, Rich Mitchell¹

¹MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow, UK

²Centre for Research on Environment, Society and Health, School of GeoSciences, University of Edinburgh, Edinburgh, UK

³Department of Public Health Sciences, NHS Health Scotland, Edinburgh, UK

Abstract

Background—Identifying factors shaping knowledge of and attitudes toward tobacco products in pre-adolescence is a key component supporting tobacco control policies aimed at preventing smoking initiation. This study quantified exposure to tobacco retailing environments within the individual-level activity spaces of children across a socioeconomic gradient.

Methods—One week of GPS tracking data were collected at 10 second intervals from a nationally-representative sample of 10-11-year-olds (n=692). Proximity of GPS locations (n=~16M) to the nearest tobacco retailer (n=9030) was measured and exposure defined when a child came within 10m of a retailer. Duration, frequency, timing, and source of exposure were compared across income-deprivation quintiles, along with retail density within children's home neighbourhoods.

Results—On average, children were exposed to tobacco retailing for 22.7 minutes (95%CI 16.8—28.6) per week in 42.7 (35.2—50.1) independent encounters. However, children from the most deprived areas accumulated 6 times the duration and 7 times the frequency of exposure as children from the least deprived areas. Home neighbourhood retail densities were 2.6 times higher in deprived areas, yet the average number of businesses encountered did not differ. Most exposure came from convenience stores (35%) and newsagents (15%), with temporal peaks before and after school hours.

Conclusions—By accounting for individual mobility, we showed that children in socially disadvantaged areas accumulate higher levels of exposure to tobacco retailing than expected from disparities in home neighbourhood densities. Reducing tobacco outlet availability, particularly in areas frequently used by children, might be crucial to policies aimed at creating 'tobacco free' generations.

Contributors FC, NS, JP and RM designed the study. FC devised methodology, extracted data, conducted analysis, and wrote the manuscript. All authors contributed to draft revision and approved the final manuscript.

Background

There is growing acceptance that tobacco ‘endgame’ strategies—which seek to end, rather than control, the tobacco pandemic—are needed to reduce the global burden of preventable disease^{1–3}. Endgame goals vary internationally, but typically set a target for reducing smoking prevalence to less than 5% of the population⁴. A variety of tobacco-related interventions will be required to achieve these ambitions, and will almost certainly have to include measures designed to reduce the local supply of tobacco products⁴. Most adult smokers start during adolescence⁵, so mitigating against risk factors connected to smoking initiation during adolescence has been identified as a priority in tobacco control policies⁶. However, much of the research into the availability of tobacco products has focused on adults and adolescents^{7–12}, and less is known about exposure among younger children. This is a key omission because pre-adolescence is a significant formative period during which knowledge and attitudes to health-related behaviours, including smoking, become ‘hard-wired’¹³.

The availability of tobacco products has been identified as a potential causal factor in promoting smoking initiation and as a barrier to cessation^{14,15}. It is well established that tobacco retailing is disproportionately located in more socially deprived neighbourhoods^{16–20}, where smoking prevalence and premature deaths attributable to tobacco are also higher^{21,22}. Research suggests that ubiquitous availability of tobacco normalises and reinforces smoking in the local population, which in turn may make young people in the area more likely to become smokers themselves^{2,15,16}. Early smoking experience is strongly linked to later behaviour^{23–25}. Two-thirds of youths who initiate smoking aged 11 years become regular smokers versus less than half of those who initiate aged 16²⁶. Even a single smoking experience at age 11 is associated with an increased risk of smoking in the future compared with those who never smoked at this age²⁷. Hence early childhood interventions, such as those designed to de-normalise smoking behaviours by reducing tobacco availability in socially disadvantaged areas, should benefit disadvantaged children who are already more vulnerable to smoking²⁸.

Research linking exposure to tobacco retailing and youth smoking has typically quantified exposure within local neighbourhoods delimited using fixed areal units, such as census tracts, postcodes, or distance buffers from schools and/or homes^{12,17,19,29,30}. However, such methods are potentially biased by the areal units for which data are reported, and may not account for highly variable movements of individuals during their daily activities³¹. For example, measuring exposure within an individual’s residential neighbourhood can lead to considerable underestimates compared to those based on an individual’s daily movements^{32,33}. To overcome this, researchers are increasingly quantifying environmental exposures, such as to food or tobacco retail environments, within individual “activity spaces”, i.e. the set of locations visited in the course of daily activities and routes used to access them^{33–36}. Importantly, novel research linking individual-level mobility patterns to point-of-sale tobacco marketing exposure has revealed substantial differences in when and where individuals encounter tobacco^{35,36}. Kirchner et al. conclude that 1) fixed measures of exposure environments fail to account for differences in the mobility, preferences, and behaviour of individuals as they interact with the built environment; and 2) quantifying

individual-level exposure can identify previously unrecognized patterns of association among individual mobility, the built environment, and behavioural outcomes^{35,36}.

The focus of this study is Scotland where recent tobacco control policies—including banning point-of-sale tobacco product displays in shops; raising the legal purchase age to 18-years-old; and making it an offence to buy tobacco for under 18s—have led to significant declines in smoking in Scotland in the last decade^{37,38}. Adolescent smoking rates are at a historical low, with just 2% of 13-year-olds and 9% of 15-year-olds reporting regular smoking³⁹. However, rates of smoking in 13- and 15-year-olds remain higher in the most deprived areas^{37–39}. If the government’s aim of making Scotland tobacco-free by 2034 is to be achieved it is clear that further action to reduce inequalities in smoking is necessary³⁸.

In this paper, we determine if individual mobility patterns of children exacerbate exposure to tobacco retailing above what would be expected based on tobacco outlet density (TOD) alone. To achieve this, we provide a nationally representative assessment of daily exposure to tobacco retailing within the individual-level activity spaces of pre-adolescent children (n=692) in Scotland. One limitation highlighted by Kirchner et al. was that the low frequency of geospatial locations recorded (once every 15 minutes) in their study meant some exposures may have been missed, and exposure duration could not be estimated³⁶. Here, we use location data collected every ten seconds to quantify real-time exposure duration and make comparisons across area-level income deprivation quintiles. We calculated traditional measures of TOD in the home environment to determine if socioeconomic inequalities in exposure duration reflect those in TOD. In addition, we quantify the frequency of independent exposures, the number of unique retailers encountered per day, and the timing and source (i.e. outlet type) of exposures.

Methods

Calculating individual-level exposure of children to tobacco retailing took the following steps: i. geocoding tobacco retailer locations; ii. measuring proximity of children’s GPS locations to the nearest tobacco retailer; iii. calculating mean hourly exposure rates to derive daily and weekly rates for comparison across area-level deprivation quintiles.

Tobacco retail data

The addresses of all premises registered for tobacco sales in 2015–2016 were obtained from the Scottish Tobacco Retailers Register (n=9043) and cleaned to remove duplicates, resulting in 9030 premises. The longitude/latitude coordinates for each address were geocoded using the R package⁴⁰ `ggmap`⁴¹. Most addresses (91%) were geocoded to rooftop accuracy, but those that failed (n=830; 9%) were manually geocoded using Google Maps.

Neighbourhood deprivation

We obtained an indicator of socioeconomic deprivation for the data zone (a commonly used census data reporting unit comprising 500–1000 residents) containing each participant’s home address. The measure came from the Scottish Government’s Scottish Index of Multiple Deprivation (SIMD) 2016, a tool for measuring area-level deprivation. The SIMD is made from 7 domains that characterise social, economic and physical environment in the

area, ranging from education to crime. Following previous precedent, we used the income deprivation domain to measure area level deprivation¹⁹. This domain indicates the proportion of population in each area experiencing income deprivation as measured by receipt of means-tested benefits and government support. Eligibility for means tested benefits is based on income and savings, and benefits are used to top-up income if it is below a certain level.

Child activity space data

We used data from participants in the ‘Studying Physical Activity in Children’s Environments across Scotland’ (SPACES) study⁴², who were recruited from the Growing Up in Scotland (GUS) study—a nationally representative longitudinal cohort study originating in 2005. From a possible 2,402 children who participated in GUS sweep 8 interviews, 2,162 consented to be approached by SPACES researchers, of which 51% (n=1,096) consented to take part. Participants were provided with an accelerometer (ActiGraph GT3X+) and a GPS (QstarzSTARZ BT-Q1000XT; Qstarz International Co., Ltd, Taiwan) and asked to wear them over eight consecutive days between May 2015 and May 2016, when the participants were 10-11-years old. SPACES inclusion criteria required at least four weekdays of accelerometer data and 1 day of weekend data, resulting in a subset of 774 participants. Of these, 692 participants (381 female, 311 male) met our inclusion criteria of providing at least one hour of GPS data (Table 1).

Quantifying exposure

The straight-line distance from each GPS location to every retailer location was measured using the geosphere package⁴³ in R, and the nearest tobacco retailer retained along with information regarding retailer outlet type. Locations were classed as “exposed” when distance to nearest retailer was ≤ 10 m. The 10m threshold was used because this is the distance a child walking at 1m sec^{-1} (3.6kph) would travel between each GPS location. Each exposed location represented a 10-second epoch and duration of exposure in minutes was calculated by multiplying counts of locations by 10, then dividing by 60. The frequency of independent exposures was also quantified. Independent exposures occurred when an exposed location was preceded by an unexposed location and thus gives a measure of encounter rates with retailers. The unique identifier of retailers on the register was used to quantify the number of unique retailers encountered by participants.

Participants were asked to wear GPS devices during waking hours, leading to variation in wear time per day. To account for this, we standardised rates of exposure (duration and frequency) per hour of wear for weekdays and weekend days. Hourly exposure rates of each participant were then averaged to provide the mean hourly rate per day type per child. Mean hourly rates were multiplied by 16 hours to calculate the daily exposure in an average week or weekend day (0600-2200) for each participant. Rates were average across week/end day types and used to scale estimates per average week.

Comparison our sample with national level demographic distributions (Supplementary material) indicate slight under-representation of children from low-middle-income households (£10,000—£29,000) and the two most socially deprived quintiles (SIMD 1 and

2); and over-representation of high-income households (>£50,000) and the least socially deprived quintiles (4 and 5). However, after applying individual-level cross-sectional weights that were generated for all GUS respondents in sweep 8⁴², our sample could be considered nationally representative. Hourly exposure rates were weighted by each participant's unique weighting score and used as response variables in models against income-deprivation quintile.

Home environment TOD

We calculated home neighbourhood TOD as the number of tobacco outlets within 800m of each participant's geocoded home address⁹.

Data analysis

Mean weighted exposure rates (duration and frequency) of participants, home environment TOD, and mean and maximum number of unique retailers encountered were compared across income deprivation quintiles using one-way analysis of variance (ANOVA). Separate models were run for week days, weekend days, and average weeks. We controlled for season (winter: October—March) in all models, although 54-64% of participants in all income quintiles were tracked in winter (Table 1). All analyses were conducted in R using the `1me4` package⁴⁴. The proportion of total daily exposure per hour of day and the proportion of total daily exposure per retailer type were also quantified. Exposure by retailer type was compared against availability in the environment with chi-square tests, as was the distribution between most and least income deprivation quintiles. The distribution of exposure by time of day was compared between most and least income deprivation quintiles. All means are presented with 95% confidence intervals.

Results

A total 52,166 hours of GPS data were collected from 692 participants, with an average 63.0 hours (61.7—64.2) of wear time per participant across an average 6.0 (5.6—6.4) days of tracking, equalling an average 10.0 hours (9.9—10.1 hours) per participant per day (Table 1).

Duration and frequency of exposure to tobacco retailing

Our results showed that an average 10-11-year-old child was exposed to tobacco retailing for 2.7 minutes (1.9—3.4) per weekday and 4.7 minutes (3.4—5.9) per weekend day, totalling 22.7 minutes (16.8—28.6) per week (Table 2). However, a significant socioeconomic gradient existed in which children from the most income deprived areas experienced 5 times more exposure than children from the most affluent areas on weekdays, 6 times more on weekend days, and 6 times more in an average week ($P<0.001$: Table 2). An even greater disparity was apparent in the frequency of independent exposures (Table 3). While the average child encountered exposures 5.2 (4.2--6.1) times per weekday, 8.5 (6.9--10.2) time per weekend day, and 42.7 (35.2--50.1) times per week, children in the most income deprived areas encountered exposures 7 times more frequently per weekday and week than children in the least deprived areas (and 6 times on weekends: $P<0.001$: Table 3). The total

number of businesses encountered by each child was higher in the most deprived areas 6.7 (5.3—8.1) than the least deprived 6.0 (5.3—6.7), but not significantly so ($P=0.63$).

Tobacco outlet density in the home environment

The average number of retailers within 800m of participant's homes was 6.2 (5.6—6.7). Home environments of participants in the most deprived quintile had significantly more retailers (11.8; 10.1—13.4) than those in the least deprived areas (4.5; 3.7—5.2; $P<0.001$). The mean density in the most deprived areas was 2.6 times greater than that in the least deprived.

Source of exposure by outlet type

We found a significant difference between the distribution of exposure source across all income-deprivation levels and the availability of those sources in the environment ($P<0.001$). Overall, most exposure during a week came from convenience stores (35.0%) and newsagents (14.5%), although the level of exposure was roughly proportionate with the availability of these outlets (37.5% and 15.3%, respectively; Table 4). Exposure from supermarkets (9.8%) was significantly higher than expected given their availability (5.4%), particularly on weekends (13.6%). Exposure from off-licences, hotels, and businesses classed as "other retail" (e.g. discount stores) was also greater than expected given their availability.

We found significant differences between the distribution of exposure sources of children in the most deprived areas compared to those in the least deprived areas, and with their availability in the environment (both $P<0.001$). Children in deprived areas got significantly more exposure from convenience stores (41.0%) than children in the least deprived areas (28.1%). However, this reflected differences in the availability of convenience stores, which were 3 times more numerous in the most deprived areas ($n=929$) than the least ($n=306$). Children in deprived areas also got almost three times more exposure from supermarkets (13.2%), particularly on weekends (21.7%), than availability in these areas (4.8%) would predict. Children in deprived areas got less exposure from newsagents (12.7%) or public houses (3.9%) than expected given their availability (17.6% and 7.6%, respectively). Whereas, children from the least deprived areas got more exposure from these two sources (15.1% and 11.8%, respectively) than expected given their availability (11.1% and 9.7%, respectively).

Timing of exposures

Considerable peaks were seen in the timing of exposure for children from across all income deprivation levels. On weekdays, 46% of total exposure occurred after immediately school between 1500-1800, with 10% occurring before school between 0800-0900 (Figure 1a). Rates of exposure were reduced during school hours (0900-1500). On weekends, exposure was elevated between 1200-1700 when 59% of exposure occurred (Figure 1b).

Despite following a similar temporal trend, the hourly distribution of exposure was significantly different on weekdays and weekend days between children from income deprived and non-deprived areas (both $P<0.001$). The weekday morning (0800-0900) and

afternoon (1500-1600) peaks were higher among children from income-deprived areas. Weekend days also saw a higher peak in exposure during the hours 1200-1500 among those from income deprived areas compared to those from non-deprived areas.

Discussion

This is the first large-scale (n=692 participants) study to quantify exposure to tobacco retailing environments within the individual daily activity-spaces of pre-adolescent youths, and socioeconomic associations therein. As such, it represents a significant advancement in our understanding of how often tobacco retailers are encountered in an under-studied, yet key, demographic group. We found that an average 10-11-year old child in Scotland is exposed to tobacco retailing for 22.7 minutes (16.8—28.6) per week. Most notable, however, was the significant socioeconomic gradient in exposure, in which children from areas with the most income deprivation accumulated 6 times the duration, and 7 times the frequency, of exposure than children from areas with the least income deprivation. In other words, children in income deprived areas typically experienced more exposure in one weekend day (13.0 minutes: 5.8—20.2) as those from non-income deprived areas experienced in a whole week (11.3 minutes: 7.4—15.1). From a public health perspective, this is a concern given that exposure to tobacco products is a potential pathway to smoking initiation^{14,15}. It means that children from income deprived areas, who are already vulnerable to smoking initiation⁴⁵, experience the most exposure to tobacco products prior to adolescence, a critical period of addiction vulnerability⁴⁶. Additionally, the magnitude of the socioeconomic inequality in exposure revealed by our study is considerably larger than the 2.6-fold difference in tobacco retailer density in the home neighbourhood. This strongly suggests that static aerial measures, such as outlet density, may underestimate exposure inequalities compared with use of activity spaces that account for interactions between individual mobility and environment^{35,36}.

Simulation studies show that socioeconomic inequalities in smoking prevalence will persist in 2034 if the UK continues with “business as usual” tobacco control policies, with smoking rates of <3% in the upper income quintile smoking compared to 15% in the lowest income quintile⁴⁷. Radical actions are therefore required if the ‘tobacco free generation’ ambition is to be realised. Our results suggest that targeting policies to address the timing and type of retailer selling tobacco, or the spatial distribution of retailers, may be ways to reduce the gap. We found that a third of all exposure came from convenience stores, rising to over 40% in deprived areas, which reflected their availability. Exposure from supermarkets was disproportionate to availability across all income deprivation levels, particularly on weekends when children presumably accompany their parents grocery shopping. Interestingly, children from deprived areas got less exposure from newsagents, while the opposite was true for the least deprived, which may reflect differences in spending-power between quintiles. Clear temporal trends were also apparent, with peaks just before and after school hours on weekdays, and around midday into early afternoon on weekends. Extended exposure after the morning peak into school hours among those from income deprived areas may suggest the schools they attend have tobacco retailers close by.

Policy implications

Possible policy responses to our results are to prohibit sales of tobacco either in shops frequented regularly by children (e.g. convenience stores, newsagents, supermarkets), or at the times of day when children are more likely to visit (e.g. before and after school hours). Previous studies suggest that such policies may be heavily resisted, however. In a feasibility study to determine willingness of New Zealand convenience store owners to stop selling tobacco, or restrict hours of sale, almost all (93%) refused to do so voluntarily⁴⁸. This was primarily because tobacco is perceived as a key product for small local businesses for generating footfall⁴⁸. Reducing the availability of tobacco in communities may therefore require a combination of building public consensus and legislation to disincentivise retailers from selling tobacco products. Encouragingly, policy options such as banning sale of tobacco products near schools can be effective at reducing retailer density in lower income areas and reducing socioeconomic disparities while receiving strong public support^{49,50}. Determining policy interventions that are most effective in reducing overall exposure and socioeconomic inequalities is therefore a priority for future research.

Strengths and limitations

The main strength of our study lies in our quantifying individual-level exposure within child activity spaces using precise child and retailer location data from a large and nationally representative sample of children. This offers a significant advantage over previous studies adopting neighbourhood or density measures, which assume exposure by virtue of residential or school location. Collecting GPS data at 10-second intervals allowed us to quantify continuous real-time exposure, unlike previous studies quantifying exposure to tobacco retailing with GPS data collected at 15- or 30-minute intervals^{35,36}. Our methodology takes our understanding further by providing additional insight into the temporal distribution and the sources of exposure. Additionally, we now have a baseline of tobacco exposure for our sample who will be followed up longitudinally as part of GUS, allowing us to track their future smoking trajectories. Our use of an area-based measure of income deprivation also meant we were able to explore how differences in exposure are driven by the positive skew in retailer density towards more deprived areas.

Our study was limited, however, in that we do not know whether the children entered a shop or what the prominence and visibility of tobacco products was within shops. We also did not remove GPS locations at speeds indicative of travel by bicycle or motor-vehicle. We do not know how successive exposures accumulate and influence subliminally—or what a suitable threshold speed would be. Instead we assume that all exposure adds environmental cues to the social normalising process of tobacco availability. In addition, we know little of how a spatial concentration of outlets may relate to other smoking stimuli in the environment to further normalise smoking behaviours. Finally, children from income deprived areas were less well represented in the sample than those from less-deprived areas due to non-responses by those approached to be involved in the study.

Conclusions

Our study highlights how exposure can be more precisely quantified in tobacco studies to better understand everyday encounters with tobacco retailing. In doing so, our findings raise important questions regarding children's exposure to the tobacco retailing environment, and the significant inequalities therein. Understanding of the timing, frequency, duration, and source of tobacco retail exposure provides some of the evidence required to open the debate on tobacco retailing in Scotland. Reducing exposure through licensing, restricting sales in 'child spaces', or restricting sale times may become essential elements of a strategy to eliminate the tobacco epidemic.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors thank Linsay Gray, Rebecca Mancy, Jon Olsen, Laura MacDonald, and Natalie Nicholls for comments that greatly improved previous drafts of this manuscript.

Funding This research was funded by NHS Health Scotland. FC and RM are part of the Neighbourhoods and Communities Programme supported by the Medical Research Council (MC_UU_12017/10) and the Chief Scientist Office (SPHSU10).

References

1. McDaniel PA, Smith EA, Malone RE. The tobacco endgame: A qualitative review and synthesis. *Tob Control*. 2016; doi: 10.1136/tobaccocontrol-2015-052356
2. Malone RE. Tobacco endgames: what they are and are not, issues for tobacco control strategic planning and a possible US scenario. *Tob Control*. 2013; 22(suppl 1):i42–4. DOI: 10.1136/tobaccocontrol-2012-050820 [PubMed: 23591508]
3. Beaglehole R, Bonita R, Yach D, Mackay J, Reddy KS. A tobacco-free world: A call to action to phase out the sale of tobacco products by 2040. *Lancet*. 2015; 385(9972):doi: 10.1016/S0140-6736(15)60133-7
4. Moon G, Barnett R, Pearce J, Thompson L, Twigg L. The tobacco endgame: The neglected role of place and environment. *Health Place*. 2018; 53:271–278. DOI: 10.1016/j.healthplace.2018.06.012 [PubMed: 30238907]
5. U.S. Department of Health and Human Services. Preventing Tobacco Use Among Youth and Young Adults. A Report of the Surgeon General. 2012; doi: 10.1056/NEJMs1405092
6. WHO. WHO Report on the Global Tobacco Epidemic: Raising Taxes on Tobacco. 2015
7. Frohlich KL, Potvin L, Chabot P, Corin E. A theoretical and empirical analysis of context: Neighbourhoods, smoking and youth. *Soc Sci Med*. 2002; doi: 10.1016/S0277-9536(01)00122-8
8. Henriksen L, Feighery EC, Schleicher NC, Cowling DW, Kline RS, Fortmann SP. Is adolescent smoking related to the density and proximity of tobacco outlets and retail cigarette advertising near schools? *Prev Med (Baltim)*. 2008; 47(2):210–214. DOI: 10.1016/j.ypmed.2008.04.008
9. Lipperman-Kreda S, Grube JW, Friend KB. Local Tobacco Policy and Tobacco Outlet Density: Associations With Youth Smoking. *J Adolesc Heal*. 2012; 50(6):547–552. DOI: 10.1016/j.jadohealth.2011.08.015
10. Pearce J, Rind E, Shortt N, Tisch C, Mitchell R. Tobacco retail environments and social inequalities in individual-level smoking and cessation among Scottish adults. *Nicotine Tob Res*. 2016; 18(2):138–146. DOI: 10.1093/ntr/ntv089 [PubMed: 25895953]

11. Mennis J, Mason M, Way T, Zaharakis N. The role of tobacco outlet density in a smoking cessation intervention for urban youth. *Health Place*. 2016; 38:39–47. DOI: 10.1016/j.healthplace.2015.12.008
12. Schleicher NC, Johnson TO, Fortmann SP, Henriksen L. Tobacco outlet density near home and school: Associations with smoking and norms among US teens. *Prev Med (Baltim)*. 2016; 91:287–293. DOI: 10.1016/j.ypmed.2016.08.027
13. Marmot M, Friel S, Bell R, Houweling TA, Taylor S. Closing the gap in a generation: health equity through action on the social determinants of health. *Lancet*. 2008; doi: 10.1016/S0140-6736(08)61690-6
14. Pearce J, Barnett R, Moon G. Sociospatial inequalities in health-related behaviours: Pathways linking place and smoking. *Prog Hum Geogr*. 2012; 36(1):3–24. DOI: 10.1177/0309132511402710
15. Tunstall H, Shortt NK, Niedzwiedz CL, Richardson EA, Mitchell RJ, Pearce JR. Tobacco outlet density and tobacco knowledge, beliefs, purchasing behaviours and price among adolescents in Scotland. *Soc Sci Med*. 2018; 206:1–13. DOI: 10.1016/j.socscimed.2017.11.046 [PubMed: 29677578]
16. Loomis BR, Kim AE, Goetz JL, Juster HR. Density of tobacco retailers and its association with sociodemographic characteristics of communities across New York. *Public Health*. 2013; doi: 10.1016/j.puhe.2013.01.013
17. Robertson L, McGee R, Marsh L, Hoek J. A systematic review on the impact of point-of-sale tobacco promotion on smoking. *Nicotine Tob Res*. 2015; 17(1):2–17. DOI: 10.1093/ntr/ntu168 [PubMed: 25173775]
18. Wood L, Pereira G, Middleton N, Foster S. Socioeconomic area disparities in tobacco retail outlet density: a Western Australian analysis. *Med J Aust*. 2013; doi: 10.5694/mja1.2.11539
19. Shortt NK, Tisch C, Pearce J, et al. A cross-sectional analysis of the relationship between tobacco and alcohol outlet density and neighbourhood deprivation. *BMC Public Health*. 2015; 15(1):1014. doi: 10.1186/s12889-015-2321-1 [PubMed: 26437967]
20. Macdonald L, Olsen JR, Shortt NK, Ellaway A. Do 'environmental bads' such as alcohol, fast food, tobacco, and gambling outlets cluster and co-locate in more deprived areas in Glasgow City, Scotland? *Health Place*. 2018; 51:224–231. DOI: 10.1016/j.healthplace.2018.04.008
21. Taulbut, M, Gordon, D, M, K. Tobacco Smoking in Scotland: An Epidemiology Briefing. Edinburgh: NHS Health Scotland and Scottish Public Health Observatory; 2008.
22. The Scottish Government. *Scottish Health Survey 2016: Volume 1: Main*. 2016.
23. Patton GC, Carlin JB, Coffey C, Wolfe R, Hibbert M, Bowes G. The course of early smoking: A population-based cohort study over three years. *Addiction*. 1998; doi: 10.1046/j.1360-0443.1998.938125113.x
24. Choi WS, Gilpin EA, Farkas AJ, Pierce JP. Determining the probability of future smoking among adolescents. *Addiction*. 2001; doi: 10.1046/j.1360-0443.2001.96231315.x
25. Jackson C, Dickinson D. Cigarette consumption during childhood and persistence of smoking through adolescence. *Arch Pediatr Adolesc Med*. 2004; doi: 10.1001/archpedi.158.11.1050
26. Lynch, BS, Bonnie, RJ. *Growing up Tobacco Free: Preventing Nicotine Addiction in Children and Youths*. Washington DC: National Academy Press; 1994.
27. Fidler JA, Wardle J, Brodersen NH, Jarvis MJ, West R. Vulnerability to smoking after trying a single cigarette can lie dormant for three years or more. *Tob Control*. 2006; 15(3):205–209. DOI: 10.1136/tc.2005.014894 [PubMed: 16728751]
28. Purcell KR, O'Rourke K, Rivis M. Tobacco control approaches and inequity-how far have we come and where are we going? *Health Promot Int*. 2015; doi: 10.1093/heapro/dav075
29. Shortt NK, Tisch C, Pearce J, Richardson EA, Mitchell R. The density of tobacco retailers in home and school environments and relationship with adolescent smoking behaviours in Scotland. *Tob Control*. 2016; 25(1):75–82. DOI: 10.1136/tobaccocontrol-2013-051473 [PubMed: 25370699]
30. Finan LJ, Lipperman-Kreda S, Abadi M, et al. Tobacco outlet density and adolescents' cigarette smoking: a meta-analysis. *Tob Control*. 2018; 0:1–7. DOI: 10.1136/tobaccocontrol-2017-054065

31. Fotheringham, AS, Brunson, C, Charlton, M. Quantitative Geography : Perspectives on Spatial Data Analysis. SAGE Publications; 2000. https://books.google.co.uk/books/about/Quantitative_Geography.html?id=semXiMy6T-ToC&redir_esc=y [Accessed April 10, 2019]
32. Shareck M, Kestens Y, Vallée J, Datta G, Frohlich KL. The added value of accounting for activity space when examining the association between tobacco retailer availability and smoking among young adults. *Tob Control*. 2016; 25(4):406–412. DOI: 10.1136/tobaccocontrol-2014-052194 [PubMed: 26032269]
33. Lipperman-Kreda S, Morrison C, Grube JW, Gaidus A. Youth activity spaces and daily exposure to tobacco outlets. *Heal Place*. 2015; doi: 10.1016/j.healthplace.2015.03.013
34. Christian WJ. Using geospatial technologies to explore activity-based retail food environments. *Spat Spatiotemporal Epidemiol*. 2012; 3(4):287–295. DOI: 10.1016/J.SSTE.2012.09.001 [PubMed: 23149325]
35. Kirchner, TR; Vallone, D; Cantrell, J; , et al. Individual mobility patterns and real-time geo-spatial exposure to point-of-sale tobacco marketing. *Proceedings of ACM Wireless Health*; New York NY: ACM; 2012. 1–8. 2012
36. Kirchner TR, Cantrell J, Anesetti-Rothermel A, Ganz O, Vallone DM, Abrams DB. Geospatial Exposure to Point-of-Sale Tobacco Real-Time Craving and Smoking-Cessation Outcomes. *Am J Prev Med*. 2013; 45(4):379–385. DOI: 10.1016/j.amepre.2013.05.016 [PubMed: 24050412]
37. Mclean J, Christie S, Hinchliffe S, et al. The Scottish Health Survey 2017 Edition. Volume: 1 Main Report. 2017; 1
38. Reid, G; Rennick, L; Laird, Y; Arnot, J; Mcateer, J. [Accessed April 19, 2019] Review of ‘Creating a Tobacco-Free Generation: A Tobacco Control Strategy for Scotland’. 2017. <http://www.healthscotland.scot/media/1545/review-of-creating-a-tobacco-free-generation-a-tobacco-control-policy-for-scotland.pdf>
39. NHS. [Accessed April 8, 2019] Scottish Schools Adolescent Lifestyle and Substance Use Survey (SALSUS) National Report: Smoking, Drinking and Drug Use among 13 and 15 Year Olds in Scotland in 2013. 2015. https://www.isdscotland.org/Health-Topics/Public-Health/Publications/2014-11-25/SALSUS_2013_Smoking_Report.pdf
40. R Development Core Team R, R Core Team. R: A language and environment for statistical computing. *R A Lang Environ Stat Comput*. 2017; doi: 10.1016/j.jssas.2015.06.002
41. Kahle D, Wickham H. ggmap : Spatial Visualization with ggpl. *R J*. 2013; doi: 10.1023/A:1009843930701
42. Mccrorie P, Mitchell R, Ellaway A. Comparison of two methods to assess physical activity prevalence in children: an observational study using a nationally representative sample of Scottish children aged 10-11 years. *BMJ Open*. 2018; 8doi: 10.1136/bmjopen-2017-018369
43. Hijmans RJ, Williams E, Vennes C. geosphere: Spherical Trigonometry. R package version 1.2–28. *Packag Geosph*. 2012
44. Bates, D, Maechler, M, Bolker, B. , et al. Package “lme4”. *R Found Stat Comput*; Vienna, Austria: 2018.
45. Levin KA, Dundas R, Miller M, McCartney G. Socioeconomic and geographic inequalities in adolescent smoking: A multilevel cross-sectional study of 15 year olds in Scotland. *Soc Sci Med*. 2014; 107:162–170. DOI: 10.1016/j.socscimed.2014.02.016 [PubMed: 24607678]
46. Chambers RA, Taylor JR, Potenza MN. Developmental Neurocircuitry of Motivation in Adolescence: A Critical Period of Addiction Vulnerability. *Am J Psychiatry*. 2003; 160(6):1041–1052. DOI: 10.1176/appi.ajp.160.6.1041 [PubMed: 12777258]
47. Hunt D, Knuchel-Takano A, Jaccard A, et al. Modelling the implications of reducing smoking prevalence: the public health and economic benefits of achieving a ‘tobacco-free’ UK. *Tob Control*. 2017; 27(2)doi: 10.1136/tobaccocontrol-2016-053507
48. Paynter J, Glover M, Bullen C, Sonia D. An intervention to reduce the number of convenience stores selling tobacco: feasibility study. *Tob Control*. 2016; 25(3):319–324. DOI: 10.1136/tobaccocontrol-2014-052045 [PubMed: 25873646]
49. Ribisl KM, Luke DA, Bohannon DL, Sorg AA, Moreland-Russell S. Reducing disparities in tobacco retailer density by banning tobacco product sales near schools. *Nicotine Tob Res*. 2017; 19(2):239–244. DOI: 10.1093/ntr/ntw185 [PubMed: 27613900]

50. Whyte G, Gendall P, Hoek J. Advancing the retail endgame: public perceptions of retail policy interventions. *Tob Control*. 2014; 23(2):160–166. DOI: 10.1136/tobaccocontrol-2013-051065 [PubMed: 23842946]

What this study adds

This study is significant because it reveals how much greater socioeconomic disparities in tobacco retail exposure become when individual mobility is accounted for. By implementing cutting-edge methodology for measuring continuous real-time exposure to tobacco retailing we were able to identify socioeconomic inequalities of greater magnitude than disparities in neighbourhood measures of density would indicate. This forms a significant contribution to the policy debate on tobacco availability. Our findings highlight a need to take interactions between individual patterns of mobility and the retail environment into account when considering any supply-side intervention. However, the observed socio-economic gradient in exposure (as measured by income deprivation level) suggests that any moves to either reduce retail outlets, or restrict time of sales, will have a greater impact on, and indeed benefit to, more deprived income groups who suffer the greatest amount of tobacco-related harm.

Table 1
Unweighted sociodemographic characteristics and summary of GPS data of 692 study participants.

	Income deprivation quintile					
	Overall	1 (Most Deprived)	2	3	4	5 (Least Deprived)
Sex: male	311 (45%)	26 (44%)	33 (39%)	58 (41%)	85 (46%)	109 (48%)
Sex: female	381	33	52	82	98	116
Season: winter	450 (63%)	38 (64%)	59 (69%)	76 (54%)	106 (58%)	151 (67%)
Season: summer	262	21	26	64	77	74
Urban: 1	176 (25%)	18 (31%)	20 (24%)	17 (12%)	37 (20%)	84 (37%)
2	248	36	42	48	48	74
3	83	1	10	24	17	31
4	20	2	2	8	6	2
5	106	2	6	20	48	30
Rural: 6	59 (9%)	0 (0%)	5 (6%)	23 (16%)	27 (15%)	4 (2%)
Tracking effort: wear hours (mean ± 95% CI)	63.0 (61.7--64.2)	57.9 (53.4--62.4)	58.0 (53.6--62.4)	65.0 (62.4--67.6)	63.9 (61.5--66.3)	64.1 (62.0--66.3)
Tracking effort: wear days (mean ± 95% CI)	6.0 (5.6--6.4)	6.0 (5.8--6.3)	6.3 (6.1--6.5)	6.1 (6.0--6.3)	6.2 (6.1--6.4)	6.2 (6.1--6.3)

Table 2
Mean duration of exposure per average day and week with 95% confidence intervals in parenthesis.

Income deprivation quintiles	Weekday	Weekend	Week
All income levels	2.7 (1.9--3.4)	4.7 (3.4--5.9)	22.7 (16.8--28.6)
1 (most deprived)	7.3 (4.6--10.0)	13.0 (5.8--20.2)	63.4 (38.7--88.1)
2	5.8 (1.9--9.7)	9.2 (4.1--14.3)	45.6 (17.6--73.7)
3	2.4 (0.1--4.7)	4.5 (1.1--7.9)	21.1 (2.5--39.8)
4	1.5 (0.9--2.2)	3.1(1.5--4.7)	14.0 (9.3--18.7)
5 (least deprived)	1.4 (0.8--1.9)	2.2 (1.4--3.0)	11.3 (7.4--15.1)
<i>ANOVA</i>	<i>P < 0.001</i>	<i>P < 0.001</i>	<i>P < 0.001</i>

Table 3
Mean frequency of independent exposures per day and week with 95% confidence intervals in parenthesis.

Income deprivation quintiles	Weekday	Weekend	Week
All income levels	5.2 (4.2--6.1)	8.5 (6.9--10.2)	42.7 (35.2--50.1)
1 (most deprived)	18.1 (11.6--24.5)	27.3 (15.3--39.3)	149.2 (96.5--201.9)
2	8.2 (5.1--11.3)	12.9 (7.8--17.9)	63.3 (42.8--83.8)
3	3.4 (1.6--5.2)	7.0 (3.1--10.9)	30.5 (14.8--46.2)
4	4.0 (2.2--5.8)	5.9 (4.1--7.8)	32.5 (20.3--44.7)
5 (least deprived)	2.7 (2.0--3.4)	5.0 (3.8--6.3)	22.8 (18.4--27.3)
<i>ANOVA</i>	<i>P < 0.001</i>	<i>P < 0.001</i>	<i>P < 0.001</i>

Table 4
The percentage of independent exposures by retailer type and availability of retailer types by income deprivation.

Retailer type	All income quintiles			Most deprived income quintile			Least deprived income quintile					
	Weekday	Weekend	Week Availability	Weekday	Weekend	Week Availability	Weekday	Weekend	Week Availability			
Convenience Store	40.9	25.5	35.0	37.5	45.4	34.6	41.0	42.9	34.3	18.6	28.1	35.8
Newsagent	14.5	14.6	14.5	15.3	15.1	9.2	12.7	17.6	14.1	16.8	15.1	11.1
Public House	9.2	12.3	10.4	10.6	5.1	2.0	3.9	7.6	10.4	14.0	11.8	9.7
Supermarket	7.5	13.6	9.8	5.4	7.3	21.7	13.2	4.8	11.0	16.6	13.2	7.3
Off-licence	8.1	8.8	8.4	5.9	9.3	10.8	9.9	8.7	6.4	5.6	6.1	4.8
Hotel	5.9	5.3	5.7	3.9	0.4	0.4	0.4	0.6	10.3	6.4	8.7	8.0
Other retail	4.2	7.1	5.3	4.2	6.4	9.2	7.6	4.5	3.5	7.0	4.9	2.3
Forecourt Garage	3.6	5.9	4.5	6.9	4.6	7.4	5.7	3.9	3.8	4.1	3.9	10.2
Other catering	2.8	2.5	2.7	4.2	4.0	2.2	3.3	5.4	2.6	3.8	3.1	3.5
Restaurant	0.6	1.4	0.9	1.2	0.3	0.4	0.3	1.1	0.3	2.9	1.3	1.8
Nightclub	0.7	0.8	0.7	0.8	0.4	1.2	0.8	0.4	0.6	0.8	0.7	0.9
Entertainment venue	0.6	0.6	0.6	1.2	0.7	0.2	0.5	0.8	0.8	1.4	1.0	2.1
Private Club	0.5	0.5	0.5	1.1	0.3	0.4	0.3	0.7	1.1	0.8	1.0	1.3
Specialist tobacconists	0.2	0.3	0.2	0.3	0.3	0.0	0.2	0.3	0.5	0.2	0.4	0.0
Sports Club	0.3	0.1	0.2	0.9	0.3	0.2	0.3	0.5	0.3	0.2	0.3	1.3
Mobile trader	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0