



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to the Editor

Active smoking is not associated with severity of coronavirus disease 2019 (COVID-19)



ARTICLE INFO

Keywords:

Smoking

Cigarette

Coronavirus

COVID-19

In the ongoing coronavirus disease 2019 (COVID-19) pandemic, some unique clinical features have been described [1]. In a report of 44,672 cases from China, a case fatality rate of 2.8% for males versus only 1.7% for females was observed [2]. It has been hypothesized that this may be due a higher prevalence of comorbidities among males, in particular the rate of smoking, which is reported to be 52.1% in men and 2.7% among women in China [3]. However, decreased levels of angiotensin converting enzyme 2 (ACE2), the reported host receptor of the virus responsible of COVID-19 (severe acute respiratory syndrome coronavirus 2; SARS-CoV-2), are observed in smokers [4,5]. In this article, we aimed to investigate the association between active smoking and severity of COVID-19 illness.

An electronic search was conducted in Medline (PubMed interface), Scopus and Web of Science, using the keywords “smoking” OR “cigarette” AND “coronavirus 2019” OR “COVID-19” OR “2019-nCoV” OR “SARS-CoV-2”, between 2019 and present time (i.e., March 9, 2020), without language restriction. The title, abstract and full text of all documents captured with these search criteria were scrutinized, and

those reporting the rate of active smokers in COVID-19 patients with clinically validated definition of severe disease were included in this meta-analysis. The reference list of the identified studies was also analyzed (forward and backward citation tracking) for detecting other potentially eligible articles.

A meta-analysis was then performed, with estimation of the odds ratio (OR) and its 95% confidence interval (95% CI) in patients with or without severe forms of COVID-19. The statistical analysis was carried out using MetaXL, software Version 5.3 (EpiGear International Pty Ltd., Sunrise Beach, Australia). The study was carried out in accordance with the declaration of Helsinki and with the term of local legislation.

Overall, 27 documents could be initially identified based on our search criteria and from the reference lists, 22 of which were excluded after title, abstract or full text reading, since they were review articles ($n=4$), commentaries or other editorial materials ($n=2$), they did not deal with COVID-19 disease ($n=11$), or did not provided the rate of active smokers in patients with or without severe disease ($n=5$). Therefore, 5 studies could finally be included in our meta-analysis,

Table 1
Characteristics of the studies included.

Authors	Setting	Sample size	Outcome	Severe patients			Non-severe patients		
				n (%)	Age (yrs)*	Women (%)	n (%)	Age (yrs)*	Women (%)
Guan W et al., 2020 to ICU, mechanical ventilation, death.	China	1085	Admission						
	172 (16%)	52 (40–65)	42%	913 (84%)	45 (34–57)	42%			
Huang C et al., 2020 Liu W et al., 2020 to ICU, mechanical ventilation, death.	China	44	ICU Care	13 (30%)	49 (41–61)	15%	31 (70%)	49 (41–58)	32%
	China	78	Admission						
	11 (14%)	66 (51–70)	36%	67 (86%)	37 (32–41)	52%			
Yang X et al., 2020 Zhang JJ et al., 2020	China	50	Death	32 (64%)	65 (45–87)	34%	18 (36%)	52 (26–77)	30%
	China	142	Respiratory distress/ insufficiency	60 (42%)	64 (25–87)	43%	82 (58%)	52 (26–78)	54%

* Median and interquartile range.




Fig. 1. Odds ratio (OR) and 95% confidence interval (95% CI) of active smoking between coronavirus disease 2019 (COVID-19) patients with or without severe disease. .

totaling 1399 COVID-19 patients, 288 of whom (20.6%) with severe disease [6–10]. The essential characteristics of the five included studies are shown in Table 1, whilst the individual and pooled OR of smoking for predicting severe COVID-19 is shown in Fig. 1. Overall, in only one study [8] active smoking was found to be a significant predictor of COVID-19 severity, whilst in the other four studies the association was not statistically significant. Despite a trend towards higher risk was appreciable, no significant association could neither be found between active smoking and severity of COVID-19 when data of individual studies were pooled (OR, 1.69; 95% CI, 0.41–6.92; $p=0.254$). No significant association could also be found when the large study by Guan et al. (89.5% of all samples size) [6] was excluded from statistical analysis (OR, 4.35; 95% 0.86–21.86; $p=0.129$; I^2 29%, $p=0.24$).

In conclusion, the results of this preliminary meta-analysis based on Chinese patients suggest that active smoking does not apparently seem to be significantly associated with enhanced risk of progressing towards severe disease in COVID-19.

Declaration of Competing Interest

None

References

- [1] Mattiuzzi C, Lippi G. Which lessons shall we learn from the 2019 novel coronavirus outbreak? *Ann Transl Med* 2020;8:48.
- [2] The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The

- epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) — China. 2020. *China CDC Wkly* 2020;2:113–22.
- [3] Parascandola M, Xiao L. Tobacco and the lung cancer epidemic in China. *Transl Lung Cancer Res* 2019;8(Suppl 1):S21–30. <https://doi.org/10.21037/tlcr.2019.03.12>.
- [4] Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor recognition by novel coronavirus from wuhan: an analysis based on decade-long structural studies of sars. *J Virol* 2020 Jan 29. <https://doi.org/10.1128/JVI.00127-20>. pii: JVI.00127-20[Epub ahead of print].
- [5] Oakes JM, Fuchs RM, Gardner JD, Lazartigues E, Yue X. Nicotine and the renin-angiotensin system. *Am J Physiol Regul Integr Comp Physiol* 2018;315(5):R895–906.
- [6] Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in china. *N Engl J Med* 2020 Feb 28. <https://doi.org/10.1056/NEJMoa2002032>. [Epub ahead of print].
- [7] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in wuhan. *China. Lancet* 2020;395:497–506.
- [8] Liu W, Tao Z-W, Lei W, Ming-Li Y, Kui L, Ling Z, et al. Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chin Med J February* 2020. <https://doi.org/10.1097/CM9.0000000000000775>.
- [9] Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in wuhan, China: a single-centered, retrospective, observational study. *The Lancet Respiratory Medicine* 2020;0(0). [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5).
- [10] Zhang JJ, Dong X, Cao YY, Yuan YD, Yang YB, Yan YQ, et al. Clinical characteristics of 140 patients infected by SARS-CoV-2 in wuhan, China. *Allergy* 2020 Feb 19. <https://doi.org/10.1111/all.14238>. [Epub ahead of print].

Giuseppe Lippi^{a,*}, Brandon Michael Henry^b

^a Section of Clinical Biochemistry Department of Neuroscience Biomedicine and Movement, University of Verona, Verona, Italy

^b Cardiac Intensive Care Unit The Heart Institute Cincinnati Children's Hospital Medical Center, Ohio, United States

E-mail address: giuseppe.lippi@univr.it (G. Lippi).

* Corresponding author. Section of Clinical Biochemistry, University Hospital of Verona, Piazzale LA Scuro, 37134 Verona, Italy