

# Characterization of cold stress responses in different rapeseed ecotypes based on metabolomics and transcriptomics analyses

Hongju Jian\*, Ling Xie\*, Yanhua Wang, Yanru Cao, Mengyuan Wan, Dianqiu Lv, Jiana Li, Kun Lu, Xinfu Xu and Liezhao Liu

College of Agronomy and Biotechnology, Southwest University, Chongqing, China

\* These authors contributed equally to this work.

## ABSTRACT

The winter oilseed ecotype is more tolerant to low temperature than the spring ecotype. Transcriptome and metabolome analyses of leaf samples of five spring *Brassica napus* L. (*B. napus*) ecotype lines and five winter *B. napus* ecotype lines treated at 4 °C and 28 °C were performed. A total of 25,460 differentially expressed genes (DEGs) of the spring oilseed ecotype and 28,512 DEGs of the winter oilseed ecotype were identified after cold stress; there were 41 differentially expressed metabolites (DEMs) in the spring and 47 in the winter oilseed ecotypes. Moreover, more than 46.2% DEGs were commonly detected in both ecotypes, and the extent of the changes were much more pronounced in the winter than spring ecotype. By contrast, only six DEMs were detected in both the spring and winter oilseed ecotypes. Eighty-one DEMs mainly belonged to primary metabolites, including amino acids, organic acids and sugars. The large number of specific genes and metabolites emphasizes the complex regulatory mechanisms involved in the cold stress response in oilseed rape. Furthermore, these data suggest that lipid, ABA, secondary metabolism, signal transduction and transcription factors may play distinct roles in the spring and winter ecotypes in response to cold stress. Differences in gene expression and metabolite levels after cold stress treatment may have contributed to the cold tolerance of the different oilseed ecotypes.

Submitted 26 September 2019

Accepted 6 February 2020

Published 31 March 2020

Corresponding author

Liezhao Liu, liezhao@swu.edu.cn

Academic editor

Ana Ribeiro-Barros

Additional Information and  
Declarations can be found on  
page 23

DOI 10.7717/peerj.8704

© Copyright

2020 Jian et al.

Distributed under

Creative Commons CC-BY 4.0

**OPEN ACCESS**

**Subjects** Agricultural Science, Plant Science

**Keywords** Cold stress, *Brassica napus*, Metabolome, Transcriptomics, Molecular mechanisms

## INTRODUCTION

As one of the most important environmental stresses, cold stress significantly affects plant growth, development and distribution, and it can be classified as chilling (<20 °C) and freezing (<0 °C) temperatures (Chinnusamy, Zhu & Zhu, 2007; Miura & Furumoto, 2013). Cell membrane stiffness, protein complex instability and photosynthesis weakness are detected under chilling stress, and much more serious injuries occur under freezing stress conditions. Many physiology and biochemistry processes can be influenced by cold stress, which is dependent on the plant species, stage of plant development, and length of the stress period. Many plant species, including *Brassica napus* L. (*B. napus*), have the ability to

acclimate to the cold. However, this ability can be increased after exposure to chilling stress, and the capacity of acclimation varies greatly among species. Cold acclimation mechanisms have been fully explored and have expanded our understanding of *Arabidopsis thaliana* (*A. thaliana*), while information regarding the molecular basis of this process in *B. napus* is largely lacking despite their similar genetics.

Numerous cold-regulated genes have been identified in diverse species, including *A. thaliana* (Zhao *et al.*, 2009), *Oryza sativa* (Chen *et al.*, 2003; Maruyama *et al.*, 2014; He *et al.*, 2016; Xiaochuang *et al.*, 2017), *Zea mays* (Bilska-Kos, Szczepanik & Sowinski, 2016; Riva-Roveda *et al.*, 2016), *Prunus persica* (Sanhueza *et al.*, 2015; Bustamante *et al.*, 2016), and many other species (Khodakovskaya *et al.*, 2006; Kosova *et al.*, 2012; Tian *et al.*, 2013; Zhao *et al.*, 2014; Liu *et al.*, 2017), using RNA sequencing (RNA-Seq) technology. The ICE1-CBF-COR genetic network response to cold acclimation is the best understood pathway in *Arabidopsis* (Cook *et al.*, 2004; Chinnusamy, Zhu & Zhu, 2007). C-repeat/dehydration responsive element (DREB)-binding factors (e.g., CBF1, -2, -3, -4, -6) belonging to the AP2/ERF domain family are significantly induced in response to cold stress (Chinnusamy *et al.*, 2003, 2007; Cook *et al.*, 2004). In addition, other genes involved in cold stress response have also characterized using transcriptome analysis (Kaplan *et al.*, 2004; Wu *et al.*, 2016). Genes involved in lipid metabolism (Welti *et al.*, 2002; Testerink & Munnik, 2005; D'Angeli & Altamura, 2016), ABA biosynthesis and signal transduction (Lee, Henderson & Zhu, 2005; Chen *et al.*, 2014), Ca<sup>2+</sup> signal transduction (Miura & Furumoto, 2013; Shi, Ding & Yang, 2015), transcription factors (TFs) (Lindemose *et al.*, 2013) and various hormones have been detected after cold stress (Stuempfle, Nindl & Kamimori, 2010). All of these findings have expanded our understanding of molecular mechanisms in response to cold stress tolerance (Chinnusamy, Zhu & Zhu, 2007). Nevertheless, many cold stress phenotypes cannot be explained by the functions of a single gene, highlighting the need to identify additional regulatory networks and pathways, including key metabolites.

In addition to the very sophisticated gene regulatory networks, plants have evolved various complex metabolite response patterns in the low temperature stress response (Kaplan *et al.*, 2004). Increasing evidence has shown that gene expression and biochemical pathways are altered under cold stress conditions. Sugars, lipids and amino acids, for example, have been widely reported to be involved in the cold stress response, not only for protein synthesis process but as precursors of other key metabolites in the response to cold stress (Cook *et al.*, 2004). As key functions of the metabolite response to cold stress, many more kinds of metabolites that are responsive to cold stress must be characterized in further studies.

Transcriptome sequencing technology has matured and can provide very efficient and high-throughput screening of low temperature response genes. In previous studies, genes involved in the cell wall and membrane stabilization, calcium signaling, hormones, TFs and mitogen-activated protein kinase (MAPK) pathways, and soluble sugar and protein biosynthesis and metabolic pathways were significantly altered after cold stress using RNA-Seq technologies (Wang *et al.*, 2013a, 2013b; Chen *et al.*, 2014; Ren *et al.*, 2014; Fu *et al.*, 2016; Gong *et al.*, 2018). Recent technological advances have made it possible to

explore the molecular mechanisms underlying the response to low temperature stress using transcriptome and metabolomics techniques (Kaplan *et al.*, 2007; Maruyama *et al.*, 2014; Wu *et al.*, 2016; Zhang *et al.*, 2016). By integrating transcriptome and metabolome data, a broader and more comprehensive understanding of genes and metabolites co-regulation of low temperature stress in Arabidopsis has been performed (Kaplan *et al.*, 2007). In rice, genes encoding proteins involved in sucrose-starch metabolism and the glyoxylate cycle were upregulated, and these changes were correlated with the accumulation of glucose, fructose and sucrose in rice after exposure to cold or dehydration, as revealed by joint metabolite, phytohormone, and gene transcript analyses (Maruyama *et al.*, 2014). In *A. thaliana*, flavonoids are regarded as determinants of freezing tolerance and cold acclimation using 20 mutant lines based on qRT-PCR, LC-MS and GC-MS analysis (Schulz *et al.*, 2016). Carboxylates, amino acids, carbohydrates and sugar alcohols were also made contributing to reduce freezing tolerance in the *pgm* mutant (Hoermiller *et al.*, 2017). Joint RNA-Seq and metabolomics analyses have revealed a reactive oxygen species-dominated dynamic model response to chilling stress in two subspecies of Asian cultivated rice (*Oryza sativa*) with significant divergence in chilling tolerance (Zhang *et al.*, 2016). Integrative analysis of transcripts and metabolites has revealed that many transcripts and metabolites are rearranged to cope with low temperature stress in *Dendrobium officinale* (Wu *et al.*, 2016).

As the third largest oil crop in the world, rapeseed is the most important sources of vegetable oil in China. Rapeseed is the only overwintering oil crop in China. The research and breeding of cold-resistant varieties play an important role in the safety of edible oil in China. Until now, many valuable studies have been performed to examine the morphology, physiology and molecular biology of cold resistance (Burbulis *et al.*, 2010; Chen *et al.*, 2011), but few studies have addressed the cold resistance regulatory network mechanisms at the transcriptome and metabolome levels in rapeseed. ABA and IP3/Ca<sup>2+</sup> signal transduction pathways have been identified as key actors in response to low temperature in rapeseed (Xian *et al.*, 2017). To our knowledge, the elucidation of candidate genes or the metabolite response to low temperature in rapeseed using metabolomics and transcriptomics analysis is still very lacking.

In this study, we aim to explore the molecular mechanisms in different rapeseed ecotypes at transcriptome and metabolome levels from a wide perspective and seek clues linking genes and metabolites in response to low temperature stress. Each rapeseed ecotype (spring and winter) containing five lines were used in our study. The spring rapeseed ecotype was more sensitive to low temperature than the winter rapeseed ecotype. Many cold response genes and metabolites were detected, and common ones had greater degrees of change in the winter than in the spring rapeseed ecotypes.

## MATERIALS AND METHODS

### Plant materials and cold treatments

Five winter (N5, N7, N36, N49 and N60) and five spring ecotypes (M10, M11, N240, N244 and N249) of *B. napus* were used in this study. Seeds of each line were germinated and cultivated in a petri dish under normal conditions. After 10 days, the plants were moved to

plastic pots in a greenhouse with a long-day photoperiod (28 °C with 16 h light and 24 °C with 8 h dark) until the plants grew to the five- or six-leaf stage. Half plants of each line were grown under normal conditions and used as control plants. The remaining plants of each line were moved to 4 °C for 12 h with the same photoperiod. Leaves of control and treatment plants were harvested at 0 and 12 h after cold stress. All leaves were immediately frozen in liquid nitrogen and stored at –80 °C until use. Every sample consisted of leaves from more than five plants grown under the same conditions. Three biological replicates were performed, and 12 samples including control samples for winter (28W) and spring (28S) ecotypes, cold stress samples for winter (4W) and spring (4S) ecotypes were obtained and used for RNA-Seq and metabolic profiling.

### RNA extraction and RNA-Seq

Total RNA of all samples was extracted using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. An Agilent 2100 Bioanalyzer (Agilent, Santa Clara, CA, USA) and a spectrophotometer (NanoDrop, Wilmington, DE, USA) were used to detect the integrity, quality and quantity of total RNA. Then, high-quality RNA was sent to Genedenovo Biotechnology Co., Ltd. (Guangzhou, China) and sequenced on the Illumina sequencing platform. Briefly, mRNA was extracted using dynabeads oligo (dT) and fragmented by fragmentation buffer. Double-stranded cDNAs were synthesized using reverse-transcriptase and random hexamer primers, and the cDNA fragments were purified using a QIA quick PCR extraction kit. These purified fragments were washed with EB buffer for end reparation of poly (A) addition and then ligated to sequencing adapters. Following agarose gel electrophoresis and extraction of cDNA from gels, the cDNA fragments were purified and enriched by PCR to construct the final cDNA library. The cDNA library was sequenced on the Illumina sequencing platform (Illumina HiSeq™ 2500) using the paired-end technology.

### Sequence processing and analysis

After removal of adapter and low quality sequences, clean reads were obtained and mapped to the *B. napus* reference genome (<http://www.genoscope.cns.fr/brassicapapus/>). They were then assembled using TopHat 2.0.0 and DEGseq. FPKM (fragments per kilobase of exon per million mapped fragments) was used to calculate gene expression levels, and differentially expressed genes (DEGs) were screened with Cuffdiff using the following two criteria: (i) false discovery rate *P*-value correction of <0.05 and (ii)  $|\log_2(\text{fold change})| > 1$  (Trapnell *et al.*, 2010). Based on blastp analysis, homologs of TFs and hormone genes in *B. napus* were screened from our previous study (Jian *et al.*, 2018), and possible TF and plant hormone genes among these DEGs were also obtained. In addition, all sequencing data were uploaded to NCBI under accession number PRJNA530291.

### qRT-PCR analysis

Here, 1 µg of total RNA was used for cDNA synthesis using a Transcriptor First-Strand cDNA Synthesis Kit (Roche, Basel, Switzerland). qRT-PCR was performed on a Bio-Rad CFX96 Real-time System with SYBR® Green PCR Supermix (CA, USA). Each reaction



of 20  $\mu\text{L}$  contained 10  $\mu\text{L}$  of SYBR Green premix (2X) (Roche, Basel, Switzerland), 0.5  $\mu\text{L}$  of each primer (10  $\mu\text{M}$ ), 7.0  $\mu\text{L}$  of  $\text{H}_2\text{O}$  and 2  $\mu\text{L}$  of cDNA. Three biological replications with three technical replicates were used for each reaction. The following program for qPCR was used: 98 °C for 30 s, followed by 40 cycles of 98 °C for 10 s and 60 °C for 30 s. *BnActin7* and *Bn26S* gene were used as a control. All primers used in this study are shown in Table S1.

### Metabolite extraction

The metabolite extraction protocol is based on previous studies with a few modifications (Wang *et al.*, 2018; Ying *et al.*, 2020). The cryopreserved biomaterial samples were removed and vacuum freeze-dried. The dried sample was ground in MM 400 (Retsch, Haan, Germany) at 30 Hz for 1.5 min, 100 mg of the powder was weighed, and 1.0 ml 70% methanol containing 0.1 mg/l lidocaine was used as an internal standard for overnight extraction at 4 °C. After extraction, the sample was centrifuged at 10,000 $\times$ g for 10 min, the supernatant was absorbed, and the sample was filtered with a microporous membrane (0.22- $\mu\text{m}$  pore size), and stored in the injection bottle for LC-MS analysis. The quality control (QC) sample was prepared by mixing the sample extracts and used to analyze the reproducibility of the samples under the same treatment method.

### Metabolite detection

The data acquisition instrument system mainly included ultra-high performance liquid chromatography (Ultra Performance Liquid Chromatography, UPLC) (Shim-pack UFLC SHIMADZU CBM20A, <http://www.shimadzu.com.cn/>) and tandem mass spectrometry (Tandem mass spectrometry, MS/MS) (Applied Biosystems 4500 QTRAP, <http://www.appliedbiosystems.com.cn/>). Briefly, 5  $\mu\text{L}$  of each sample was injected onto a Waters ACQUITY UPLC HSS T3 C18 column (2.1 mm  $\times$  100 mm, 1.8  $\mu\text{m}$ ) operating at 40 °C and flow rate of 0.4 mL/min. The mobile phases used were acidified water (0.04% acetic acid) (Phase A) and acidified acetonitrile (0.04 % acetic acid) (Phase B). Compounds were separated using the following gradient: 95:5 Phase A/Phase B at 0 min; 5:95 Phase A/Phase B at 11.0 min; 5:95 Phase A/Phase B at 12.0 min; 95:5 Phase A/Phase B at 12.1 min; 95:5 Phase A/Phase B at 15.0 min (Chen *et al.*, 2013). The effluent was connected to an ESI-triple quadrupole-linear ion trap (QTRAP)-MS.

Mass spectrometry information was measured on API 4500 QTRAP LC/MS/MS system both in positive and negative mode, the main parameters of the linear ion trap and triple quadrupole rod were as follows: electrospray ion source (electrospray ionization, ESI) temperature 550 °C, mass spectrometry voltage 5.5 kV, curtain gas (curtain gas, CUR) 25 psi. The collision-induced ionization (collision-activated dissociation, CD) parameter was set to high. In the qualitative analysis of the metabolite, the isotopic signals, repetitive signals containing  $\text{K}^+$  ions,  $\text{Na}^+$  ions and  $\text{NH}_4^+$  ions were removed. Metabolites were analyzed with reference to internal database, MassBank (<http://www.massbank.jp/>), KNAPSACk (<http://kanaya.naist.jp/KNAPsAcK/>), HMDB (<http://www.hmdb.ca/>) (Wishart *et al.*, 2013), MoTo DB (<https://omictools.com/moto-db-tool>) and METLIN (<http://metlin.scripps.edu/index.php>) (Zhu *et al.*, 2013), as well as other available mass

spectrometry public databases (*Chen et al., 2013*). In the triple quad (QQQ), each ion pair was scanned and detected according to the optimized declustering voltage (declustering potential, DP) and collision energy (collision energy, CE). The  $m/z$  range was set between 50 and 1,000. After mass spectrometry analysis, the original data in wiff format could be opened and browsed using Analyst 1.6.1, which could be used for qualitative and relative quantitative analysis. Cinematographic peak areas were determined and corrected to ensure the accuracy of the metabolite relative quantitative analysis.

### Multivariate statistical analysis

Principal component analysis (PCA) carried out by SPSS for Windows (Version 18.0; SPSS Inc., Chicago, IL, USA), partial least squares discriminant analysis (PLS-DA) and orthogonal partial least-squares discriminant analysis (OPLS-DA) were performed by R software package (<http://bioconductor.org/packages/release/bioc/html/ropls.html>) with data from 12 samples to observe differences in metabolic composition between two *B. napus* ecotypes after cold stress. The loading plot helped to identify the metabolites that contributed the most to the changes in metabolite patterns between the comparison groups. The variables far from the origin in the transverse coordinate direction contributed more to the distinction between the two groups of samples. In addition, the variables in a certain position of the load graph often provided an important contribution to the samples in the same position on the score graph. We combined the multivariate statistical analysis of the VIP value of OPLS-DA and the univariate statistical analysis of the  $T$  test  $P$  value to screen significantly different metabolites among different comparison groups. The thresholds used to determine a significant difference were  $VIP \geq 1$  and  $T$ -test  $P < 0.05$ .

## RESULTS

### Transcriptome analysis for winter and spring ecotypes of *B. napus* in response to cold stress

To detect transcriptome differences in response to cold treatment between different oilseed ecotypes, we performed an RNA-Seq analysis of each five winter and spring ecotypes of *B. napus* using Illumina-based  $2 \times 150$ -bp paired-end read sequencing. In total, 598,775,190 clean reads with 587,373,972 high-quality reads were obtained from 12 libraries (Table 1). In our study, all high-quality reads were mapped to the assembled *B. napus* genome of “Darmor-bzh” (*Chalhoub et al., 2014*) after mapping to Ribosome. Finally, more than 80.83–82.23% of the clean reads in 12 libraries could be mapped on the reference genome (Table S2).

In total, 36,014 (4W), 43,467 (28W), 37,610 (4S) and 43,997 (28S) genes were detected in our study with an FPKM value  $\geq 1$ . Moreover, almost half (44.2–49.9%) of the genes in the four libraries showed moderate expression levels ( $3 < \text{FPKM} < 15$ ), while only a small percentage (~5%) of the genes expressed at high levels ( $\text{FPKM} > 60$ ) (Fig. 1A; Table S3). In addition, more than 66.6% (32,229/48,356) of the genes were detected in all four libraries, with 35,601 genes commonly detected in the 28S and 4S comparison, and 34,158 genes commonly detected in the 28W and 4W comparison, respectively

**Table 1** Basic statistics of RNA-seq reads obtained from 12 libraries.

Sample	Clean data (bp)	Clean reads num	HQ clean data (bp)	HQ clean reads num (%)	Adapter (%)	Low quality (%)	poly A (%)	N (%)
28S1	7,248,282,600	48,321,884	7,036,617,191	47,491,286 (98.28)	301,530 (0.62)	524,164 (1.08)	11 (0)	2,441 (0.01)
28S2	6,382,584,900	42,550,566	6,163,481,119	41,679,946 (97.95)	240,588 (0.57)	629,556 (1.48)	8 (0)	230 (0.00)
28S3	6,636,738,600	44,244,924	6,417,371,237	43,394,204 (98.08)	265,830 (0.60)	584,444 (1.32)	3 (0)	220 (0.00)
4S1	6,222,662,700	41,484,418	6,008,313,279	40,646,634 (97.98)	266,786 (0.64)	570,552 (1.38)	7 (0)	216 (0.00)
4S2	7,840,323,600	52,268,824	7,581,545,439	51,217,238 (97.99)	305,172 (0.58)	730,956 (1.40)	15 (0)	7,714 (0.03)
4S3	7,932,055,800	52,880,372	7,683,042,395	51,881,556 (98.11)	310,230 (0.59)	672,336 (1.27)	14 (0)	8,111 (0.03)
28W1	7,404,752,400	49,365,016	7,180,601,844	48,458,954 (98.16)	283,500 (0.57)	607,390 (1.23)	10 (0)	7,576 (0.03)
28W2	7,129,653,000	47,531,020	6,897,653,432	46,579,764 (98.00)	270,716 (0.57)	666,158 (1.4)	19 (0)	7,172 (0.03)
28W3	8,087,136,900	53,914,246	7,843,866,798	52,953,200 (98.22)	301,946 (0.56)	642,532 (1.19)	13 (0)	8,271 (0.03)
4W1	9,033,641,700	60,224,278	8,752,058,406	59,093,420 (98.12)	386,554 (0.64)	726,226 (1.21)	8 (0)	9,031 (0.03)
4W2	8,177,355,000	54,515,700	7,920,965,399	53,481,182 (98.1)	339,772 (0.62)	678,208 (1.24)	2 (0)	8,267 (0.03)
4W3	7,721,091,300	51,473,942	7,481,296,627	50,496,588 (98.1)	333,700 (0.65)	627,856 (1.22)	7 (0)	7,892 (0.03)

**Note:**


28S1, 28S2 and 28S3 were three replications for 28S; 4S1 4S2 and 4S3 were three replications for 4S; 28W1, 28W2 and 28W3 were three replications for 28W; 4W1, 4W2 and 4W3 were three replications for 4W.

(Figs. 1B–1D; Table S3). Furthermore, high Pearson correlations were detected among biological replicates (Fig. S1).

### Identification of DEGs in response to cold stress

In this study, 25,461 DEGs with 5,853 upregulated and 19,607 downregulated genes between the control and cold-treated spring ecotypes were detected. For winter ecotypes, there were 5,912 upregulated and 22,600 downregulated DEGs (Fig. 2A; Table S4). Interestingly, among the upregulated genes, 3,343 were commonly detected in two ecotypes, and 2,569 and 2,510 genes were specific to winter and spring ecotypes, respectively (Fig. 2B; Table S4). For downregulated genes, 13,703 genes were common in both winter and spring ecotypes, and 8,897 and 5,904 genes were specific to winter and spring ecotypes (Fig. 2C; Table S4). In addition, most of the DEGs (67.4% and 71.2% upregulated genes, and 78.2% and 76.5% downregulated genes in winter and spring ecotypes, respectively) had a  $|\text{fold change}| \leq 8$  and <4% DEGs had a  $|\text{fold change}| \geq 1,024$  (Fig. 2D; Table S4).

As critical regulators, TFs make great contributions in response to various (a) biotic stresses in plants. In this study, 2,346 TFs genes were identified as DEGs in two ecotypes, with 1,855 and 1,585 genes in the winter and spring ecotypes, respectively. Among them, 1,094 genes including 300 upregulated and 794 downregulated genes, were commonly detected in both two ecotypes, of which 761 included 204 upregulated and 557 downregulated genes and 491 included 180 upregulated and 311 downregulated genes in the winter and spring ecotypes, respectively (Fig. 3A). These 2,346 TF genes were unevenly distributed in 53 TF families, of which the top four families, namely *bHLH*, *ERF*, *MYB* and *WRKY*, made up the majority in all groups. Interestingly, most of the *bHLH* family members were downregulated in both two ecotypes after cold stress (Table S5).




**Figure 1** Expression statistics of winter and spring *B. napus* ecotypes in normal and cold stress conditions. (A) Statistical analysis of gene expression identified in the four libraries. (B) Venn diagram of genes detected in the four libraries. (C) Venn diagram of genes detected in the spring *B. napus* ecotype between cold stress and normal conditions. (D) Venn diagram of genes detected in the winter *B. napus* ecotype between cold stress and normal conditions.

Full-size DOI: 10.7717/peerj.8704/fig-1

Plant hormones are not only involved in plant development, but they also have crucial functions in abiotic stresses. In total, 434 genes, including 159 upregulated and 275 downregulated, involved in hormone metabolism or signal transduction process were identified. In addition, 409 genes, including 127 upregulated and 282 downregulated, and 393 genes, including 110 upregulated and 283 downregulated, were screened in winter and spring ecotypes after cold stress, respectively (Fig. 3B). Of them, 268 genes, including 78 up- and 190 downregulated, were commonly detected in both ecotypes after cold stress (Fig. 3B). These hormone genes, ABA, auxin and ET genes made up the majority (Table S6).

In Arabidopsis, 320 genes were identified as key factors involved in low or high temperature stresses. In this study, 568 (234 upregulated and 334 downregulated) genes in total and 284 (138 upregulated and 146 downregulated) genes in common were detected in two ecotypes of *B. napus* after cold stress. In the winter ecotype, 436 genes, including 190 upregulated and 246 downregulated, and 416 genes, including 182 upregulated and 234 downregulated, were screened in winter and spring ecotypes, respectively (Fig. 3C). Among them, key genes encoded calcineurin B-like protein 1 (CBL1), cold regulated 15b




**Figure 2** DEGs identified in the two *B. napus* ecotypes after cold stress. (A) Statistical analysis of DEGs identified in winter and spring *B. napus* ecotypes after cold stress. (B) Venn diagram of upregulated gene between spring and winter *B. napus* ecotypes after cold stress. (C) Venn diagram of downregulated gene between spring and winter *B. napus* ecotypes after cold stress. (D) Fold change statistics of DEGs identified in spring and winter *B. napus* ecotypes after cold stress.

Full-size DOI: 10.7717/peerj.8704/fig-2

(COR15B), cold regulated 15a (COR15A), cold regulated 413 plasma membrane 1 (COR413-PM1), C-repeat/DRE binding factor 1 (CBF1), CBF2, CBF4, EARLY RESPONSE TO DEHYDRATION 14 (ERD14), ERD10, ERD7, LOW-TEMPERATURE-INDUCED 65 (LTI65) and LTI41, were upregulated in both ecotypes of *B. napus* after cold stress (Table S7). Genes encoding beta-6 tubulin (TUB6), cold acclimation protein WCOR413 family, cold shock domain protein 3 (CSP3), COR413-PM2, CPL1, ERD5, ERD6, HIGH EXPRESSION OF OSMOTICALLY RESPONSIVE GENES 1 (HOS1), INDUCER OF CBF EXPRESSION 1 (ICE1), MYB15 and WRKY33 were downregulated in both ecotypes of *B. napus* after cold stress (Table S7).

When plants perceive a low external temperature, they transmit the signal from the cell membrane to the cell, which causes a change in the calcium ion concentration on the cell membrane and triggers a phosphorylation cascade of kinases. The calcium concentration increased to promote the upregulation of ICE1 or ICE1-like, CAMTA1-3 and ZAT12, which directly or indirectly promoted the upregulation of CBFs, followed by adaption to low temperature stress through RAP2.1, 2.6 and ZAT10 (Fig. 4A). In this process, the two ecotypes of rapeseed displayed similar expression patterns, but the degree of change was slightly different (Fig. 4B). Several of them were confirmed by qRT-PCR analysis (Fig. 5).






**Figure 3** Venn diagram of DEGs encoding TFs (A), hormone biosynthesis and signal transductions (B) and temperature responsive proteins (C) between winter and spring *B. napus* ecotypes after cold stress. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674\_img.jpg\) DOI: 10.7717/peerj.8704/fig-3](https://doi.org/10.7717/peerj.8704/fig-3)

High correlation coefficients were detected between RNA-Seq and qRT-PCR technologies, suggesting that the RNA-Seq data were reliable.


### Functional analysis of DEGs involved in the response to cold stress

To classify the functions of the DEGs in the two ecotypes after cold stress, GO was used according to biological process, cellular component and molecular function (Fig. S2).

Among all DEG comparisons, “metabolic process” and “cell process”, “cell part” and “cell”



Full-size [DOI: 10.7717/peerj.8704/fig-4](https://doi.org/10.7717/peerj.8704/fig-4)




Full-size [DOI: 10.7717/peerj.8704/fig-5](https://doi.org/10.7717/peerj.8704/fig-5)

and “catalytic activity” and “binding” were dominant in “biological process”, “cellular component” and “molecular function”, respectively.

In addition, DEGs identified in the two ecotypes after cold stress were used for KEGG pathway analysis to explore their potential functions. In total, 27 and 44 significantly

A Up-regulated	B Down-regulated
Circadian rhythm – plant, Carotenoid biosynthesis, Amino sugar and nucleotide sugar metabolism, Plant-pathogen interaction, Phosphatidylinositol signaling system, Inositol phosphate metabolism, Endocytosis, Flavonoid biosynthesis, Starch and sucrose metabolism, Arginine and proline metabolism, Phenylpropanoid biosynthesis, Glycolysis / Gluconeogenesis, Glycerophospholipid metabolism, Ascorbate and aldarate metabolism, Cysteine and methionine metabolism, Phenylalanine metabolism, Glycerolipid metabolism, Biosynthesis of secondary metabolites, beta-Alanine metabolism, Degradation of aromatic compounds, Flavone and flavonol biosynthesis, Metabolic pathways, Butanoate metabolism, <b>Plant hormone signal transduction</b> , <b>Microbial metabolism in diverse environments</b> , <b>Glutathione metabolism</b> , <b>Stilbenoid, diarylheptanoid and gingerol biosynthesis</b>	Metabolic pathways, Biosynthesis of secondary metabolites, Biosynthesis of antibiotics, N-Glycan biosynthesis, Non-homologous end-joining, Other glycan degradation, Porphyrin and chlorophyll metabolism, Biosynthesis of amino acids, Purine metabolism, Photosynthesis, Pantothenate and CoA biosynthesis, Glycosaminoglycan degradation, Propanoate metabolism, Base excision repair, Arginine biosynthesis, Glycosphingolipid biosynthesis - globo series, Valine, leucine and isoleucine biosynthesis, Glycosphingolipid biosynthesis - ganglio series, Thiamine metabolism, Glycosylphosphatidylinositol(GPI)-anchor biosynthesis, Fatty acid metabolism, Valine, leucine and isoleucine degradation, <b>Microbial metabolism in diverse environments</b> , Steroid biosynthesis, Lysine biosynthesis, Photosynthesis - antenna proteins, Monobactam biosynthesis, Sulfur metabolism, 2-Oxocarboxylic acid metabolism, <b>Brassinosteroid biosynthesis</b> , Fatty acid elongation, <b>Glycerophospholipid metabolism</b> , Phenylalanine, tyrosine and tryptophan biosynthesis, Terpenoid backbone biosynthesis, <b>Plant-pathogen interaction</b> , <b>Folate biosynthesis</b> , Lipoic acid metabolism, <b>RNA degradation</b> , <b>Regulation of autophagy</b> , <b>Glucosinolate biosynthesis</b> , Alanine, aspartate and glutamate metabolism, Fatty acid biosynthesis, <b>Riboflavin metabolism</b> , alpha-Linolenic acid metabolism

**Figure 6** Summary of functional categories that were significantly enriched within up- and downregulated genes in the winter and spring *B. napus* ecotypes. The red font denotes categories that were only detected in the winter *B. napus* ecotype, and the blue font indicates categories were only detected in the spring *B. napus* ecotype. (A) Functional categories of up-regulated DEGs. (B) Functional categories of down-regulated DEGs. [Full-size](#)  DOI: 10.7717/peerj.8704/fig-6

enriched pathways were detected among the up- and downregulated genes, respectively (Fig. 6; Table S8). The upregulated genes that were commonly detected in the two ecotypes were mainly involved in primary metabolism, including amino acid metabolism (arginine, proline, phenylalanine, cysteine and methionine metabolism), carbohydrate metabolism (amino sugar and nucleotide sugar, inositol phosphate, starch and sucrose, glycolysis/ gluconeogenesis, ascorbate and aldarate and butanoate metabolism) lipid metabolism (glycerophospholipid and glycerolipid metabolism) and secondary metabolism (flavonoid biosynthesis, phenylpropanoid biosynthesis, flavone and flavonol biosynthesis and carotenoid biosynthesis), suggesting that this reprogramming of primary and secondary metabolism might represent a response to cold stress (Fig. 6; Table S8). In addition, we observed signal transduction (phosphatidylinositol signaling system), transport and catabolism (endocytosis) and environmental adaptation (circadian rhythm—plant and plant–pathogen interaction). Moreover, plant hormone signal transduction was only detected in the winter *B. napus* ecotype and glutathione metabolism and stilbenoid, diarylheptanoid and gingerol biosynthesis were only detected in the spring ecotype (Fig. 6; Table S8), suggesting that hormone signaling and secondary metabolism were involved in the cold stress response in different *B. napus* ecotypes.

Conversely, among the downregulated genes commonly detected in the two ecotypes were pathways involved in amino acid metabolism (arginine biosynthesis, valine, leucine and isoleucine biosynthesis, lysine biosynthesis and valine, leucine and isoleucine degradation, phenylalanine, tyrosine and tryptophan biosynthesis and alanine, aspartate and glutamate metabolism), carbohydrate metabolism (propanoate metabolism), lipid metabolism (steroid biosynthesis, fatty acid elongation, fatty acid biosynthesis and alpha-linolenic acid

metabolism), nucleotide metabolism (purine metabolism) and glycan biosynthesis and metabolism (Fig. 6; Table S8). In addition, energy metabolism genes (photosynthesis, sulfur metabolism and photosynthesis-antenna proteins) also seemed to be downregulated. Plant–pathogen interaction pathways were only detected in the spring *B. napus* ecotype after cold stress (Fig. 6; Table S8). In the winter *B. napus* ecotype, eight pathways were also detected: glucosinolate biosynthesis, RNA degradation, microbial metabolism in diverse environments, glycerophospholipid metabolism, folate biosynthesis, riboflavin metabolism, brassinosteroid biosynthesis and regulation of autophagy (Fig. 6; Table S8), suggesting that different pathways were involved in different ecotypes after cold stress.


To further explore the molecular mechanisms underlying the response to cold stress in the two *B. napus* ecotypes, the software MapMan was used to classify the DEGs involved in the metabolic pathways (Fig. 7). As shown in Fig. 7, genes involved in photosynthesis and circadian rhythm were mostly suppressed, while cell wall, lipids, and biosynthesis of secondary metabolites were mostly activated. All DEGs involved in different metabolic pathways are summarized in Tables S9 and S10.

### Metabolome analyses

To detect the effects of DEGs in response to cold stress at the metabolite level, metabolite profiling of winter and spring *B. napus* ecotypes using LC–MS was performed in this study. In total, 559 metabolites were distributed in 30 classes (Table S11). Among them, the top three classes, namely organic acids (74, 13.2%), amino acid derivatives (55, 9.8%) and nucleotides and their derivatives (52, 9.3%), made up more than 30% of all the metabolites (Fig. 8A). Among these metabolites, 81 were differentially accumulated metabolites (DAMs) using  $VIP \geq 1$  and  $T$ -test  $P < 0.05$ , with 35 increased and 46 decreased after cold stress. In addition, 47 (including 24 increased and 23 decreased) and 40 (including 15 increased and 25 decreased) metabolites were differentially accumulated in winter and spring *B. napus* ecotypes after cold stress (Figs. 8B–8D; Table S12). Among these differentially accumulated metabolites, lipids and amino acids were the most prominent in winter and spring *B. napus* ecotypes, respectively (Figs. 8E and 8F). Four metabolites increased, namely lysoPC 18:2 (lipids glycerophospholipids, pmb0852), lysoPC 18:3 (lipids glycerophospholipids, pmb0865), D-(+)-sucrose (carbohydrates, pme0519) and dihydromyricetin (flavonol, pme2898), and other two metabolites, guanosine (nucleotide and its derivatives, pme1175) and pyridoxine (vitamins, pme1383), decreased in both winter and spring *B. napus* ecotypes (Figs. 8C and 8D; Table S12). Interestingly, L(-)-malic acid (organic acids, pme2033) was increased in spring oilseed rape ecotype, while it was decreased in the winter oilseed rape ecotype (Figs. 8C and 8D; Table S12), suggesting that the metabolites might have critical roles in response to cold stress.


### Combined analysis of gene expression and metabolite data

To screen key genes and metabolites involved in the response to cold stress, joint data from gene expression and metabolism using pairwise correlations and O2PLS were assessed. A pairwise correlation analysis of all DEGs and DAMs was performed, and the top 50 DEGs and top 50 DAMs are displayed in Fig. 9; Table S13. Two latent variables were




**Figure 7** Overview of DEGs involved in different metabolic processes. The images were drawn using MapMan and show different functional categories of DEGs identified in this study. Each box depicts an individual gene. Red and green represent up-DEGs and down-DEGs, respectively. The scale bar represents fold change values. (A) DEGs identified in the spring *B. napus* ecotype after cold stress. (B) DEGs identified in winter *B. napus* ecotype after cold stress. [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac\_img.jpg\) DOI: 10.7717/peerj.8704/fig-7](https://doi.org/10.7717/peerj.8704/fig-7)





**Figure 8** Metabolites identified in the winter and spring *B. napus* ecotypes under cold and normal conditions. (A) Metabolites identified in this study belonged to different classes. (B) Venn diagram of differentially accumulated metabolites between the winter and spring *B. napus* ecotypes after cold stress. (C) Heatmap of differentially accumulated metabolites in the spring *B. napus* ecotype after cold stress. (D) Heatmap of differentially accumulated metabolites in the winter *B. napus* ecotype after cold stress. (E) Differentially accumulated metabolites in the winter *B. napus* ecotype after cold stress mainly belonged to four classes. (F) Differentially accumulated metabolites in the spring *B. napus* ecotype after cold stress mainly belonged to four classes. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674\_img.jpg\) DOI: 10.7717/peerj.8704/fig-8](https://doi.org/10.7717/peerj.8704/fig-8)



**Figure 9** Loading plots of DEGs and DEMs identified in the spring (A and B) and winter (C and D) *B. napus* ecotypes after cold stress. Red circles represent the top 10 genes and 10 metabolites. (A) Metabolite joint loading plot of spring *B. napus* ecotypes. (B) Transcript joint loading plot of spring *B. napus* ecotypes. (C) Metabolite joint loading plot of winter *B. napus* ecotypes. (D) Transcript joint loading plot of winter *B. napus* ecotypes. [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02\_img.jpg\) DOI: 10.7717/peerj.8704/fig-9](https://doi.org/10.7717/peerj.8704/fig-9)

**Table 2** Top 10 metabolites in winter and spring *B. napus* ecotypes based on O2PLS analysis.

Type	ID	VIP	Metabolite name	Class
Spring <i>B. napus</i>	pme3246	-1.53	Coniferin	Hydroxycinnamoyl derivatives
	pmb3064	-1.26	3- <i>O-p</i> -coumaroyl quinic acid <i>O</i> -hexoside	Quinate and its derivatives
	pmb0859	-0.79	LysoPC 18:1 (2n isomer)	Lipids_Glycerophospholipids
	pme0020	-0.54	L-Phenylalanine	Amino acids
	pmb2444	0.39	MAG (18:3) isomer1	Lipids_Glycerolipids
	pmb0852	0.50	LysoPC 18:2	Lipids_Glycerophospholipids
	pme3212	0.94	Quercetin 3- <i>O</i> -glucoside (Isotrifoliin)	Flavonol
	pme2594	1.00	4-Pyridoxic acid	Pyridine derivatives
	pme3130	1.05	Quercetin 4'- <i>O</i> -glucoside (Spiraeoside)	Flavonol
	pmb2654	1.79	Anthranilate <i>O</i> -hexosyl- <i>O</i> -hexoside	Benzoic acid derivatives
Winter <i>B. napus</i>	pmb2325	-1.38	MAG (18:3) isomer2	Lipids_Glycerolipids
	pme0486	-1.01	Methylmalonic acid	Organic acids
	pme1097	-0.99	Adenine	Nucleotide and its derivatives
	pme0022	-0.89	L-Threonine	Amino acids
	pme2427	-0.62	<i>N</i> -Acetyl-L-tyrosine	Amino acid derivatives
	pme0233	-0.49	Kynurenic acid	Organic acids
	pme0519	0.43	D-(+)-Sucrose	Carbohydrates
	pme1363	0.60	Nicotinic acid adenine dinucleotide	Nucleotide and its derivatives
	pmb0824	0.79	Syringic acid <i>O</i> -feruloyl- <i>O</i> -hexoside	Benzoic acid derivatives
	pme2898	1.68	Dihydromyricetin	Flavonol

identified using O2PLS analysis, with 98.0–98.3% of the total variation in the transcript dataset and 97.4–98.3% of the total variation in the metabolite dataset (Table S14). To visualize the importance of the elements in the two omics, a loading plot of different omics was drawn for all transcriptional and metabolic data (Tables S15 and S16). As shown in Figs. 9A–9C and 9B–9D, the loading coefficient thresholds were higher for the metabolomics dataset than for the transcriptomic dataset, and the top 10 genes and metabolites are represented as a red circular plot (Figs. 9A–9C and 9B–9D). In the winter *B. napus* ecotype, four metabolites (D-(+)-sucrose, nicotinic acid adenine dinucleotide, syringic acid *O*-feruloyl-*O*-hexoside and dihydromyricetin) were increased, and the rest of them (MAG (18:3) isomer2, methylmalonic acid, adenine, L-threonine, *N*-acetyl-L-tyrosine and kynurenic acid) were decreased (Table 2). Interestingly, all ten genes were downregulated, two of which were involved in protein posttranslational modifications, three in the cell, one in amino acid metabolism, one in RNA regulation, one in glycolysis and two encoded unknown proteins (Table 3). In the spring *B. napus* ecotype, four metabolites (Coniferin, 3-*O-p*-coumaroyl quinic acid *O*-hexoside, LysoPC 18:1 (2n isomer) and L-phenylalanine) were decreased and six metabolites (MAG (18:3) isomer1,

**Table 3** Top 10 genes in winter and spring *B. napus* ecotypes based on O2PLS analysis.

Type	Gene ID	Log2 fold change	Bin code	Bin name
Spring <i>B. napus</i>	BnaC04g53300D	-1.01	-	-
	BnaA06g29540D	1.03	27.3.25	RNA.regulation of transcription
	BnaA06g07470D	1.07	27.3.3	RNA.regulation of transcription
	BnaA05g33810D	1.07	8.3	TCA/org transformation
	BnaA05g04890D	1.15	10.5.1.1	cell wall.cell wall proteins
	BnaC09g48200D	1.16	27.3.67	RNA.regulation of transcription
	BnaC05g19620D	1.31	26.2	misc.UDP glucosyl and glucuronyl transferases
	BnaC04g24040D	1.32	-	-
	BnaC08g00430D	1.45	16.8.1.21	secondary metabolism.flavonoids.anthocyanins
	BnaA06g05310D	1.58	15.2	metal handling.binding
Winter <i>B. napus</i>	BnaC09g50620D	-1.03	29.4	protein.postranslational modification
	BnaC01g40370D	-1.00	29.4	protein.postranslational modification
	BnaCnng50490D	-1.42	31.1	cell.organization
	BnaC02g02500D	-1.19	31.4	cell.vesicle transport
	BnaC01g38970D	-1.27	31.4	cell.vesicle transport
	BnaC01g38880D	-1.59	13.1.4.5.1	amino acid metabolism
	BnaC02g01650D	-1.17	27.3.73	RNA.regulation of transcription
	BnaC08g45130D	-1.04	4.1.6	glycolysis.cytosolic branch
	BnaC02g02430D	-1.32	-	-
	XLOC_042135	-1.15	-	-

LysoPC 18:2, quercetin 3-*O*-glucoside (isotrifoliin), 4-pyridoxic acid, quercetin 4'-*O*-glucoside (spiraeoside) and anthranilate *O*-hexosyl-*O*-hexoside) were increased (Table 2). Among the top 10 genes, nine were upregulated and one was downregulated with unknown functions. Nine upregulated genes were found to encode MYB111, RAV1, CARBONIC ANHYDRASE 1, FASCICLIN-LIKE ARABINO GALACTAN PROTEIN 8, NICOTIANAMINE SYNTHASE 3, UDP glucosyl and glucuronyl transferases, anthocyanins, anthocyanin 5-aromatic acyltransferase, eukaryotic aspartyl protease family protein and one unknown function protein (Table 3).

## DISCUSSION

As one major adverse environmental factor, low temperature decreases crop production worldwide and has become a great concern for crop breeders (Khodakovskaya et al., 2006; Chinnusamy, Zhu & Zhu, 2007). Plants have evolved fine regulatory mechanisms in response to low temperature stress, including changes in gene expression and metabolites (Maruyama et al., 2014). In this study, alterations of genes and metabolites after cold stress in both winter and spring *B. napus* ecotypes were detected using RNA-Seq and LC-MS analysis, respectively.

## Roles of lipid metabolism in response to cold stress

The plasma membrane performs crucial roles in response to low temperature stress, and lipids are the major component of the plasma membrane (D'Angeli & Altamura, 2016). Genes involved in lipid metabolism were differentially expressed after cold stress, as shown by the enrichment of fatty acid biosynthesis (ko00061), fatty acid elongation (ko00062), alpha-linolenic acid metabolism (ko00592), steroid biosynthesis (ko00100), glycerolipid metabolism (ko00561) and glycerophospholipid metabolism (ko00564) pathways (Fig. 6; Table S7). Key genes encoding proteins, including sterol methyltransferase 3 (SMT3), sterol 1 (STE1), lipase 1 (LIP1), phospholipase D alpha 1 (PLDALPHA1), diacylglycerol kinase 1 (DGK1), glycerol-3-phosphate acyltransferase 4 (GPAT4) and lysophosphatidyl acyltransferase 2 (LPAT2), were detected as DEGs after cold stress. These genes were involved in the cold response by altering the phospholipid and lipid composition (Li et al., 2004; Testerink & Munnik, 2005; Rodriguez-Vargas et al., 2007). In addition, 19 phospholipid signaling genes were cold-regulated, most of which (15 genes) were upregulated in both winter and spring *B. napus* ecotypes. These genes included three PLD2s, two INOSITOL (1, 4 and 5) P3 5-PHOSPHATASE IIs (IP5P2), two PLC-like phosphodiesterase superfamily protein genes (plcA), four DGK1s, three phospholipase C1s (PLC1), three DGK2s and two inositol monophosphatase family protein genes (SAL3). Taken together, these differentially expressed cold-regulated genes indicated that phospholipid second messengers play critical roles in response to cold signaling. Diacylglycerol (DAG) is generated from phosphatidylinositol 4, 5-bisphosphate by PLCs and converted to phosphatidic acid (PA) by DGKs (Munnik, 2001). PLDs are used to catalyze the transformation of phosphatidylcholine and phosphatidylethanolamine to generate PA. An increasing number of reports suggest that PA plays key roles in cold signaling (Ruelland et al., 2002; Welte et al., 2002; Li et al., 2004). In the metabolism analysis, lysoPC 18:2 (pmb0852) and lysoPC 18:3 (2n isomer, pmb0865) were accumulated in both winter and spring *B. napus* ecotypes. Interestingly, three glycerolipids, namely MAG (18:2, pmb0890), MAG (18:3) isomer1 (pmb2444) and MAG (18:3) isomer4 (pmb1656), were significantly accumulated only in the spring *B. napus* ecotype, while eight glycerophospholipids, including lysoPC 16:0 (pmb0855), lysoPE 16:0 (pmb3117), lysoPE 16:0 (2n isomer, pmd0160), lysoPC 15:0 (pmb2319), lysoPC 18:1 (pmb0882), lysoPC 18:2 (2n isomer, pmb0873), lysoPE 18:1 (pmb3121) and lysoPE 18:2 (2n isomer, pmb0874), were significantly accumulated only in the winter *B. napus* ecotype. However, lysoPC 18:1 (2n isomer, pmb0859) was decreased in the spring *B. napus* ecotype, and MAG (18:3) isomer2 (pmb2325) and 14, 15-dehydrocrepenynic acid (pma0461) were decreased in the winter *B. napus* ecotype after cold stress. These findings suggest that lipid metabolism may have crucial functions in response to cold stress.

## ABA signal transduction components have critical roles in response to cold stress

Increasing evidence has revealed the types of mechanisms involved in cold stress response. As an important stress phytohormone, ABA participates in regulating diverse abiotic stresses. COR genes are precisely regulated by both ABA-dependent and -independent



pathways. Hormone homeostasis may be altered under cold stress compared with normal conditions (Lee, Henderson & Zhu, 2005). ABA levels are increased after cold stress in potato (Chen, Li & Brenner, 1983). Plant freezing tolerance can be enhanced by exogenous application of ABA at a normal temperature, while the functions of ABA in the cold regulation process remain elusive.

To explore the involvement of ABA in the cold stress response in *B. napus*, we detected the expression patterns of ABA biosynthesis and the ABA signaling process in two *B. napus* ecotypes after cold stress. In this study, 27 genes encoding the key rate-limiting enzymes of ABA biosynthesis, including nine-*cis*-epoxycarotenoid dioxygenase 3 (NCED3), NCED4, NCED5, NCED9, ABA DEFICIENT 1 (ABA1), ABA2, abscisic aldehyde oxidase 3 (AAO3), aldehyde oxidase 1 (AAO1), AAO2 and carotenoid cleavage dioxygenase 1 (CCD1), were observed in the two *B. napus* ecotypes after cold stress (Table S5). Among them, 16 genes, 10 down- and 6 upregulated genes were commonly detected in the two *B. napus* ecotypes after cold stress, while five and six ABA biosynthesis genes were specific to the winter and spring *B. napus* ecotype after cold stress (Table S5), respectively. These data are consistent with previous studies conducted in *Arabidopsis* (Lang et al., 1994; Park et al., 2008) and *Populus euphratica* (Chen et al., 2014), indicating that ABA biosynthesis is regulated by cold. In ABA signal transduction, many ABA-induced genes, encoding ABA INSENSITIVE 1 (ABI1), ABI5, ABA-responsive element binding protein 3 (AREB3), abscisic acid responsive element-binding factor 1 (ABF1), ABF2, ABF3, highly ABA-induced PP2C gene 1 (HAI1), HAI2, HAI3, OPEN STOMATA 1 (OST1), PYR1-like 4 (PYL4), PYL6, PYL7, PYL10, regulatory component of ABA receptor 1 (RCAR1), RCAR3, SNF1-related protein kinase 2.2 (SNRK2.2), SNRK2.5, SNRK2.7 and SNRK2.10, were differentially expressed in winter and/or spring *B. napus* ecotype(s) after cold stress (Table S5). ABF1, which binds to the ABA-responsive element in ABA-responsive promoters, was continually induced by cold in *Arabidopsis* (Lee, Henderson & Zhu, 2005). One copy of *BnABF1* and two copies of *BnABF3s* were commonly induced in both winter and spring *B. napus* ecotypes, while two copies of *BnABF2s* were downregulated and only observed in the winter *B. napus* ecotype after cold stress (Table S5). Regardless, our data indicate that ABA functions as a crucial mediator of the cold stress response in *B. napus*.

### Alterations in secondary metabolism in response to cold stress

Genes involved in secondary metabolism, including flavonoid biosynthesis, flavone and flavonol biosynthesis, glucosinolate biosynthesis, monobactam biosynthesis, phenylpropanoid biosynthesis and stilbenoid, diarylheptanoid and gingerol biosynthesis, may play critical roles in response to cold stress (Fig. 6; Table S7). In the flavonoid biosynthesis process, two copies of *flavonol synthase 1s* (*FLS1s*, *BnaA10g22800D* and *BnaC09g47360D*), one *FLS3* (*BnaC02g42580D*), two *TRANSPARENT TESTA 4s* (*TT4*, *BnaC02g05070D* and *BnaC09g43250D*), two *TT5s* (*BnaA09g34840D* and *BnaC08g26020D*) and one *dihydroflavonol 4-reductase* (*DFR*, *BnaC09g17150D*), were upregulated in both winter and spring *B. napus* ecotypes. Interestingly, one *FLS1* (*BnaCnng61390D*), one *DFR* (*BnaA09g15710D*), two *leucoanthocyanidin dioxygenases* (*LDOX*, *BnaC01g14310D* and

*BnaC07g37670D*), two *TT5s* (*BnaA07g37900D* and *BnaC08g26010D*) and three *TT4s* (*BnaA03g04590D*, *BnaA10g19670D* and *BnaC03g06120D*) were only upregulated in the spring *B. napus* ecotype. Unfortunately, no metabolite changes were detected in this process. *BnaA02g03370D* and *BnaC02g06990D* encoding UDP-glucosyl transferase 78D2 (UGT78D2) involved in flavone and flavonol biosynthesis were also commonly upregulated in the two *B. napus* ecotypes. Quercetin 4'-*O*-glucoside (spiraeoside) and quercetin 3-*O*-glucoside (isotrifoliin) belonging to flavonol were only increased in the spring *B. napus* ecotype, while selgin 5-*O*-hexoside (flavone) and hesperetin C-hexosyl-*O*-hexosyl-*O*-hexoside (flavone C-glycosides) were only increased in the winter *B. napus* ecotype after cold stress (Table S10). In total, 17 genes involved in monobactam biosynthesis were downregulated in both winter and spring *B. napus* ecotypes. Genes encoding ATP-SULFURYLASE 3 (APS3), APS4, aspartate kinase 1 (AK-LYS1), dihydrodipicolinate synthase 1 (DHDPS1), DHDPS2 and semialdehyde dehydrogenase family protein were detected and participated in S-assimilation and amino acid metabolism processes. Genes that were upregulated in the spring *B. napus* ecotype after cold stress included seven copies of *cytochrome P450, family 71, subfamily B, polypeptide 17s* (*CYP71B17s*), one *CYP76G1*, two *CYP81F4s*, one *CYP81D8* and three *CYP91A2s*. By contrast, chlorogenic acid (3-*O*-caffeoylquinic acid, pme0398) was decreased in the spring *B. napus* ecotype after cold stress. This metabolite can function in the oxidative stress response. Commonly upregulated in both winter and spring *B. napus* ecotypes after cold stress were lignin biosynthesis genes encoding PHE ammonia lyase 1 (PAL1), cinnamate-4-hydroxylase (C4H), cinnamyl alcohol dehydrogenase 5 (CAD5), *O*-methyltransferase 1 (OMT1), ferulic acid 5-hydroxylase 1 (FAH1), caffeoyl-CoA 3-*O*-methyltransferase (CCOAMT) and hydroxycinnamoyl-CoA shikimate/quinic acid hydroxycinnamoyl transferase (HCT). Coniferin (pme3246), chlorogenic acid (3-*O*-caffeoylquinic acid, pme0398), L-(–)-tyrosine (pme0030) and L-phenylalanine (pme0020) were decreased in the spring *B. napus* ecotype after cold stress.

### Signal transduction in response to cold stress

The signal transduction pathway makes great contributions in response to various environmental stresses, including cold stress (Janska et al., 2010; D'Angeli & Altamura, 2016; Tanou et al., 2017). As an important messenger, Ca<sup>2+</sup> plays a crucial role in stress signaling (Ye et al., 2013). The Ca<sup>2+</sup> level increases rapidly under low temperature stress and activates various signaling pathways in response to cold stress and cold-induced gene regulatory processes (Monroy & Dhindsa, 1995; Dodd, Kudla & Sanders, 2010; Reddy et al., 2011; Shi, Ding & Yang, 2015). Calcium signals are sensed by three major family proteins in higher plants: calmodulins (CaMs), calcium-dependent protein kinases (CDPK) and calcineurin B-like (CBL) proteins (Sheen, 1996; Luan, 2009; Boudsocq & Sheen, 2013; Miura & Furumoto, 2013). The MAPKs cascade and receptor-like protein kinases (RLKs) also contribute in the perception of signals in the external environment and have important roles in environmental stress responses (Morris & Walker, 2003; Lehti-Shiu et al., 2009). The calmodulin-like *OsMSR2* gene is significantly induced by low temperature, and its over-expression in *Arabidopsis* enhances the salt and drought

tolerance ability of the plants (Xu et al., 2011). Five calmodulin genes, two CDPK genes and one CBL gene in *Camellia sinensis* were identified to take part in signal transduction after cold stress using RNA-Seq analysis (Wang et al., 2013b). To determine whether these gene families were involved in cold response in *B. napus*, we explored these genes in the list of DEGs. The results showed that 135 genes, including 44 MAPK cascade genes, 35 CDPKs, 38 CaMs, 16 CIPKs and 2 Ca<sup>2+</sup>-ATPase genes, were differentially expressed, with 68 upregulated and 67 downregulated in winter *B. napus* ecotype after cold stress. In the spring *B. napus* ecotype, 117 genes, including 38 MAPK cascade genes, 44 CDPKs, 19 CaMs, 15 CIPKs and 1 Ca<sup>2+</sup>-ATPase genes, were differentially expressed, with 34 upregulated and 83 downregulated after cold stress (Table S17). Among them, 54 genes (including 25 MAPK cascade genes, 26 CDPKs, 14 CaMs, 13 CIPKs and 1 Ca<sup>2+</sup>-ATPase gene) were commonly detected, with 27 upregulated and 52 downregulated in both winter and spring *B. napus* ecotypes. Interestingly, two genes, encoding CBL1 (*BnaA01g08510D*) and ACA2 (*BnaA08g15730D*), showed opposite expression patterns between the two *B. napus* ecotypes after cold stress. These two genes were downregulated in the winter *B. napus* ecotype but upregulated in the spring *B. napus* ecotype after cold stress (Table S17). Our results provide further evidence supporting the key functions of Ca<sup>2+</sup> binding proteins and the MAPK pathway in response to cold stress in *B. napus*.

### TFs in response to cold stress

Transcription factors play key roles in various biological processes, including diverse stress responses (including cold stress), via binding to cis-acting elements of target genes (Singh, Foley & Onate-Sanchez, 2002; Yamaguchi-Shinozaki & Shinozaki, 2006; Chinnusamy, Zhu & Sunkar, 2010; Lindemose et al., 2013). In this study, various kinds of TF genes were differentially expressed after cold stress (Fig. 3; Table S5), and many of them are involved in the cold stress response in plants (Jakoby et al., 2002; Toledo-Ortiz, Huq & Quail, 2003; Mukhopadhyay, Vij & Tyagi, 2004; Hu et al., 2008; Rushton et al., 2010; Chen et al., 2014; Jiang et al., 2017). Many cold-related (COR) genes contain dehydration-responsive elements/C-repeat elements (DRE/CRT, A/GCCGAC) and myeloblastosis (MYB, C/TAACNA/G) in their promoters and are regulated by AP2/ERF and MYB TFs (Stockinger, Gilmour & Thomashow, 1997; Yamaguchi-Shinozaki & Shinozaki, 2005). CBF TFs, including DREB1A/CBF3, DREB1B/CBF1 and DREB1C/CBF2, belong to the AP2/ERF family, which is divided into the AP2, RAV, ERF and DREB subfamilies (Yamasaki & Randall, 2016; Vazquez-Hernandez et al., 2017). Additionally, AP2/ERF play crucial regulatory functions in response to cold stress, especially DREB proteins, which regulate ABA-independent genes involved in low temperature stress in *A. thaliana* (Du et al., 2016). DREB2B also enhances tolerance to various abiotic stresses (including cold stress) by increasing proline levels in both yeast and tobacco (Li et al., 2014). In our study, 211 *BnAP2/ERF* genes were differentially expressed, and two *BnCBF1s*, five *BnCBF2s* (*BnaC03g71900D* was only upregulated in the spring ecotype) and two *BnCBF4s* (*BnaA10g07630D* was only upregulated in the spring ecotype) were strongly upregulated in both winter and spring *B. napus* ecotypes after cold stress (Table S5). By contrast, no *BnCBF3s* was found to be upregulated in the two *B. napus* ecotypes after cold stress

(Table S5). Two copies of *BnDREB2Bs*, *BnaA05g27930D* were overrepresented in both ecotypes, and *BnaAnng23490D* was only upregulated in the winter *B. napus* ecotype after cold stress (Table S5). *BnERF7s* and *BnERF38s* were overrepresented, while *BnERF1s*, 2s, 5s, 6s and 13s were downregulated after cold stress in the two *B. napus* ecotypes (Table S5).

Several COR genes are not directly regulated by the CBF/DREB family (Fowler & Thomashow, 2002; Kreps et al., 2002). WRKY and NAC TFs also play key roles in response to cold stress (Hu et al., 2008; Yokotani et al., 2013; Yang et al., 2015; Zhuo et al., 2018). An involvement of the WRKY gene family in the cold stress response has been suggested in Arabidopsis (Lee, Henderson & Zhu, 2005), rice (Ramamoorthy et al., 2008), soybean (Zhou et al., 2008), *Vitis vinifera* (Wang et al., 2014), cucumber (Zhang et al., 2016) and cassava (Wei et al., 2016). In our study, 150 (128 in the winter and 115 in the spring ecotype) genes were differentially regulated after cold stress, with 47 up- and 103 downregulated. Among them, 26 up- and 46 downregulated were detected in both winter and spring *B. napus* ecotypes after cold stress. Key members, such as *BnWRKY8s*, 23s, 28s, 31s, 42s, 48s and 123s, were upregulated, while *BnWRKY7s*, 51s, 70s and 74s were downregulated after cold stress in winter and/or spring *B. napus* ecotypes. Key members of NAC genes involved in plant cold responsiveness have been reported, such as *TaNAC2*, *OsNAC5* and *SsNAC23* (Hu et al., 2008; Song et al., 2011). Thirty-two and twenty-eight NAC genes were upregulated after cold stress in the winter and spring *B. napus* ecotypes, respectively, and nine of them were common to the two ecotypes. Many other kinds of TFs, such as bHLH, C2H2, MYB, MYC, MADS and b-ZIP families, were also detected as DEGs after cold stress (Table S5), suggesting that these TFs are also involved in the low temperature regulatory process (Chinnusamy et al., 2003; Yang, Dai & Zhang, 2012; Peng et al., 2013). These up- and downregulated TFs suggest very complex mechanisms of low temperature regulation.

## CONCLUSIONS

In this study, a total of 25,460 DEGs and 41 differentially expressed metabolites (DEMs) in spring oilseed ecotype and 28,512 DEGs and 47 DEMs in winter oilseed ecotype were identified by transcriptome and metabolome analyses after cold stress. In addition, more specific genes and metabolites were detected in winter ecotypes. The combined data indicate that lipid, ABA, secondary metabolism, signal transduction and TFs could be involved in the complex regulation in the spring and winter ecotypes in response to cold stress. This study provides new evidence and insights into how different ecotypes of *B. napus* respond to cold stress from metabolomics and transcriptomics levels.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the Talent Introduction Program of Southwest University (SWU019008), The National Natural Science Foundation of China (31771830, 31701335), National Key Research and Development Program of China (2016YFD0100202), China Postdoctoral Science Foundation (2019M653319), and the “111” Project (B12006).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### **Grant Disclosures**

The following grant information was disclosed by the authors:

Talent Introduction Program of Southwest University: SWU019008.

The National Natural Science Foundation of China: 31771830 and 31701335.

National Key Research and Development Program of China: 2016YFD0100202.

China Postdoctoral Science Foundation: 2019M653319.

“111” Project: B12006.

### **Competing Interests**

Kun Lu is an Academic Editor for PeerJ.

### **Author Contributions**

- Hongju Jian conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Ling Xie performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Yanhua Wang performed the experiments, prepared figures and/or tables, and approved the final draft.
- Yanru Cao performed the experiments, prepared figures and/or tables, and approved the final draft.
- Mengyuan Wan performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Dianqiu Lv conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Jiana Li conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Kun Lu conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Xinfu Xu conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Liezhao Liu conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

### **Data Availability**

The following information was supplied regarding data availability:

Data is available at the NCBI: [PRJNA530291](https://pubmed.ncbi.nlm.nih.gov/32424291/).

### **Supplemental Information**

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8704#supplemental-information>.



## REFERENCES

- Bilska-Kos A, Szczepanik J, Sowinski P. 2016. Cold induced changes in the water balance affect immunocytolocalization pattern of one of the aquaporins in the vascular system in the leaves of maize (*Zea mays* L.). *Journal of Plant Physiology* **205**:75–79 DOI [10.1016/j.jplph.2016.08.006](https://doi.org/10.1016/j.jplph.2016.08.006).
- Boudsocq M, Sheen J. 2013. CDPKs in immune and stress signaling. *Trends in Plant Science* **18**(1):30–40 DOI [10.1016/j.tplants.2012.08.008](https://doi.org/10.1016/j.tplants.2012.08.008).
- Burbulis N, Jonytiene V, Kupriene R, Blinstrubiene A, Liakas V. 2010. Effect of abscisic acid on cold tolerance in *Brassica napus* shoots cultured in vitro. *Journal of Food, Agriculture & Environment* **8**:698–701.
- Bustamante CA, Monti LL, Gabilondo J, Scossa F, Valentini G, Budde CO, Lara MV, Fernie AR, Drincovich MF. 2016. Differential metabolic rearrangements after cold storage are correlated with chilling injury resistance of peach fruits. *Frontiers in Plant Science* **7**(499):1478 DOI [10.3389/fpls.2016.01478](https://doi.org/10.3389/fpls.2016.01478).
- Chalhoub B, Denoeud F, Liu S, Parkin IA, Tang H, Wang X, Chiquet J, Belcram H, Tong C, Samans B, Correa M, Da Silva C, Just J, Falentin C, Koh CS, Le Clainche I, Bernard M, Bento P, Noel B, Labadie K, Alberti A, Charles M, Arnaud D, Guo H, Daviaud C, Alamery S, Jabbari K, Zhao M, Edger PP, Chelaifa H, Tack D, Lassalle G, Mestiri I, Schnell N, Le Paslier MC, Fan G, Renault V, Bayer PE, Golicz AA, Manoli S, Lee TH, Thi VH, Chalabi S, Hu Q, Fan C, Tollenaere R, Lu Y, Battail C, Shen J, Sidebottom CH, Wang X, Canaguier A, Chauveau A, Berard A, Deniot G, Guan M, Liu Z, Sun F, Lim YP, Lyons E, Town CD, Bancroft I, Wang X, Meng J, Ma J, Pires JC, King GJ, Brunel D, Delourme R, Renard M, Aury JM, Adams KL, Batley J, Snowdon RJ, Tost J, Edwards D, Zhou Y, Hua W, Sharpe AG, Paterson AH, Guan C, Wincker P. 2014. Early allopolyploid evolution in the post-Neolithic *Brassica napus* oilseed genome. *Science* **345**(6199):950–953 DOI [10.1126/science.1253435](https://doi.org/10.1126/science.1253435).
- Chen J-Q, Dong Y, Wang Y-J, Liu Q, Zhang J-S, Chen S-Y. 2003. An AP2/EREBP-type transcription-factor gene from rice is cold-inducible and encodes a nuclear-localized protein. *Theoretical and Applied Genetics* **107**(6):972–979 DOI [10.1007/s00122-003-1346-5](https://doi.org/10.1007/s00122-003-1346-5).
- Chen W, Gong L, Guo Z, Wang W, Zhang H, Liu X, Yu S, Xiong L, Luo J. 2013. A novel integrated method for large-scale detection, identification, and quantification of widely targeted metabolites: application in the study of rice metabolomics. *Molecular Plant* **6**(6):1769–1780 DOI [10.1093/mp/sst080](https://doi.org/10.1093/mp/sst080).
- Chen H-H, Li PH, Brenner ML. 1983. Involvement of abscisic acid in potato cold acclimation. *Plant Physiology* **71**(2):362–365 DOI [10.1104/pp.71.2.362](https://doi.org/10.1104/pp.71.2.362).
- Chen J, Tian Q, Pang T, Jiang L, Wu R, Xia X, Yin W. 2014. Deep-sequencing transcriptome analysis of low temperature perception in a desert tree, *Populus euphratica*. *BMC Genomics* **15**(1):326 DOI [10.1186/1471-2164-15-326](https://doi.org/10.1186/1471-2164-15-326).
- Chen L, Zhong H, Ren F, Guo Q-Q, Hu X-P, Li X-B. 2011. A novel cold-regulated gene, *COR25*, of *Brassica napus* is involved in plant response and tolerance to cold stress. *Plant Cell Reports* **30**(4):463–471 DOI [10.1007/s00299-010-0952-3](https://doi.org/10.1007/s00299-010-0952-3).
- Chinnusamy V, Ohta M, Kanrar S, Lee BH, Hong X, Agarwal M, Zhu JK. 2003. *ICE1*: a regulator of cold-induced transcriptome and freezing tolerance in Arabidopsis. *Genes & Development* **17**(8):1043–1054 DOI [10.1101/gad.1077503](https://doi.org/10.1101/gad.1077503).
- Chinnusamy V, Zhu JK, Sunkar R. 2010. Gene regulation during cold stress acclimation in plants. *Methods in Molecular Biology* **639**:39–55 DOI [10.1007/978-1-60761-702-0\\_3](https://doi.org/10.1007/978-1-60761-702-0_3).
- Chinnusamy V, Zhu J, Zhu J-K. 2007. Cold stress regulation of gene expression in plants. *Trends in Plant Science* **12**(10):444–451 DOI [10.1016/j.tplants.2007.07.002](https://doi.org/10.1016/j.tplants.2007.07.002).

- Cook D, Fowler S, Fiehn O, Thomashow MF. 2004.** A prominent role for the CBF cold response pathway in configuring the low-temperature metabolome of Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **101**(42):15243–15248 DOI [10.1073/pnas.0406069101](https://doi.org/10.1073/pnas.0406069101).
- Dodd AN, Kudla J, Sanders D. 2010.** The language of calcium signaling. *Annual Review of Plant Biology* **61**(1):593–620 DOI [10.1146/annurev-arplant-070109-104628](https://doi.org/10.1146/annurev-arplant-070109-104628).
- Du C, Hu K, Xian S, Liu C, Fan J, Tu J, Fu T. 2016.** Dynamic transcriptome analysis reveals AP2/ERF transcription factors responsible for cold stress in rapeseed (*Brassica napus* L.). *Molecular Genetics and Genomics* **291**(3):1053–1067 DOI [10.1007/s00438-015-1161-0](https://doi.org/10.1007/s00438-015-1161-0).
- D'Angeli S, Altamura M. 2016.** Unsaturated lipids change in olive tree drupe and seed during fruit development and in response to cold-stress and acclimation. *International Journal of Molecular Sciences* **17**(11):1889 DOI [10.3390/ijms17111889](https://doi.org/10.3390/ijms17111889).
- Fowler S, Thomashow MF. 2002.** Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell* **14**(8):1675–1690 DOI [10.1105/tpc.003483](https://doi.org/10.1105/tpc.003483).
- Fu Y-P, Liang Y, Dai Y-T, Yang C-T, Duan M-Z, Zhang Z, Hu S-N, Zhang Z-W, Li Y. 2016.** De novo sequencing and transcriptome analysis of *Pleurotus eryngii* subsp. *tuoliensis* (Bailinggu) mycelia in response to cold stimulation. *Molecules* **21**(5):560 DOI [10.3390/molecules21050560](https://doi.org/10.3390/molecules21050560).
- Gong X-X, Yan B-Y, Hu J, Yang C-P, Li Y-J, Liu J-P, Liao W-B. 2018.** Transcriptome profiling of rubber tree (*Hevea brasiliensis*) discovers candidate regulators of the cold stress response. *Genes & Genomics* **40**(11):1181–1197 DOI [10.1007/s13258-018-0681-5](https://doi.org/10.1007/s13258-018-0681-5).
- He YA, Li YP, Cui LX, Xie LX, Zheng CK, Zhou GH, Zhou JJ, Xie XZ. 2016.** Phytochrome B negatively affects cold tolerance by regulating *OsDREB1* gene expression through phytochrome interacting factor-like protein OsPIL16 in rice. *Frontiers in Plant Science* **7**:1963 DOI [10.3389/fpls.2016.01963](https://doi.org/10.3389/fpls.2016.01963).
- Hoermiller II, Naegele T, Augustin H, Stutz S, Weckwerth W, Heyer AG. 2017.** Subcellular reprogramming of metabolism during cold acclimation in *Arabidopsis thaliana*. *Plant, Cell & Environment* **40**(5):602–610 DOI [10.1111/pce.12836](https://doi.org/10.1111/pce.12836).
- Hu H, You J, Fang Y, Zhu X, Qi Z, Xiong L. 2008.** Characterization of transcription factor gene *SNAC2* conferring cold and salt tolerance in rice. *Plant Molecular Biology* **67**(1–2):169–181 DOI [10.1007/s11103-008-9309-5](https://doi.org/10.1007/s11103-008-9309-5).
- Jakoby M, Weisshaar B, Droge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T, Parcy F. 2002.** bZIP transcription factors in Arabidopsis. *Trends in Plant Science* **7**(3):106–111 DOI [10.1016/S1360-1385\(01\)02223-3](https://doi.org/10.1016/S1360-1385(01)02223-3).
- Janska A, Marsik P, Zelenkova S, Ovesna J. 2010.** Cold stress and acclimation—what is important for metabolic adjustment? *Plant Biology* **12**(3):395–405 DOI [10.1111/j.1438-8677.2009.00299.x](https://doi.org/10.1111/j.1438-8677.2009.00299.x).
- Jian H, Ma J, Wei L, Liu P, Zhang A, Yang B, Li J, Xu X, Liu L. 2018.** Integrated mRNA, sRNA, and degradome sequencing reveal oilseed rape complex responses to *Sclerotinia sclerotiorum* (Lib.) infection. *Scientific Reports* **8**(1):10987 DOI [10.1038/s41598-018-29365-y](https://doi.org/10.1038/s41598-018-29365-y).
- Jiang J, Ma S, Ye N, Jiang M, Cao J, Zhang J. 2017.** WRKY transcription factors in plant responses to stresses. *Journal of Integrative Plant Biology* **59**(2):86–101 DOI [10.1111/jipb.12513](https://doi.org/10.1111/jipb.12513).
- Kaplan F, Kopka J, Haskell DW, Zhao W, Schiller KC, Gatzke N, Sung DY, Guy CL. 2004.** Exploring the temperature-stress metabolome of Arabidopsis. *Plant Physiology* **136**(4):4159–4168 DOI [10.1104/pp.104.052142](https://doi.org/10.1104/pp.104.052142).
- Kaplan F, Kopka J, Sung DY, Zhao W, Popp M, Porat R, Guy CL. 2007.** Transcript and metabolite profiling during cold acclimation of Arabidopsis reveals an intricate relationship of

- cold-regulated gene expression with modifications in metabolite content. *Plant Journal* **50**(6):967–981 DOI [10.1111/j.1365-313X.2007.03100.x](https://doi.org/10.1111/j.1365-313X.2007.03100.x).
- Khodakovskaya M, McAvoy R, Peters J, Wu H, Li Y. 2006.** Enhanced cold tolerance in transgenic tobacco expressing a chloroplast  $\omega$ -3 fatty acid desaturase gene under the control of a cold-inducible promoter. *Planta* **223**(5):1090–1100 DOI [10.1007/s00425-005-0161-4](https://doi.org/10.1007/s00425-005-0161-4).
- Kosova K, Prasil IT, Vitamvas P, Dobrev P, Motyka V, Flokova K, Novak O, Tureckova V, Rolcik J, Pesek B, Travnickova A, Gaudinova A, Galiba G, Janda T, Vlasakova E, Prasilova P, Vankova R. 2012.** Complex phytohormone responses during the cold acclimation of two wheat cultivars differing in cold tolerance, winter Samanta and spring Sandra. *Journal of Plant Physiology* **169**(6):567–576 DOI [10.1016/j.jplph.2011.12.013](https://doi.org/10.1016/j.jplph.2011.12.013).
- Kreps JA, Wu Y, Chang H-S, Zhu T, Wang X, Harper JF. 2002.** Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiology* **130**(4):2129–2141 DOI [10.1104/pp.008532](https://doi.org/10.1104/pp.008532).
- Lang V, Mantyla E, Welin B, Sundberg B, Palva ET. 1994.** Alterations in water status, endogenous abscisic acid content, and expression of *rab18* gene during the development of freezing tolerance in *Arabidopsis thaliana*. *Plant Physiology* **104**(4):1341–1349 DOI [10.1104/pp.104.4.1341](https://doi.org/10.1104/pp.104.4.1341).
- Lee B-H, Henderson DA, Zhu J-K. 2005.** The *Arabidopsis* cold-responsive transcriptome and its regulation by ICE1. *Plant Cell* **17**(11):3155–3175 DOI [10.1105/tpc.105.035568](https://doi.org/10.1105/tpc.105.035568).
- Lehti-Shiu MD, Zou C, Hanada K, Shiu S-H. 2009.** Evolutionary history and stress regulation of plant receptor-like kinase/pelle genes. *Plant Physiology* **150**(1):12–26 DOI [10.1104/pp.108.134353](https://doi.org/10.1104/pp.108.134353).
- Li W, Li M, Zhang W, Welti R, Wang X. 2004.** The plasma membrane-bound phospholipase D $\delta$  enhances freezing tolerance in *Arabidopsis thaliana*. *Nature Biotechnology* **22**(4):427–433 DOI [10.1038/nbt949](https://doi.org/10.1038/nbt949).
- Li X, Zhang D, Li H, Wang Y, Zhang Y, Wood AJ. 2014.** *EsDREB2B*, a novel truncated DREB2-type transcription factor in the desert legume *Eremosparton songoricum*, enhances tolerance to multiple abiotic stresses in yeast and transgenic tobacco. *BMC Plant Biology* **14**(1):44 DOI [10.1186/1471-2229-14-44](https://doi.org/10.1186/1471-2229-14-44).
- Lindemose S, O’Shea C, Jensen MK, Skriver K. 2013.** Structure, function and networks of transcription factors involved in abiotic stress responses. *International Journal of Molecular Sciences* **14**(3):5842–5878 DOI [10.3390/ijms14035842](https://doi.org/10.3390/ijms14035842).
- Liu X, Shi W, Yin W, Wang J. 2017.** Distinct cold responsiveness of a *StInvInh2* gene promoter in transgenic potato tubers with contrasting resistance to cold-induced sweetening. *Plant Physiology and Biochemistry* **111**:77–84 DOI [10.1016/j.plaphy.2016.11.021](https://doi.org/10.1016/j.plaphy.2016.11.021).
- Luan S. 2009.** The CBL–CIPK network in plant calcium signaling. *Trends in Plant Science* **14**(1):37–42 DOI [10.1016/j.tplants.2008.10.005](https://doi.org/10.1016/j.tplants.2008.10.005).
- Maruyama K, Urano K, Yoshiwara K, Morishita Y, Sakurai N, Suzuki H, Kojima M, Sakakibara H, Shibata D, Saito K, Shinozaki K, Yamaguchi-Shinozaki K. 2014.** Integrated analysis of the effects of cold and dehydration on rice metabolites, phytohormones, and gene transcripts. *Plant Physiology* **164**(4):1759–1771 DOI [10.1104/pp.113.231720](https://doi.org/10.1104/pp.113.231720).
- Miura K, Furumoto T. 2013.** Cold signaling and cold response in plants. *International Journal of Molecular Sciences* **14**(3):5312–5337 DOI [10.3390/ijms14035312](https://doi.org/10.3390/ijms14035312).
- Monroy AF, Dhindsa RS. 1995.** Low-temperature signal transduction: induction of cold acclimation-specific genes of alfalfa by calcium at 25 °C. *Plant Cell* **7**(3):321–331 DOI [10.2307/3869854](https://doi.org/10.2307/3869854).

- Morris ER, Walker JC. 2003.** Receptor-like protein kinases: the keys to response. *Current Opinion in Plant Biology* **6(4)**:339–342 DOI [10.1016/S1369-5266\(03\)00055-4](https://doi.org/10.1016/S1369-5266(03)00055-4).
- Mukhopadhyay A, Vij S, Tyagi AK. 2004.** Overexpression of a zinc-finger protein gene from rice confers tolerance to cold, dehydration, and salt stress in transgenic tobacco. *Proceedings of the National Academy of Sciences of the United States of America* **101(16)**:6309–6314 DOI [10.1073/pnas.0401572101](https://doi.org/10.1073/pnas.0401572101).
- Munnik T. 2001.** Phosphatidic acid: an emerging plant lipid second messenger. *Trends in Plant Science* **6(5)**:227–233 DOI [10.1016/S1360-1385\(01\)01918-5](https://doi.org/10.1016/S1360-1385(01)01918-5).
- Park H-Y, Seok H-Y, Park B-K, Kim S-H, Goh C-H, Lee B-H, Lee C-H, Moon Y-H. 2008.** Overexpression of Arabidopsis *ZEP* enhances tolerance to osmotic stress. *Biochemical and Biophysical Research Communications* **375(1)**:80–85 DOI [10.1016/j.bbrc.2008.07.128](https://doi.org/10.1016/j.bbrc.2008.07.128).
- Peng H-H, Shan W, Kuang J-F, Lu W-J, Chen J-Y. 2013.** Molecular characterization of cold-responsive basic helix-loop-helix transcription factors MabHLHs that interact with MaICE1 in banana fruit. *Planta* **238(5)**:937–953 DOI [10.1007/s00425-013-1944-7](https://doi.org/10.1007/s00425-013-1944-7).
- Ramamoorthy R, Jiang S-Y, Kumar N, Venkatesh PN, Ramachandran S. 2008.** A comprehensive transcriptional profiling of the WRKY gene family in rice under various abiotic and phytohormone treatments. *Plant and Cell Physiology* **49(6)**:865–879 DOI [10.1093/pcp/pcn061](https://doi.org/10.1093/pcp/pcn061).
- Reddy ASN, Ali GS, Celesnik H, Day IS. 2011.** Coping with stresses: roles of calcium- and calcium/calmodulin-regulated gene expression. *Plant Cell* **23(6)**:2010–2032 DOI [10.1105/tpc.111.084988](https://doi.org/10.1105/tpc.111.084988).
- Ren L, Sun J, Chen S, Gao J, Dong B, Liu Y, Xia X, Wang Y, Liao Y, Teng N, Fang W, Guan Z, Chen F, Jiang J. 2014.** A transcriptomic analysis of *Chrysanthemum nankingense* provides insights into the basis of low temperature tolerance. *BMC Genomics* **15(1)**:844 DOI [10.1186/1471-2164-15-844](https://doi.org/10.1186/1471-2164-15-844).
- Riva-Roveda L, Escale B, Giauffret C, Perilleux C. 2016.** Maize plants can enter a standby mode to cope with chilling stress. *BMC Plant Biology* **16(1)**:212 DOI [10.1186/s12870-016-0909-y](https://doi.org/10.1186/s12870-016-0909-y).
- Rodriguez-Vargas S, Sanchez-Garcia A, Martinez-Rivas JM, Prieto JA, Randez-Gil F. 2007.** Fluidization of membrane lipids enhances the tolerance of *Saccharomyces cerevisiae* to freezing and salt stress. *Applied and Environmental Microbiology* **73(1)**:110–116 DOI [10.1128/AEM.01360-06](https://doi.org/10.1128/AEM.01360-06).
- Ruelland E, Cantrel C, Gawer M, Kader J-C, Zachowski A. 2002.** Activation of phospholipases C and D is an early response to a cold exposure in Arabidopsis suspension cells. *Plant Physiology* **130(2)**:999–1007 DOI [10.1104/pp.006080](https://doi.org/10.1104/pp.006080).
- Rushton PJ, Somssich IE, Ringler P, Shen QJ. 2010.** WRKY transcription factors. *Trends in Plant Science* **15(5)**:247–258 DOI [10.1016/j.tplants.2010.02.006](https://doi.org/10.1016/j.tplants.2010.02.006).
- Sanhueza D, Vizoso P, Balic I, Campos-Vargas R, Meneses C. 2015.** Transcriptomic analysis of fruit stored under cold conditions using controlled atmosphere in *Prunus persica* cv. “Red pearl”. *Frontiers in Plant Science* **6**:788 DOI [10.3389/fpls.2015.00788](https://doi.org/10.3389/fpls.2015.00788).
- Schulz E, Tohge T, Zuther E, Fernie AR, Hinch DK. 2016.** Flavonoids are determinants of freezing tolerance and cold acclimation in *Arabidopsis thaliana*. *Scientific Reports* **6(1)**:34027 DOI [10.1038/srep34027](https://doi.org/10.1038/srep34027).
- Sheen J. 1996.** Ca<sup>2+</sup>-dependent protein kinases and stress signal transduction in plants. *Science* **274(5294)**:1900–1902 DOI [10.1126/science.274.5294.1900](https://doi.org/10.1126/science.274.5294.1900).
- Shi Y, Ding Y, Yang S. 2015.** Cold signal transduction and its interplay with phytohormones during cold acclimation. *Plant and Cell Physiology* **56(1)**:7–15 DOI [10.1093/pcp/pcu115](https://doi.org/10.1093/pcp/pcu115).

- Singh K, Foley RC, Onate-Sanchez L. 2002.** Transcription factors in plant defense and stress responses. *Current Opinion in Plant Biology* 5(5):430–436  
DOI 10.1016/S1369-5266(02)00289-3.
- Song S-Y, Chen Y, Chen J, Dai X-Y, Zhang W-H. 2011.** Physiological mechanisms underlying OsNAC5-dependent tolerance of rice plants to abiotic stress. *Planta* 234(2):331–345  
DOI 10.1007/s00425-011-1403-2.
- Stockinger EJ, Gilmour SJ, Thomashow MF. 1997.** *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a *cis*-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proceedings of the National Academy of Sciences of the United States of America* 94(3):1035–1040 DOI 10.1073/pnas.94.3.1035.
- Stuempfle KJ, Nindl BC, Kamimori GH. 2010.** Stress hormone responses to an ultraendurance race in the cold. *Wilderness & Environmental Medicine* 21(1):22–27  
DOI 10.1016/j.wem.2009.12.020.
- Tanou G, Minas IS, Scossa F, Belghazi M, Xanthopoulou A, Ganopoulos I, Madesis P, Fernie A, Molassiotis A. 2017.** Exploring priming responses involved in peach fruit acclimation to cold stress. *Scientific Reports* 7(1):11358 DOI 10.1038/s41598-017-11933-3.
- Testerink C, Munnik T. 2005.** Phosphatidic acid: a multifunctional stress signaling lipid in plants. *Trends in Plant Science* 10(8):368–375 DOI 10.1016/j.tplants.2005.06.002.
- Tian D-Q, Pan X-Y, Yu Y-M, Wang W-Y, Zhang F, Ge Y-Y, Shen X-L, Shen F-Q, Liu X-J. 2013.** De novo characterization of the *Anthurium* transcriptome and analysis of its digital gene expression under cold stress. *BMC Genomics* 14(1):827 DOI 10.1186/1471-2164-14-827.
- Toledo-Ortiz G, Huq E, Quail PH. 2003.** The *Arabidopsis* basic/helix-loop-helix transcription factor family. *Plant Cell* 15(8):1749–1770 DOI 10.1105/tpc.013839.
- Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, Van Baren MJ, Salzberg SL, Wold BJ, Pachter L. 2010.** Transcript assembly and quantification by RNA-seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nature Biotechnology* 28(5):511–515 DOI 10.1038/nbt.1621.
- Vazquez-Hernandez M, Romero I, Escribano MI, Merodio C, Sanchez-Ballesta MT. 2017.** Deciphering the role of CBF/DREB transcription factors and dehydrins in maintaining the quality of table grapes cv. autumn royal treated with high CO<sub>2</sub> levels and stored at 0 °C. *Frontiers in Plant Science* 8:1591 DOI 10.3389/fpls.2017.01591.
- Wang Y, Zhang X, Yang S, Yuan Y. 2018.** Lignin involvement in programmed changes in peach-fruit texture indicated by metabolite and transcriptome analyses. *Journal of Agricultural and Food Chemistry* 66(48):12627–12640 DOI 10.1021/acs.jafc.8b04284.
- Wang H, Zou Z, Wang S, Gong M. 2013a.** Global analysis of transcriptome responses and gene expression profiles to cold stress of *Jatropha curcas* L. *PLOS ONE* 8(12):e82817  
DOI 10.1371/journal.pone.0082817.
- Wang L, Zhu W, Fang L, Sun X, Su L, Liang Z, Wang N, Londo JP, Li S, Xin H. 2014.** Genome-wide identification of WRKY family genes and their response to cold stress in *Vitis vinifera*. *BMC Plant Biology* 14(1):103 DOI 10.1186/1471-2229-14-103.
- Wang X-C, Zhao Q-Y, Ma C-L, Zhang Z-H, Cao H-L, Kong Y-M, Yue C, Hao X-Y, Chen L, Ma J-Q, Jin J-Q, Li X, Yang Y-J. 2013b.** Global transcriptome profiles of *Camellia sinensis* during cold acclimation. *BMC Genomics* 14(1):415 DOI 10.1186/1471-2164-14-415.
- Wei Y, Shi H, Xia Z, Tie W, Ding Z, Yan Y, Wang W, Hu W, Li K. 2016.** Genome-wide identification and expression analysis of the WRKY gene family in *Cassava*. *Frontiers in Plant Science* 7:25 DOI 10.3389/fpls.2016.00025.



- Welti R, Li W, Li M, Sang Y, Biesiada H, Zhou H-E, Rajashekar CB, Williams TD, Wang X. 2002. Profiling membrane lipids in plant stress responses: role of phospholipase D alpha in freezing-induced lipid changes in Arabidopsis. *Journal of Biological Chemistry* 277(35):31994–32002 DOI 10.1074/jbc.M205375200.
- Wishart DS, Jewison T, Guo AC, Wilson M, Knox C, Liu Y, Djoumbou Y, Mandal R, Aziat F, Dong E, Bouatra S, Sinelnikov I, Arndt D, Xia J, Liu P, Yallou F, Bjorn Dahl T, Perez-Pineiro R, Eisner R, Allen F, Neveu V, Greiner R, Scalbert A. 2013. HMDB 3.0—the human metabolome database in 2013. *Nucleic Acids Research* 41(D1):D801–D807 DOI 10.1093/nar/gks1065.
- Wu ZG, Jiang W, Chen SL, Mantri N, Tao ZM, Jiang CX. 2016. Insights from the cold transcriptome and metabolome of *Dendrobium officinale*: global reprogramming of metabolic and gene regulation networks during cold acclimation. *Frontiers in Plant Science* 7:1653 DOI 10.3389/fpls.2016.01653.
- Xian M, Luo T, Khan MN, Hu L, Xu Z. 2017. Identifying differentially expressed genes associated with tolerance against low temperature stress in *Brassica napus* through transcriptome analysis. *International Journal of Agriculture and Biology* 19(2):273–281 DOI 10.17957/IJAB/15.0274.
- Xiaochuang C, Chu Z, Lianfeng Z, Junhua Z, Hussain S, Lianghuan W, Qianyu J. 2017. Glycine increases cold tolerance in rice via the regulation of N uptake, physiological characteristics, and photosynthesis. *Plant Physiology and Biochemistry* 112:251–260 DOI 10.1016/j.plaphy.2017.01.008.
- Xu G-Y, Rocha PSCF, Wang M-L, Xu M-L, Cui Y-C, Li L-Y, Zhu Y-X, Xia X. 2011. A novel rice calmodulin-like gene, *OsMSR2*, enhances drought and salt tolerance and increases ABA sensitivity in Arabidopsis. *Planta* 234(1):47–59 DOI 10.1007/s00425-011-1386-z.
- Yamaguchi-Shinozaki K, Shinozaki K. 2005. Organization of *cis*-acting regulatory elements in osmotic- and cold-stress-responsive promoters. *Trends in Plant Science* 10(2):88–94 DOI 10.1016/j.tplants.2004.12.012.
- Yamaguchi-Shinozaki K, Shinozaki K. 2006. Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annual Review of Plant Biology* 57(1):781–803 DOI 10.1146/annurev.arplant.57.032905.105444.
- Yamasaki Y, Randall SK. 2016. Functionality of soybean CBF/DREB1 transcription factors. *Plant Science* 246:80–90 DOI 10.1016/j.plantsci.2016.02.007.
- Yang A, Dai X, Zhang W-H. 2012. A R2R3-type MYB gene, *OsMYB2*, is involved in salt, cold, and dehydration tolerance in rice. *Journal of Experimental Botany* 63(7):2541–2556 DOI 10.1093/jxb/err431.
- Yang X, Wang X, Ji L, Yi Z, Fu C, Ran J, Hu R, Zhou G. 2015. Overexpression of a *Miscanthus lutarioriparius* NAC gene *MLNAC5* confers enhanced drought and cold tolerance in Arabidopsis. *Plant Cell Reports* 34(6):943–958 DOI 10.1007/s00299-015-1756-2.
- Ye C-Y, Yang X, Xia X, Yin W. 2013. Comparative analysis of cation/proton antiporter superfamily in plants. *Gene* 521(2):245–251 DOI 10.1016/j.gene.2013.03.104.
- Ying S, Su M, Wu Y, Zhou L, Zhang Y. 2020. Trichome regulator *SLMX1-like* directly manipulates primary metabolism in tomato fruit. *Plant Biotechnology Journal* 18(2):354–363 DOI 10.1111/pbi.13202.
- Yokotani N, Sato Y, Tanabe S, Chujo T, Shimizu T, Okada K, Yamane H, Shimono M, Sugano S, Takatsuji H, Kaku H, Minami E, Nishizawa Y. 2013. *WRKY76* is a rice transcriptional repressor playing opposite roles in blast disease resistance and cold stress tolerance. *Journal of Experimental Botany* 64(16):5085–5097 DOI 10.1093/jxb/ert298.

- Zhang J, Luo W, Zhao Y, Xu Y, Song S, Chong K. 2016.** Comparative metabolomic analysis reveals a reactive oxygen species-dominated dynamic model underlying chilling environment adaptation and tolerance in rice. *New Phytologist* **211**(4):1295–1310 DOI [10.1111/nph.14011](https://doi.org/10.1111/nph.14011).
- Zhao M, Liu W, Xia X, Wang T, Zhang W-H. 2014.** Cold acclimation-induced freezing tolerance of *Medicago truncatula* seedlings is negatively regulated by ethylene. *Physiologia Plantarum* **152**(1):115–129 DOI [10.1111/ppl.12161](https://doi.org/10.1111/ppl.12161).
- Zhao L, Liu F, Xu W, Di C, Zhou S, Xue Y, Yu J, Su Z. 2009.** Increased expression of *OsSPX1* enhances cold/subfreezing tolerance in tobacco and *Arabidopsis thaliana*. *Plant Biotechnology Journal* **7**(6):550–561 DOI [10.1111/j.1467-7652.2009.00423.x](https://doi.org/10.1111/j.1467-7652.2009.00423.x).
- Zhou Q-Y, Tian A-G, Zou H-F, Xie Z-M, Lei G, Huang J, Wang C-M, Wang H-W, Zhang J-S, Chen S-Y. 2008.** Soybean WRKY-type transcription factor genes, *GmWRKY13*, *GmWRKY21*, and *GmWRKY54*, confer differential tolerance to abiotic stresses in transgenic *Arabidopsis* plants. *Plant Biotechnology Journal* **6**(5):486–503 DOI [10.1111/j.1467-7652.2008.00336.x](https://doi.org/10.1111/j.1467-7652.2008.00336.x).
- Zhu Z-J, Schultz AW, Wang J, Johnson CH, Yannone SM, Patti GJ, Siuzdak G. 2013.** Liquid chromatography quadrupole time-of-flight mass spectrometry characterization of metabolites guided by the METLIN database. *Nature Protocols* **8**(3):451–460 DOI [10.1038/nprot.2013.004](https://doi.org/10.1038/nprot.2013.004).
- Zhuo X, Zheng T, Zhang Z, Zhang Y, Jiang L, Ahmad S, Sun L, Wang J, Cheng T, Zhang Q. 2018.** Genome-wide analysis of the NAC transcription factor gene family reveals differential expression patterns and cold-stress responses in the woody plant *Prunus mume*. *Genes* **9**(10):494 DOI [10.3390/genes9100494](https://doi.org/10.3390/genes9100494).