



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Table 2 Laboratory results for patients with EHI

Patient	CK (U/L)	CK-MB ($\mu\text{g/L}$)	Troponin T ($\mu\text{g/L}$)	Myoglobin ($\mu\text{g/L}$)	Aldolase (U/L)	Creatinine ($\mu\text{mol/L}$)
1	3333	105	0.10	192	247	73
2	1395	7.7	0.07	1123	14.1	163
3	913	6.1	0.05	466	12.3	145
4	858	5.3	0.19	921	129	159
5	1115	13.4	0.07	142	23.3	82
6	567	6.9	0.05	203	15.6	104
7	489	8.9	0.07	<21	7.2	83
Laboratory results	Reference range					
CK (U/L)	40-120					
CK-MB ($\mu\text{g/L}$)	0.5-5.0					
Troponin T ($\mu\text{g/L}$)	<0.03					
Myoglobin ($\mu\text{g/L}$)	27.0-55.6					
Aldolase (U/L)	2-12					
Creatinine ($\mu\text{mol/L}$)	63-110					
CK (U/L)	40-120					

References

- [1] Smith JE. Cooling methods used in the treatment of exertional heat illness. *Br J Sports Med* 2005;39:503-7.
- [2] Bouchama A, Knochel JP. Heat stroke. *New Engl J Med* 2002;346:1978-88.
- [3] Armstrong LE, Crago AE, Adams R, et al. Whole-body cooling of hyperthermic runners: comparison of two field therapies. *Am J Emerg Med* 1996;14:355-8.
- [4] Gaffin SL, Gardner JW, Flinn SD. Cooling methods for heatstroke victims. *Ann Intern Med* 2000;132:678.
- [5] Weiner JS, Khogali M. A physiological body-cooling unit for treatment of heat stroke. *Lancet* 1980;1:507-9.
- [6] Armstrong LE, Casa DJ, Millard-Stafford M, Moran DS, et al. American College of Sports Medicine position stand. Exertional heat illness during training and competition. *Med Sci Sports Exerc* 2007;39:556-72.
- [7] SAF Medical Corps Preventive Medicine Directives-PM 20; 2006. Accessed May 14, 2009.
- [8] Khogali M, Weiner JS. Heat stroke: report on 18 cases. *Lancet* 1980;2:276-8.
- [9] Heled Y, Rav-Acha M, Shani Y, et al. The "golden hour" for heatstroke treatment. *Mil Med* 2004;169:184-6.
- [10] Callahan ML. Emergency management of heat illness. *Emergency Physician Series*. North Chicago (Ill): Abbott Laboratories; 1979. p. 1-23.
- [11] Richards D, Richards R, Schofield PJ, et al. Management of heat exhaustion in Sydney's the sun city to surf fun runners. *Med J Aust* 1979;2:457-61.
- [12] Mager M, Hubbard RW, Kerstein MD. Survey and analysis of the medical experience for CAX 8-80. Natick (Mass): Army Research Institute of Environmental Medicine, Technical Report; 1980.
- [13] Maron BJ, Doerer JJ, Haas TS, et al. Sudden deaths in young competitive athletes: analysis of 1866 deaths in the United States, 1980-2006. *Circulation* 2009;119:1085-92 [Epub 2009 Feb 16].
- [14] Pease S, Bouadma L, Kermarrec N, et al. Early organ dysfunction course, cooling time and outcome in classic heatstroke. *Intensive Care Med* 2009;35:1454-8 [Epub 2009 Apr 29].
- [15] Costrini, Pitt HA, Gustafson AB, et al. Cardiovascular and metabolic manifestations of heat stroke and severe heat exhaustion. *Am J Med* 1979;66:296-302.
- [16] Kew M, Bersohn I, Seftel H, et al. Liver damage in heatstroke. *Am Med* 1970;49:192-202.
- [17] Ohman EM, Armstrong PW, Christenson RH, et al. Cardiac troponin T levels for risk stratification in acute ischemia. *N Engl J Med* 1996;335:1333-41.
- [18] Douglas PS, O'Toole ML, Hiller WD, et al. Cardiac fatigue after prolonged exercise. *Circulation* 1987;76:1206-13.

The impact of H1N1 influenza A virus pandemic on the emergency medical service in Kobe

To the Editor,

Pandemic H1N1 2009 influenza virus infection has been identified as the cause of a widespread outbreak of febrile respiratory tract infection. On June 11, 2009, the World Health Organization raised its pandemic alert to the highest level, phase 6.

The H1N1 influenza virus infection rapidly spread all over Japan after the first case was confirmed in Kobe, Japan, on May 16. The outbreak of this virus had a great impact on the Japanese society. There was a shortage in masks and antiseptic [1,2]. More than 4800 schools had been closed, medical services were swamped, many companies were forced into bankruptcy, and the stock market price declined [3,4]. Kobe suffered the most severe impact of the H1N1 influenza in Japan.

Previous studies on severe acute respiratory syndrome [5] and the seasonal influenza [6] reported that emergency departments (EDs) were overcrowded in the epidemic and that the number of ED ambulance diversion increased. Emergency medical service (EMS) was greatly influenced by the epidemic, whereas it remains unknown whether EMS was influenced by the H1N1 influenza virus pandemic.

In Japan, the emergency paramedics are allowed to triage patients. When the emergency paramedics contact with patients, they negotiate with hospitalists to find a hospital where physicians can provide the medical support. To




Fig. 1 The number of febrile patients who called an ambulance remained unchanged during the study period.

investigate the impacts of the H1N1 influenza pandemic on EMS, we studied the situation of EMS during the epidemic provided by the emergency paramedics.

Using an ambulance dispatch data of the Fire Prevention Bureau of Kobe, we extracted data on patients with fever higher than 37.5°C who called an ambulance between April 16 and June 16. We examined the following variables including the time between EMS call and arrival at hospital, that between arrival to the site and the departure from it, and the number of times of negotiation by paramedics with hospitalists for patients' acceptance.

To evaluate the impact of the pandemic on the EMS system, we studied the longitudinal changes in these variables during the pandemic. We defined the period from April 16 to May 16 as the *preepidemic stage* and that from May 17 to June 16th as the *postepidemic stage*. We compared these variables in preepidemic stage with those in postepidemic stage.

Mann-Whitney *U* test was used for statistical analysis, and *P* value of less than .05 was considered statistically significant.

An ambulance was called by 961 patients during the study period, 469 and 492 patients in the preepidemic and postepidemic periods, respectively (Fig. 1). There was no significant difference between the 2 periods (Table 1).

The interval between EMS call and arrival at hospital was significantly longer in the postepidemic period (mean, 28.8 minutes; range, 10.2-97.8 minutes) than in the preepidemic period (mean, 31.3 minutes; range, 9.4-128 minutes) ($P < .001$) (Fig. 2 and Table 1).

The number of hospitals with which the paramedics negotiated for patients' transfer was comparable between the 2 periods (Fig. 3 and Table 1), whereas the sojourn time at the site was prolonged in the postepidemic period (mean, 17.0 minutes; range, 4.0-70 minutes) compared with the preepidemic period (mean, 14.0 minutes; range, 1.0-

Table 1 The interval between EMS call and arrival at hospital and the sojourn time at the site were significantly longer in the postepidemic period than in the preepidemic period

	Preepidemic	Postepidemic	<i>P</i>
	Average (SD)	Average (SD)	
No. of patients who called ambulance (/d)	16.17 (5.015)	15.87 (3.748)	.794 ^a
No. of hospitals with which the paramedics negotiated for patients' transfer (/patient)	1.35 (0.960)	1.34 (0.902)	.778 ^a
	Mean (range)	Mean (range)	<i>P</i>
The interval between EMS call and arrival at hospital (min)	28.8 (10.2-97.8)	31.3 (9.40-128)	<.001 ^b
The sojourn time at the site (min.)	14.0 (4.00-70.0)	17.0 (1.00-86.0)	<.001 ^b

^a *t* test.

^b Mann-Whitney *U* test.




Fig. 2 The interval between EMS call and arrival at hospital was significantly longer in the postepidemic period than in the preepidemic period.

86 minutes). The difference was statistically significant ($P < .001$) (Fig. 4 and Table 1).

This study showed that H1N1 influenza pandemic did not have a great impact on the EMS system of Kobe.

Interestingly, the number of febrile patients who called an ambulance remained unchanged during the study period. This finding was in contrast to the situation in the EDs and the clinics, which were overcrowded with patients who had been scared by the repetitive press reports. Patients with H1N1 influenza occasionally developed severe manifestations. It is reasonable to assume that most patients with suspected H1N1 influenza infection might have visited clinic at its early stages on foot.

It is interesting that paramedics needed an extra time of 3 minutes to triage patients. Considering that the number of hospitals with which the paramedics negotiated for patients' transfer was comparable between the 2 periods, this extra time was required to provide the hospitalists with precise information on febrile patients, including symptoms and records of overseas travel. This finding is consistent with the fact that paramedics were required to fill in a form about patients' information during the study period. It is unlikely that hospitalists refused patients with possible H1N1 influenza infection because of fears of in-hospital spread of the infection.




Fig. 3 The number of hospitals with which the paramedics negotiated for patients' transfer was comparable between the 2 periods.




Fig. 4 The sojourn time at the site was prolonged in the postepidemic period compared with the preepidemic period.

This is a small-sized retrospective study, and unrecognized bias might have existed. Nationwide survey will be necessary to evaluate the true impact of H1N1 influenza pandemic on EMS system. However, it is likely that H1N1 influenza pandemic did not have a great impact on supply and demand of EMS system. H5N1 influenza virus is another concern, although we should be careful in applying the results of this study to predicting its influence on the EMS. There are considerable differences in the virulence between the 2 types of influenza viruses.

Masaharu Tsubokura MD
 Department of Hematology
 Teikyo University Chiba Medical Center
 Ichihara, Japan
 E-mail address: tsubokura-tyk@umin.ac.jp

Haruka Nakada JD
 Tomoko Matsumura MD
 Yuko Kodama RN
 Masahiro Kami MD
 Division of Social Communication System
 for Advanced Clinical Research
 Institute of Medical Science
 University of Tokyo, Tokyo, Japan

doi:10.1016/j.ajem.2009.10.013

References

- [1] Spread of swine flu puts Japan in crisis mode. NY Times 2009.
- [2] Masks, gargles enjoying surge in popularity. Jpn Times Online 2009.
- [3] Flu virus starts to take toll on businesses. Travel industry seen facing biggest risk. Jpn Times Online 2009.
- [4] Flu scare could harm economy. Jpn Times Online 2009.
- [5] Heiber M, Lou WY. Effect of the SARS outbreak on visits to a community hospital emergency department. *Can J Emerg Med* 2006;8(5):323-8.
- [6] Menec VH, Black C, MacWilliam L, Aoki FY. The impact of influenza-associated respiratory illnesses on hospitalizations, physician visits, emergency room visits, and mortality. *Can J Public Health* 2003;94(1):59-63.

In response to

To the Editor,

We thank *** for their insightful comments on our article concerning the diagnosis and management of preexcitation in the emergency department [1]. We would like to clarify the following points:

Atrial fibrillation can indeed present with an apparently regular rhythm and at rates that would seem concordant with simple reentrant tachycardia (approximately 150 beats/min). Yet in the scenario presented by ****, which attempts to differentiate atrial fibrillation with ventricular preexcitation from supraventricular tachycardia with aberrant conduction in the bundle of His, the QRS complex configuration should readily differentiate the 2, at least in the classic case of aberrant conduction with left or right bundle branch blockade, and should not be a major concern (preexcitation should not give a bundle branch block morphology).

The difficulty occurs then when one is not presented with a regular wide-complex tachycardia with a classic QRS complex morphology for bundle branch block. In this case, clinicians must strongly consider ventricular tachycardia (VT) in the differential diagnosis. Fortunately, some guidance exists in this situation. Steurer et al [2] studied this scenario and found that any of the following criteria identified VT (versus ventricular preexcitation) with a 75%