



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Original Article

**Coxsackievirus B3 Infection Triggers Autophagy through 3 Pathways of Endoplasmic Reticulum Stress***

LUO Xiao Nuan^{1,&}, YAO Hai Lan^{2,&}, SONG Juan¹, SONG Qin Qin¹,
SHI Bing Tian¹, XIA Dong¹, and HAN Jun^{1,#}

1. State Key Laboratory of Infectious Disease Prevention and Control, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102206, China; 2. Molecular Immunology Laboratory, Capital Institute of Pediatrics, Beijing 100020, China

Abstract

Objective Autophagy is a highly conserved intracellular degradation pathway. Many picornaviruses induce autophagy to benefit viral replication, but an understanding of how autophagy occurs remains incomplete. In this study, we explored whether coxsackievirus B3 (CVB3) infection induced autophagy through endoplasmic reticulum (ER) stress.

Methods In CVB3-infected HeLa cells, the specific molecules of ER stress and autophagy were detected using Western blotting, reverse transcription polymerase chain reaction (RT-PCR), and confocal microscopy. Then PKR-like ER protein kinase (PERK) inhibitor, inositol-requiring protein-1 (IRE1) inhibitor, or activating transcription factor-6 (ATF6) inhibitor worked on CVB3-infected cells, their effect on autophagy was assessed by Western blotting for detecting microtubule-associated protein light chain 3 (LC3).

Results CVB3 infection induced ER stress, and ER stress sensors PERK/eIF2 α , IRE1/XBP1, and ATF6 were activated. CVB3 infection increased the accumulation of green fluorescent protein (GFP)-LC3 punctuation and induced the conversion from LC3-I to phosphatidylethanolamine-conjugated LC3-1 (LC3-II). CVB3 infection still decreased the expression of mammalian target of rapamycin (mTOR) and p-mTOR. Inhibition of PERK, IRE1, or ATF6 significantly decreased the ratio of LC3-II to LC3-I in CVB3-infected HeLa cells.

Conclusion CVB3 infection induced autophagy through ER stress in HeLa cells, and PERK, IRE1, and ATF6a pathways participated in the regulation of autophagy. Our data suggested that ER stress may inhibit mTOR signaling pathway to induce autophagy during CVB3 infection.

Key words: Coxsackievirus B3 (CVB3); Autophagy; Endoplasmic reticulum (ER) stress; Unfolded protein response (UPR)

Biomed Environ Sci, 2018; 31(12): 867-875

doi: 10.3967/bes2018.115

ISSN: 0895-3988

www.besjournal.com (full text)

CN: 11-2816/Q

Copyright ©2018 by China CDC

*This study was supported by the China Mega-project for Infectious Disease [2018ZX10102001, 2018ZX10711001, 2018ZX10734401, and 2018ZX10734404]; and the SKLID Development Grant [2011SKLID104].

[&]LUO Xiao Nuan and YAO Hai Lan contributed equally to this work and should be considered as co-first authors.

[#]Correspondence should be addressed to HAN Jun, professor, PhD, Tel: 86-10-58900680, E-mail: hanjun_sci@163.com

Biographical notes of first authors: LUO Xiao Nuan, female, born in 1989, PhD, majoring in immunology; YAO Hai Lan, female, born in 1975, PhD, majoring in immunology.

INTRODUCTION

The genus *enterovirus* is a member of the family *picornavirus*, a historically ancient virus species. Enteroviruses includes polioviruses, coxsackieviruses A and B, echoviruses, and recently identified enteroviruses^[1,2]. Most enteroviruses enter the body through the respiratory or gastrointestinal tract and usually cause mild diseases; however, they can cause encephalitis, paralysis, and myocarditis^[3]. Coxsackievirus B3 (CVB3) is one major pathogen of viral myocarditis, a condition that may evolve into dilated cardiomyopathy. The pathogenic mechanisms of CVB3-induced myocarditis include directly attacking the heart tissue and those involving the host immune response to CVB3 infection^[4].

The endoplasmic reticulum (ER) is an essential organelle of eukaryotic cells responsible for protein folding and modification, lipid biosynthesis, and calcium homeostasis^[5]. When circumstances change, such as with glucose deprivation, hypoxia, aberrant Ca²⁺ regulation, and pathogen invasion, the unfolded or misfolded proteins accumulate in the ER, resulting in ER stress^[6]. In response to ER stress, an intracellular adaptive mechanism called the unfolded protein response (UPR) is activated. UPR recovers ER homeostasis by attenuating protein synthesis, upregulating protein folding and trafficking machinery, and activating the ER-associated degradation system. However, when ER stress is so prolonged and severe that ER function cannot be restored, UPR will remove the damaged cells by activating apoptosis^[7]. ER stress and ER stress-induced UPR play important roles in the course of virus infection^[8]. Many plus-strand RNA viruses, including CVB3, Japanese encephalitis virus, dengue virus, and hepatitis C virus, may modify the ER membrane, exploit the ER, or reduce the ER calcium concentration during their infection. Then viral infection leads to imbalance of ER function and, subsequently, activates ER stress^[9-12].

Autophagy is a highly conserved catabolic mechanism that occurs under normal conditions to maintain cellular homeostasis by clearing unwanted cytosolic materials and liberating energy. During the process, a double-membraned structure called an autophagosome is produced that engulfs damaged organelles and long-lived proteins, then fuses with lysosome to degrade and recycle the entrapped contents^[13]. Many studies have focused on the relationship between autophagy and virus infection.

Autophagy is a protective cellular mechanism that defends the host cells against viral infection. Autophagy directly captures and degrades intracellular viral cargo to restrict viral infection. Furthermore, autophagy activates innate and acquired immunities by delivering viral components to endosomal pattern recognition receptors and major histocompatibility complex (MHC)^[14-16]. However, to defend the antiviral role of autophagy, many viruses have evolved various ways to escape, impair or even enhance autophagy. Many RNA viruses infection, including poliovirus, CVB3, foot-and-mouth disease virus, mouse hepatitis virus, and influenza A virus, contribute autophagy and use the membranes of autophagic vacuoles for their RNA replication^[17-19].

Autophagy and ER stress are 2 independent biological processes. However, an increasing number of reports indicate that ER stress may also trigger autophagy. Autophagy alleviates ER stress by removal of the aggregated protein and the portions of the expanded ER^[5,6]. Additionally, several forms of virus-activated ER stress subsequently regulates autophagy^[20-22]. Previous studies have shown that CVB3 infection activates ER stress and autophagy^[9,23], but the relationship of ER stress and autophagy is still not completely understood. In our study, we have demonstrated that CVB3 infection activated all 3 pathways of ER stress contributing to autophagy.

METHODS

Cell and Virus

HeLa cells were purchased from the American Type Culture Collection (ATCC) and grew in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum, penicillin, and streptomycin at 37 °C in 5% CO₂. CVB3 was obtained from ATCC and propagated in HeLa cells. The virus titer was calculated by plaque assay. Briefly, HeLa cells were plated in 6-well plates [(7-9) × 10⁵ cells/well] and grew overnight to approximately 90% confluency, washed with PBS and overlaid with 500 μL of virus-containing samples 10-fold serially diluted in serum-free DMEM. Then the cells were incubated at 37 °C for 1 h and shaken gently every 15 min, washed with PBS and overlaid with 2 mL DMEM supplied with 0.7% agar and 2% FBS. After 72 h, the cells were fixed with 4% paraformaldehyde and stained with 1% crystal violet. Subsequently, the plaques were counted and the viral titer was calculated.

ER Stress Inhibition

HeLa cells grown to approximately 80%-90% confluency were infected with CVB3 at an multiplicity of infection (MOI) of 10 for 1 h, then washed with PBS and cultured in DMEM supplemented with 2% FBS until they were harvested at various times. To inhibit ER stress, HeLa cells were pretreated with 1 μ mol/L PRK-like ER protein kinase (PERK) inhibitor GSK2656157 (S7033, Selleck), 300 μ mol/L ATF6 inhibitor AEBSF (78431, Thermo Scientific) or 5 μ mol/L IRE1 inhibitor STF-083010 (ab146176, Abcam) for 2 h, then the cells were infected with CVB3 at an MOI of 10. At 1 h post-infection, HeLa cells were incubated in fresh maintenance medium in the presence of GSK2656157, AEBSF or STF-083010 until being harvested.

Western Blot Analysis

The cells were lysed in lysis buffer (P0013, Beyotime) on ice for 10 min. The cell lysates were centrifuged at 13,000 \times g for 10 min at 4 °C and the supernatant was preserved. Then the protein sample was separated by SDS-PAGE and transferred to nitrocellulose membranes. The membranes were blocked in PBS containing 5% nonfat dry milk and 0.1% Tween-20 at room temperature for 1 h, and subsequently incubated overnight at 4 °C with one of the following primary antibodies: antibodies against GRP78 (#3177), PERK (#3192), eIF2 α (#5324), p-eIF2 α (#3398), IRE1 (#3294), p62 (#5114), mammalian target of rapamycin (mTOR) (#2983), and p-mTOR (#5536) were purchased from Cell Signaling Technology, antibodies against p-IRE1 (ab124945), ATF6 (ab122897), and β -actin (ab8226) were purchased from Abcam, antibody against LC3 (L8918) was purchased from Sigma, and antibody against VP1 (M7064) was purchased from Dako. After washed with PBS containing 0.1% Tween-20, the membranes were then reacted with Horseradish peroxidase (HRP)-labeled anti-rabbit IgG (ab6721, Abcam) or anti-mouse IgG (ab6728, Abcam) at room temperature for 1 h. Finally, the specific protein bands were visualized using chemiluminescence detection kit (NEL103E001EA, PerkinElmer).

Semiquantitative RT-PCR

Total RNA was extracted using Trizol (15596018, Invitrogen). First-strand complementary DNAs (cDNAs) were synthesized using PrimeScript RT Master Mix (RR036A, TaKaRa). PCR was performed

using Premix Taq™ (R004A, TaKaRa) with the corresponding primer pairs: primers for XBP1 (human) are 5'-CTGGAACAGCAAGTGGTAGA-3' and 5'-CTGGATCCTTCTGGGTAGAC-3'; primers for β -actin (human) are 5'-TTAGTTGCGTTACACCCTTCTTG-3' and 5'-TCACCTTCACCGTCCAGTTT-3'. The condition of PCR was 30 cycles of 98 °C for 10 s, 55 °C for 30 s, and 72 °C for 1 min.

Confocal Microscopy

HeLa cells were seeded in 12-well plates and grown to 60%-70% confluency. Then the cells were transfected with plasmid GFP-LC3 (11060, Addgene) using X-tremeGENE HP DNA Transfection reagent (06366244001, Roche). At 24 h post-transfection, HeLa cells were infected with CVB3 at an MOI of 10 for 8 h, fixed with 4% paraformaldehyde and stained with 4',6-diamidino-2-phenylindole (DAPI). The fluorescence of GFP-LC3 was visualized using a confocal fluorescence microscope.

Statistical Analysis

All experiments were performed at least in 3 independent experiments and the results were presented as means \pm standard deviation (SD). The Student's *t* test was used to analyze data. Statistical significance was set at $P < 0.05$.

RESULTS

CVB3 Infection Activated ER Stress

The upregulation of glucose regulated protein 78 kD (GRP78) is an indicator of ER stress in mammalian cells^[24,25]. To determine whether ER stress was activated by CVB3 infection, we first identified the expression of GRP78 in CVB3-infected cells. HeLa cells were infected with CVB3 at an MOI of 10, and the cells were collected at 6, 8, and 10 h postinfection. The expression levels of GRP78 in CVB3-infected HeLa cells were detected by Western blotting. As shown in Figure 1, compared with control cells, GRP78 expression gradually increased and reached a high level at 10 h postinfection. These results confirmed that CVB3 infection triggered ER stress and activated UPR in HeLa cells.

CVB3 Infection Induced Activation of the ATF6 Pathway

ATF6 is an ER membrane protein with a relative molecular mass of about 90 kD. Under ER stress,

ATF6 is translocated to the Golgi apparatus, where it is cleaved to produce an N-terminal-activated form (p50ATF6). As an active transcription factor, p50ATF6 then enters the nucleus and moderates ER stress-associated gene expression^[26,27]. To investigate whether the ATF6 pathway was activated by CVB3 infection, HeLa cells were collected at 6, 8, and 10 h postinfection. Both full-length ATF6 and cleaved ATF6 were detected by Western blotting. Our results showed that full-length ATF6 (p90ATF6) expression decreased with extension of CVB3 infection. Meanwhile, cleaved ATF6 (p50ATF6) could be clearly detected at each time point in CVB3-infected HeLa cells, and the expression of p50ATF6 significantly increased with extension of infection time (Figure 2). Thus, the results suggested that the ATF6 pathway was activated by CVB3 infection.

CVB3 Infection Induced Activation of the IRE1- XBP1 Pathway

Under ER stress, IRE1 is activated by oligomerization and autophosphorylation and exhibits both endoribonuclease and kinase activity. The IRE1 RNase domain directly splices a 26-nucleotide intron from XBP1 mRNA. Then the spliced

XBP1 (XBP1s) mRNA is translated into a potent transcription factor that increases the expression of ER stress-associated proteins^[28,29]. To explore whether CVB3 infection triggered the activation of IRE1, HeLa cells were inoculated with CVB3 and collected at 6, 8, and 10 h postinfection. The expressions of IRE1 and p-IRE1 were detected by Western blotting. Our results showed that CVB3 increased the phosphorylation level of IRE1. The expression levels of p-IRE1 increased gradually and reached a peak at 10 h postinfection (Figure 3A). To investigate whether CVB3 infection activated the IRE1-XBP1 pathway, both the full-length and spliced XBP1 mRNA were amplified by RT-PCR. The results showed that spliced XBP1 mRNA appeared and increased with the extension of the infection time in CVB3-infected HeLa cells (Figure 3B). Taken together, the results indicated that the IRE1-XBP1 pathway was fully activated in CVB3-infected HeLa cells.

CVB3 Infection Induced PERK- eIF2 α Pathway

The PERK-eIF2 α pathway is an important sensor of ER stress. During ER stress, PERK is activated by dimerization and autophosphorylation. Activated PERK phosphorylates eIF2 α and subsequently effectively blocks protein synthesis^[26]. To understand whether the PERK-eIF2 α pathway was activated by CVB3 infection, the phosphorylation levels of PERK and eIF2 α were detected by Western blot. Western blotting showed that more PERK (phosphorylated PERK) migrated at a higher molecular weight in CVB3-infected cells than in mock-infected cells (Figure 4A), which was consistent with previous researches indicating the phosphorylated PERK shifts




Figure 1. ER stress was induced in CVB3-infected HeLa cells. (A) HeLa cells were infected with CVB3 at MOI of 10. The expression levels of GRP78, VP1 and actin were detected by Western blotting at 6, 8, and 10 h postinfection. 1: HeLa cells, 2: CVB3-infected HeLa cells. (B) Gray scanning analysis of GRP78 to actin. Experiment was repeated three independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Figure 2. ATF6 pathway was activated in CVB3-infected HeLa cells. After HeLa cells were inoculated with CVB3 at MOI of 10, the expression levels of full-length ATF6, cleaved ATF6 and actin were detected by Western blotting at 6, 8, and 10 h postinfection. 1: HeLa cells, 2: CVB3-infected HeLa cells.

slower in polyacrylamide gels than the latent form of the kinase^[30,31]. The result showed that CVB3 infection activated PERK. Consistently, phosphorylated eIF2 α obviously appeared after CVB3 infection and increased in a time-dependent manner (Figure 4A). These results suggested that CVB3 infection activated the PERK-eIF2 α pathway of ER stress.

CVB3 Infection Induced Autophagy

In mammals, LC3 is synthesized as a precursor form and is cleaved at its COOH terminus by the protease ATG4, referred to as LC3-I. LC3-I is conjugated to phosphatidylethanolamine (PE) to form LC3-II, which specifically targets to the autophagosome membrane and participates in the formation of autophagosomes^[13,23]. Therefore the conversion of LC3-I to LC3-II is a key step in autophagy^[23]. To determine whether autophagy was induced by CVB3 infection, we first measured the expression level of LC3-I and LC3-II by Western blot assay at 6, 8, and 10 h postinfection. As shown in Figure 5A and 5B, CVB3 induced the conversion from LC3-I to LC3-II, and the expression of LC3-II reached its highest level at 10 h postinfection, which demonstrated that CVB3 infection induced

autophagy. To further confirm the above results, we examined the distribution of LC3 in CVB3-infected HeLa cells using confocal microscopy. The results showed that CVB3 infection contributed to the obvious accumulation of green fluorescent dots from GFP-LC3 expression (Figure 6), which again showed that CVB3 infection increased formation of autophagosomes.

During autophagy, the autophagy receptor SQSTM1/p62 is delivered to lysosomes for degradation. p62 is used as an indicator of autophagic flux or autophagy-mediated protein degradation^[32]. To determine whether p62 was degraded during CVB3-induced autophagy, HeLa cells were collected at 3 time points postinfection and the expression of p62 was detected using Western blotting. In CVB3-infected cells, p62 expression gradually decreased and reached a minimum level at 10 h postinfection (Figure 5A and 5C). The results demonstrated that CVB3 infection promoted protein degradation by lysosomes.

CVB3 Infection Decreased the Kinase Activity of mTOR

mTOR1 is established as a potent repressor of autophagy in eukaryotes^[33]. The PI3K-AKT signaling axis




Figure 3. IRE1-XBP1 pathway was activated in CVB3-infected HeLa cells. (A) The expression levels of IRE1, p-IRE1, and actin were detected by Western blotting at 6, 8, and 10 h postinfection. (B) The relative XBP1 mRNA splicing was detected at 6, 8, and 10 h postinfection by RT-PCR. HeLa cells were infected with CVB3 at MOI of 10. 1: HeLa cells, 2: CVB3-infected HeLa cells.




Figure 4. CVB3 infection induced PERK-eIF2 α pathway. (A) The expression levels of PERK, p-PERK, eIF2 α , p-eIF2 α , and actin were detected by Western blotting at 6, 8, and 10 h post infection. HeLa cells were infected with CVB3 at MOI of 10. 1: HeLa cells, 2: HeLa cells infected with CVB3. (B) Gray scanning analysis of p-eIF2 α to actin. Experiment was repeated three independent experiments. *** $P < 0.001$.

phosphorylates mTOR1 at Ser2448. Phosphorylated mTOR1 exhibits the kinase activity and inhibits autophagy^[34]. To understand the molecular basis of CVB3-induced autophagy, we detected the expression of mTOR1 and p-mTOR1. Our results showed that CVB3 infection decreased the expression of total mTOR1 and p-mTOR1 (Ser2448) in a time-dependent manner (Figure 7), which indicated that CVB3 infection may inhibit the kinase activity of mTOR1 to induce autophagy.

CVB3 Infection Induced Autophagy through 3 Sensors Pathways of UPR

To explore whether CVB3 infection induced autophagy through UPR, ATF6, IRE1, and PERK signal

pathway was inhibited by the ATF6 inhibitor AEBF^[35,36], the IRE1 inhibitor STF-083010^[37,38], and the PERK inhibitor GSK2656157^[31,39]. Our results showed that these inhibitors obviously decreased the formation of cleaved ATF6 protein, spliced XBP1 mRNA, and phosphorylated PERK protein in CVB3-infected cells (Figure 8A, 8C and 8E). Next, the expression levels of LC3-I and LC3-II were detected using Western blotting. The results showed that the ratio of LC3-II to LC3-I decreased in the presence of the ATF6 inhibitor (Figure 8A and 8B), IRE1 inhibitor (Figure 8C and 8D), or PERK inhibitor (Figure 8E and 8F) in CVB3-infected cells, which demonstrated that CVB3 infection induced autophagy through the ATF6, IRE1, and PERK signal pathways.




Figure 5. CVB3 infection induced autophagy. (A) The expression levels of LC3-I/II, P62, and actin were detected by Western blotting at 6, 8, and 10 h postinfection. HeLa cells were infected with CVB3 at MOI of 10. 1: HeLa cells, 2: CVB3-infected HeLa cells. (B, C) Gray scanning analysis of LC3-II to LC3-I, and P62 to actin. Experiment was repeated three independent experiments. * $P < 0.05$, *** $P < 0.001$.




Figure 6. The fluorescence of GFP-LC3 was observed using confocal microscopy. Robust fluorescence punctate (green dots) was presented in CVB3-infected HeLa cells. Diffused fluorescence (green smears) was expressed in normal cells. Cell nuclei were stained with DAPI ($\times 400$).




Figure 7. CVB3 infection decreased expression of mTOR1 and p-mTOR1. (A) The expression levels of mTOR1, p-mTOR1, and actin were detected by Western blotting at 6, 8, and 10 h postinfection. HeLa cells were infected with CVB3 at MOI of 10. 1: HeLa cells, 2: HeLa cells. (B, C) Gray scanning analysis of p-mTOR1 to actin, and mTOR1 to actin. Experiment was repeated three independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Figure 8. CVB3 infection induced autophagy through ER stress. (A, C, and E) The expression levels of ATF6, p-PERK, LC3-I/II, and actin were detected by Western blotting, and the relative XBP1 splicing was detected by RT-PCR at 6, 8, and 10 h postinfection. (B, D, and F) Gray scanning analysis of LC3-II to LC3-I. Experiment was repeated three independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

DISCUSSION

Ongoing studies demonstrate that ER stress regulates autophagy and ATF6, IRE1, and PERK signal pathways of ER stress participate in the activation of autophagy in various ways. Under ER stress, PERK-eIF2 α upregulates the expression of transcription factor ATF4, which activates another transcription factor, CHOP. ATF4 and CHOP translocate to the nucleus to regulate autophagy-related gene expression^[5]. IRE1 interacts with tumor necrosis factor receptor-associated factor-2 and apoptosis signal-regulating kinase-1 to activate Jun-N-terminal kinase (JNK)^[6]. Then, the activated JNK phosphorylates Bcl-2 to free Bcl-2 from the Beclin-1/Bcl-2 complex, which contributes to the nucleation of autophagy. Meanwhile, IRE1-mediated XBP1 splicing also promotes autophagy through transcriptional activation of autophagy-related genes^[40,41]. Proteolytic ATF6, as an activated transcription factor, participates in regulation of the expression of GRP78 and AKT to induce autophagy^[42,43]. In our study, inhibition of IRE1, PERK, or ATF6 signaling pathways of the UPR sensor decreased the ratio of LC3-II to LC3-I in CVB3-infected HeLa cells, which demonstrated that CVB3 infection triggered autophagy through the IRE1, PERK, and ATF6 signaling pathways.

mTOR is a negative regulator of autophagy. mTOR1 inhibits autophagy through regulation of unc-51-like kinase 1 (ULK1) complex, which composed of ULK1, autophagy-related gene 13 (ATG13), and focal adhesion kinase family-interacting protein of 200 kD (FIP200) in mammals^[44,45]. In our study, CVB3 infection decreased the expression of total mTOR. A possible explanation is that CVB3 2A or 3C protease may cleave mTOR and decrease the expression of mTOR. Because there are many enterovirus protease cleavage sites on mTOR, which were predicted by NetPicoRNA 1.0 Server^[46].

Several kinases, including PERK, protein kinase double-stranded RNA-dependent (PKR), general control non-derepressible-2 (GCN2), and heme-regulated inhibitor (HRI) may phosphorylate eIF2 α to prevent protein synthesis and regulate stress-related gene expression. These kinase are respectively activated by ER stress, virus infection, uncharged tRNAs, and heme depletion^[47]. One report showed that CVB3 infection triggered activation of PKR, not PERK, to phosphorylate eIF2 α ^[9]. However, our results first demonstrated

that both PERK and PKR (data not shown) were activated during CVB3 infection.

In conclusion, this study demonstrated that CVB3 infection induced ER stress to initiate autophagy through 3 classical UPR pathways.

Received: March 8, 2018;

Accepted: November 22, 2018

REFERENCES

- Bauer L, Lyoo H, Van Der Schaar HM, et al. Direct-acting antivirals and host-targeting strategies to combat enterovirus infections. *Curr Opin Virol*, 2017; 24, 1-8.
- Huang YP, Lin TL, Chen YJ, et al. Phylogenetic analysis and development of an immunofluorescence assay for untypeable strains of coxsackievirus B3. *J Microbiol Immunol Infect*, 2014; 47, 447-54.
- Harvala H, Kalimo H, Bergelson J, et al. Tissue tropism of recombinant coxsackieviruses in an adult mouse model. *J Gen Virol*, 2005; 86, 1897-907.
- Garmaroudi FS, Marchant D, Hendry R, et al. Coxsackievirus B3 replication and pathogenesis. *Future Microbiol*, 2015; 10, 629-53.
- Smith M, Wilkinson S. ER homeostasis and autophagy. *Essays Biochem*, 2017; 61, 625-35.
- Hoyer-Hansen M, Jaattela M. Connecting endoplasmic reticulum stress to autophagy by unfolded protein response and calcium. *Cell Death Differ*, 2007; 14, 1576-82.
- Rashid HO, Yadav RK, Kim HR, et al. ER stress: Autophagy induction, inhibition and selection. *Autophagy*, 2015; 11, 1956-77.
- Li S, Kong L, Yu X. The expanding roles of endoplasmic reticulum stress in virus replication and pathogenesis. *Crit Rev Microbiol*, 2015; 41, 150-64.
- Zhang HM, Ye X, Su Y, et al. Coxsackievirus B3 infection activates the unfolded protein response and induces apoptosis through downregulation of p58IPK and activation of CHOP and SREBP1. *J Virol*, 2010; 84, 8446-59.
- Mukherjee S, Singh N, Sengupta N, et al. Japanese encephalitis virus induces human neural stem/progenitor cell death by elevating GRP78, PHB and hnRNP through ER stress. *Cell Death Dis*, 2017; 8, e2556.
- Hou JN, Chen TH, Chiang YH, et al. PERK Signal-Modulated Protein Translation Promotes the Survivability of Dengue 2 Virus-Infected Mosquito Cells and Extends Viral Replication. *Viruses*, 2017; 9, E262.
- Chan SW, Egan PA. Hepatitis C virus envelope proteins regulate CHOP via induction of the unfolded protein response. *Faseb j*, 2005; 19, 1510-2.
- Ravikumar B, Sarkar S, Davies JE, et al. Regulation of mammalian autophagy in physiology and pathophysiology. *Physiol Rev*, 2010; 90, 1383-435.
- Qian M, Fang X, Wang X. Autophagy and inflammation. *Clin Transl Med*, 2017; 6, 24.
- Jordan TX, Randall G. Manipulation or capitulation: virus interactions with autophagy. *Microbes Infect*, 2012; 14, 126-39.
- Orvedahl A, Levine B. Viral evasion of autophagy. *Autophagy*, 2008; 4, 280-5.
- Lai JK, Sam IC, Chan YF. The Autophagic Machinery in Enterovirus Infection. *Viruses*, 2016; 8, E32.

18. Choi Y, Bowman JW, Jung JU. Autophagy during viral infection - a double-edged sword. *Nat Rev Microbiol*, 2018; 16, 341-54.
19. Richards AL, Jackson WT. How positive-strand RNA viruses benefit from autophagosome maturation. *J Virol*, 2013; 87, 9966-72.
20. Sharma M, Bhattacharyya S, Sharma KB, et al. Japanese encephalitis virus activates autophagy through XBP1 and ATF6 ER stress sensors in neuronal cells. *J Gen Virol*, 2017; 98, 1027-39.
21. Lee YR, Kuo SH, Lin CY, et al. Dengue virus-induced ER stress is required for autophagy activation, viral replication, and pathogenesis both *in vitro* and *in vivo*. *Sci Rep*, 2018; 8, 489.
22. Yin H, Zhao L, Jiang X, et al. DEV induce autophagy via the endoplasmic reticulum stress related unfolded protein response. *PLoS One*, 2017; 12, e0189704.
23. Wong J, Zhang J, Si X, et al. Autophagosome supports coxsackievirus B3 replication in host cells. *J Virol*, 2008; 82, 9143-53.
24. Morris JA, Dorner AJ, Edwards CA, et al. Immunoglobulin binding protein (BiP) function is required to protect cells from endoplasmic reticulum stress but is not required for the secretion of selective proteins. *J Biol Chem*, 1997; 272, 4327-34.
25. Zhu G, Lee AS. Role of the unfolded protein response, GRP78 and GRP94 in organ homeostasis. *J Cell Physiol*, 2015; 230, 1413-20.
26. Deegan S, Saveljeva S, Gorman AM, et al. Stress-induced self-cannibalism: on the regulation of autophagy by endoplasmic reticulum stress. *Cell Mol Life Sci*, 2013; 70, 2425-41.
27. Fung TS, Liao Y, Liu DX. Regulation of Stress Responses and Translational Control by Coronavirus. *Viruses*, 2016; 8, 184.
28. Senft D, Ronai ZA. UPR, autophagy, and mitochondria crosstalk underlies the ER stress response. *Trends Biochem Sci*, 2015; 40, 141-8.
29. Jheng JR, Ho JY, Horng JT. ER stress, autophagy, and RNA viruses. *Front Microbiol*, 2014; 5, 388.
30. Jordan R, Wang L, Graczyk TM, et al. Replication of a cytopathic strain of bovine viral diarrhoea virus activates PERK and induces endoplasmic reticulum stress-mediated apoptosis of MDBK cells. *J Virol*, 2002; 76, 9588-99.
31. Krishnamoorthy J, Rajesh K, Mirzajani F, et al. Evidence for eIF2alpha phosphorylation-independent effects of GSK2656157, a novel catalytic inhibitor of PERK with clinical implications. *Cell Cycle*, 2014; 13, 801-6.
32. Liu WJ, Ye L, Huang WF, et al. p62 links the autophagy pathway and the ubiquitin-proteasome system upon ubiquitinated protein degradation. *Cell Mol Biol Lett*, 2016; 21, 29.
33. Kim YC, Guan KL. mTOR: a pharmacologic target for autophagy regulation. *J Clin Invest*, 2015; 125, 25-32.
34. Yao C, Liu BB, Qian XD, et al. Crocin induces autophagic apoptosis in hepatocellular carcinoma by inhibiting Akt/mTOR activity. *Onco Targets Ther*, 2018; 11, 2017-28.
35. Tang Y, Li J, Gao C, et al. Hepatoprotective Effect of Quercetin on Endoplasmic Reticulum Stress and Inflammation after Intense Exercise in Mice through Phosphoinositide 3-Kinase and Nuclear Factor-Kappa B. *Oxid Med Cell Longev*, 2016; 8696587.
36. Okada T, Haze K, Nadanaka S, et al. A serine protease inhibitor prevents endoplasmic reticulum stress-induced cleavage but not transport of the membrane-bound transcription factor ATF6. *J Biol Chem*, 2003; 278, 31024-32.
37. Ming J, Ruan S, Wang M, et al. A novel chemical, STF-083010, reverses tamoxifen-related drug resistance in breast cancer by inhibiting IRE1/XBP1. *Oncotarget*, 2015; 6, 40692-703.
38. Papandreou I, Denko NC, Olson M, et al. Identification of an Irelalpha endonuclease specific inhibitor with cytotoxic activity against human multiple myeloma. *Blood*, 2011; 117, 1311-4.
39. Axten JM, Romeril SP, Shu A, et al. Discovery of GSK2656157: An Optimized PERK Inhibitor Selected for Preclinical Development. *ACS Med Chem Lett*, 2013; 4, 964-8.
40. Kishino A, Hayashi K, Hidai C, et al. XBP1-FoxO1 interaction regulates ER stress-induced autophagy in auditory cells. *Sci Rep*, 2017; 7, 4442.
41. Vidal RL, Hetz C. Unspliced XBP1 controls autophagy through FoxO1. *Cell Res*, 2013; 23, 463-4.
42. Yamazaki H, Hiramatsu N, Hayakawa K, et al. Activation of the Akt-NF-kappaB pathway by subtilase cytotoxin through the ATF6 branch of the unfolded protein response. *J Immunol*, 2009; 183, 1480-7.
43. Gade P, Manjegowda SB, Nallar SC, et al. Regulation of the death-associated protein kinase 1 expression and autophagy via ATF6 requires apoptosis signal-regulating kinase 1. *Mol Cell Biol*, 2014; 34, 4033-48.
44. Ganley IG, Lam Du H, Wang J, et al. ULK1.ATG13.FIP200 complex mediates mTOR signaling and is essential for autophagy. *J Biol Chem*, 2009; 284, 12297-305.
45. Jung CH, Jun CB, Ro SH, et al. ULK-Atg13-FIP200 complexes mediate mTOR signaling to the autophagy machinery. *Mol Biol Cell*, 2009; 20, 1992-2003.
46. Blom N, Hansen J, Blaas D, et al. Cleavage site analysis in picornaviral polyproteins: discovering cellular targets by neural networks. *Protein Sci*, 1996; 5, 2203-16.
47. Donnelly N, Gorman AM, Gupta S, et al. The eIF2alpha kinases: their structures and functions. *Cell Mol Life Sci*, 2013; 70, 3493-511.