



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Published Online
February 24, 2020
[https://doi.org/10.1016/S1473-3099\(20\)30113-4](https://doi.org/10.1016/S1473-3099(20)30113-4)

We declare no competing interests. XP and DC contributed equally to this work. Patient consent was obtained.

*Xingfei Pan, Dexiong Chen, Yong Xia, Xinwei Wu, Tangsheng Li, Xueting Ou, Liyang Zhou, *Jing Liu
panxf0125@163.com or
liujing_Gl@whu.edu.cn

Department of Infectious Diseases (XP, XO, LZ), Emergency Department (DC), Department of Clinical Laboratory (YX), and Department of Radiology (TL), Third Affiliated Hospital of Guangzhou Medical University, Guangzhou, Guangdong 510150, China; Department of Microbiology, Guangzhou Center for Disease Control and Prevention, Guangzhou, Guangdong, China (XW); Department of Gastroenterology, Zhongnan Hospital of Wuhan University, Wuhan, Hubei 430071, China (JL); and Hubei Clinical Center and Key Laboratory of Intestinal and Colorectal Diseases, Wuhan, Hubei, China (JL)

- Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; published online Jan 24. DOI:10.1056/NEJMoa2001017.
- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497–506.
- Chan JF-W, Yuan S, Kok K-H, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020; **395**: 514–23.

Viral load of SARS-CoV-2 in clinical samples

An outbreak caused by a novel human coronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was first detected in Wuhan in December 2019,¹ and has since spread within China and to other countries. Real-time RT-PCR assays are recommended for diagnosis of SARS-CoV-2 infection.² However, viral dynamics in infected patients are still yet to be fully determined. Here, we report our findings from different types of clinical specimens collected from 82 infected individuals.

Serial samples (throat swabs, sputum, urine, and stool) from two patients in Beijing were collected daily after their hospitalisation (patient 1, days 3–12 post-onset; patient 2, days 4–15 post-onset). These samples were examined by an

N-gene-specific quantitative RT-PCR assay, as described elsewhere.³ The viral loads in throat swab and sputum samples peaked at around 5–6 days after symptom onset, ranging from around 10^4 to 10^7 copies per mL during this time (figure A, B). This pattern of changes in viral load is distinct from the one observed in patients with SARS, which normally peaked at around 10 days after onset.⁴ Sputum samples generally showed higher viral loads than throat swab samples. No viral RNA was detected in urine or stool samples from these two patients.

We also studied respiratory samples (nasal [n=1] and throat swabs [n=67], and sputum [n=42]) collected from 80 individuals at different stages of infection. The viral loads ranged from 641 copies per mL to 1.34×10^{11} copies per mL, with a median of 7.99×10^4 in throat samples and 7.52×10^5 in sputum samples (figure C). The only nasal swab tested in this study (taken on day 3 post-onset) showed a viral load of 1.69×10^5 copies per mL. Overall, the viral load early after onset was high ($>1 \times 10^6$ copies per mL). However, a sputum sample collected




Figure: Viral dynamics of SARS-CoV-2 in infected patients

Viral load (mean [SD]) from serial throat swab and sputum samples in patient 1 (A) and patient 2 (B). (C) Viral load (median [IQR]) in throat and sputum samples collected from 80 patients at different stages after disease onset. (D) Correlation between viral load in throat swab samples and viral load in sputum samples.

on day 8 post-onset from a patient who died had a very high viral load (1.34×10^{11} copies per mL). Notably, two individuals, who were under active surveillance because of a history of exposure to SARS-CoV-2-infected patients showed positive results on RT-PCR a day before onset, suggesting that infected individuals can be infectious before them become symptomatic.

Among the 30 pairs of throat swab and sputum samples available, viral loads were significantly correlated between the two sample types for days 1–3 ($R^2=0.50$, $p=0.022$), days 4–7 ($R^2=0.93$, $p<0.001$), and days 7–14 ($R^2=0.95$, $p=0.028$).

From 17 confirmed cases of SARS-CoV-2 infection with available

data (representing days 0–13 after onset), stool samples from nine (53%; days 0–11 after onset) were positive on RT-PCR analysis. Although the viral loads were less than those of respiratory samples (range 550 copies per mL to 1.21×10^5 copies per mL), precautionary measures should be considered when handling faecal samples.

We declare no competing interests. This work was supported by the Theme-based Research Scheme (UGC, Hong Kong).

**Yang Pan, Daitao Zhang, Peng Yang,
*Leo L M Poon, *Quanyi Wang
llmpoon@hku.hk or bjcdcxm@126.com**

Beijing Center for Disease Prevention and Control, Beijing 100013, China (YP, DZ, PY, QW); Beijing Research Center for Preventive Medicine, Beijing, China (YP, DZ, PY, QW); School of Public Health, Capital Medical University, Beijing,

China (YP, PY); and School of Public Health, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region 999077, China (LLMP)

- 1 Lu R, Zhao X, Li J, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020; published online Jan 30. [https://doi.org/10.1016/S0140-6736\(20\)30251-8](https://doi.org/10.1016/S0140-6736(20)30251-8).
- 2 WHO. Laboratory testing for 2019 novel coronavirus (2019-nCoV) in suspected human cases. Interim guidance. Jan 17, 2020. <https://www.who.int/publications-detail/laboratory-testing-for-2019-novel-coronavirus-in-suspected-human-cases-20200117> (accessed Feb 11, 2020).
- 3 Chu DKW, Pan Y, Cheng SMS, et al. Molecular diagnosis of a novel coronavirus (2019-nCoV) causing an outbreak of pneumonia. *Clin Chem* 2020; published online Jan 31. DOI:10.1093/clinchem/hvaa029.
- 4 Peiris JS, Chu CM, Cheng VC, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet* 2003; **361**: 1767–72.