



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Editorial

Middle East Respiratory Syndrome - need for increased vigilance and watchful surveillance for MERS-CoV in sub-Saharan Africa



1. Introduction and background

The past two decades have witnessed the emergence of several new and old respiratory tract infectious diseases, which threaten global health security due to their epidemic potential.^{1,2} These include multi-drug resistant TB, Severe Acute Respiratory Syndrome (SARS), avian and swine influenza and more recently the Middle East Respiratory Syndrome (MERS). MERS is a new zoonotic disease of humans caused by a coronavirus (MERS-CoV) which was first isolated in September, 2012 from a patient who died from a severe respiratory disease in Jeddah Saudi Arabia.³ Since then MERS has attracted global media attention because it is associated with a high mortality (44%) in individuals who have co-morbidities such as diabetes, chronic renal, liver or lung illnesses or in those who are immunocompromised.^{4,5}

The recent unprecedented outbreak of the MERS^{6,7} in South Korea which arose consequential to the importation of MERS-CoV by a South Korean traveler to the Middle East, alarmed global public health authorities and highlights the potential of MERS-CoV to spread across the globe and cause local outbreaks. The WHO Director general convened the ninth meeting of the Emergency Committee (EC) under the International Health Regulations regarding MERS-CoV on 16 June 2015 to discuss the Korean outbreak.⁸ As of 23rd June the total number of MERS cases reported from the Republic of South Korea now stands at 175 (94 currently receiving treatment, 54 recovered, 27 deaths).^{6,8} Of 175 cases, 80 patients and 33 hospital staff had contracted the virus nosocomially, 62 friends, colleagues and relatives had come in contact within healthcare facilities while visiting family members with MERS.

2. Global distribution of MERS cases

Virological and serological studies from several Middle Eastern, West and East African countries indicate that bats and dromedary camels are likely reservoirs of MERS-CoV.^{9–12} However, human MERS-CoV infections appear to be endemic only to countries in the Middle East where sporadic cases continue to occur in the community throughout the year.¹³ From currently available data it appears that MERS-CoV does not transmit easily from person to person and to date no sustained community transmission has been documented. As of 22nd June 2015, 1038 cases of MERS-CoV infection with 459 deaths (44% mortality) have been notified to the World Health Organization,¹³ a large majority of MERS cases have been reported from Saudi Arabia and the United Arab Emirates. MERS cases have also been detected in Algeria, Austria, China, Egypt, France, Germany, Greece, Iran, Italy, Jordan, Kuwait, Malaysia,

Netherlands, Philippines, Lebanon, Oman, Qatar, Tunisia, Turkey, Yemen, United Kingdom, and United States of America. All MERS cases reported from the USA, European and Asian countries had a history of travel to the Middle East. MERS cases continue to be reported from the Middle East with on going MERS outbreak in Hufuf, Saudi Arabia.¹³ The outbreak in Seoul, Republic of Korea, has been linked to a single individual who had travelled to Saudi Arabia. The first MERS case in Thailand was reported last week and the patient had a history of travel from the Sultanate of Oman.¹³

3. MERS and sub-Saharan Africa

Of note is the striking absence of any MERS cases (primary or travel related) reported from sub-Saharan African (SSA) countries.^{14,15} The reasons MERS-CoV predominantly affects humans in the Middle East and is not endemic in Africa where MERS-CoV-infected camels and bats are present requires further study. A likely explanation may be that this may simply reflect the lack of clinical awareness of exposure risk, diagnosis and treatment of respiratory tract infections largely remains clinically based and empiric in most SSA countries coupled with absence of surveillance.¹⁴

Every year an estimated 10 million pilgrims from over 182 countries travel to the Kingdom of Saudi Arabia to participate in Hajj pilgrimage, the mini-pilgrimage Umrah (which is performed at any time of the year), or for the month of Ramadan.¹⁶ Of these, an estimated 1 million pilgrims come from sub-Saharan African countries. There were no cases of MERS reported during the 2012, 2013 and 2014 Hajj pilgrimages or the Ramadan period.^{17–19} However, the risk of MERS-CoV spreading globally remains due to the continuous influx of pilgrims and the persistent low levels of endemic MERS-CoV transmission to humans in Saudi Arabia. There is also the possibility that MERS-CoV may mutate into a form more adaptable for human to human transmission over time.

The potential risk of MERS-CoV infection to pilgrims who visit Saudi Arabia from different regions of the world was estimated by Coker and colleagues²⁰ based on overall incidence of MERS cases in Saudi Arabia since its first discovery in 2012. Their estimates based on the most likely scenario using recent pilgrim numbers for sub-Saharan Africa are that there will be at most ten returning pilgrims each year with MERS-CoV infections. National surveillance systems should be on alert for the low but long-lasting risk of MERS-CoV infected pilgrims returning from the Umrah throughout the year, and also for the large numbers of refugees at several conflict zones in the Middle East (those migrating from Syria to Turkey and from the Yemen border into Saudi Arabia and beyond).

4. Lessons from the Korea MERS outbreak

The recent MERS outbreak in the Republic of Korea was associated with secondary, tertiary, quarternary and quinary cases of MERS-CoV transmission, though fortunately there has been no sustained community transmission.^{6,8} The Republic of Korea MERS-CoV outbreak has many similarities with that of previously reported MERS-CoV outbreaks which occurred at healthcare facilities in several cities in Saudi Arabia and from Jordan^{21–24} which were all associated with breaches and gaps in infection prevention and control protocols.

These lapses in Korean hospitals enabled MERS-CoV infected and uninfected patients, staff and visitors to mix freely in busy and crowded accident and emergency departments, within wards and multi-bed hospital rooms, with no isolation or quarantine of suspected cases. Public health measures such as enhanced contact tracing and isolation and quarantine put in place by the Korean government to control the outbreak eventually led to the decline in the numbers of MERS cases and the outbreak is being brought under control. The importance of infection controls measures was also illustrated by the Saudi Arabian hospital MERS outbreaks, where well-trained health care and workforce brought the hospital outbreaks under control quickly.^{21,22}

The WHO EC meeting noted that⁸ "... the outbreak was strongly associated with healthcare settings and that the main factors contributing to the spread of MERS-CoV in the Republic of Korea were:

- a) Lack of awareness among health care workers and the general public about MERS;
- b) Suboptimal infection prevention and control measures in hospitals;
- c) Close and prolonged contact of infected MERS patients in crowded emergency rooms and multibed rooms in hospitals;
- d) The practice of seeking care at multiple hospitals ("doctor shopping");
- e) The custom of many visitors or family members staying with infected patients in the hospital rooms facilitating secondary spread of infections among contacts."

The WHO EC⁸ referred to the outbreak as a 'wake-up' call and state that in a highly mobile world, all countries should always be prepared for the unanticipated possibility of outbreaks of MERS-CoV and other serious infectious diseases.

The Korean MERS outbreak is the largest recorded from outside the Middle East and the largest imported from a returning traveller to the Middle East, raising several important issues for global surveillance and control. It illustrates that MERS-CoV, three years after its first discovery remains an important global public health risk with many unanswered questions.²⁵ Further international spread should be anticipated and countries with weaker health systems and lack of laboratory facilities to accurately screen for MERS-CoV need to be vigilant. This will pose major challenges.²⁵ There are important lessons here for sub-Saharan African and other developing countries from where MERS-CoV cases have not yet been detected.

5. MERS-CoV surveillance in sub-Saharan Africa

As the recent Ebola Virus Disease epidemic illustrates, African countries may be very vulnerable to a Korea-like MERS-CoV outbreak, which may arise from returning pilgrims or other travellers from Saudi Arabia²⁶ or from traders between Saudi Arabia and the Horn of Africa. MERS-CoV is transmitted through MERS-CoV-infected respiratory secretions for which contact and droplet precautions are recommended.^{27–30}

The Korean MERS outbreak highlights that hospitals provide ideal conditions for amplifying MERS-CoV transmission arising from close contact between patients, healthcare and ancillary staff, relatives and other visitors, which enables spread of MERS-CoV.^{6,8} It is critical that every country should maintain a high level of vigilance and perform MERS-CoV surveillance according to widely available expert recommendations,^{27–30} whether or not MERS cases have been detected in their countries, it ensuring infection prevention and control protocols are in place at all health-care facilities. Those who travel must be educated to follow basic hygiene measures³⁰ and those develop ill health during their trip to the Middle East, or soon after their return should seek medical care and volunteer the history of travel to their healthcare provider.

Sub-Saharan African governments must pay serious attention to strengthening infection control and public health surveillance systems. All healthcare workers and travellers from Africa to the Middle East should be aware of the threat to global health security posed by MERS-CoV. Considering a diagnosis of MERS at first presentation may be difficult due to non-specific symptoms at clinical presentation. However it is important that prevention and control measures are instituted at first consideration of MERS as a diagnosis to prevent spread of MERS-CoV. Hospitals and clinics providing care for patients infected with suspected or confirmed MERS-CoV infection should take appropriate measures to decrease the risk of MERS-CoV transmission from the infected patient to other patients, doctors, nurses, allied health-care workers, relatives and visitors. Health-care workers should be educated and trained in infection prevention and control and should have continuing professional development on these issues.

Over the past decade, several surveillance systems have been introduced to monitor the emergence of new infectious pathogens.³¹ As the Ebola virus epidemic in West Africa showed, surveillance systems in African countries for infectious diseases with epidemic potential require strengthening. More effective national, regional, and international surveillance systems are required to enable rapid identification of emerging respiratory epidemics, diseases with epidemic potential, their specific microbial cause, origin, mode of acquisition, and transmission dynamics.

In light of the Republic of Korea MERS outbreak increased vigilance and surveillance for MERS-CoV should be carried out by health services in African countries using current clinical and public health guidelines for MERS-CoV.²⁸ Although resources may not allow for making an accurate diagnosis of MERS, a high degree of awareness of the possibility of MERS-CoV infection in all returning pilgrims will allow early, isolation of patients and putting in place infection control measures, avoiding a repeat of the Korea outbreak.

Sub-Saharan African countries need to protect themselves against the possible outbreaks akin to the Korean one. MERS-CoV should be included in list of pathogens by The African Network of Influenza Surveillance and Epidemiology (ANISE)³² and MERS-CoV should be made part of the Strengthening Influenza Sentinel Surveillance in Africa (SISA)³³ with national, regional and international reporting mechanisms³³ in liaison with other stakeholders involved in global infectious diseases surveillance. New, low cost, rapid, sensitive and specific diagnostic tests that can be used at all points of healthcare are required for all infectious diseases which threaten global health security.³⁴ The exact mode of transmission and pathogenesis of MERS-CoV and other novel respiratory tract viruses such as H7N9 influenza A virus require definition so that more effective prevention and management measures can be developed and introduced.³⁵ A united and coordinated global response is needed to tackle emerging respiratory tract infections and to fill major gaps²⁵ in the understanding of the epidemiology, transmission dynamics, pathogenesis prevention and control of these infectious diseases.

Declaration: All authors declare no conflicts of interest.

References

- Gautret P, Gray GC, Charrel RN, Odezulu NG, Al-Tawfiq JA, Zumla A, et al. Emerging viral respiratory tract infections—environmental risk factors and transmission. *Lancet Infect Dis* 2014 Nov; **14**(11):1113–22.
- McCloskey B, Dar O, Zumla A, Heymann DL. Emerging infectious diseases and pandemic potential: status quo and reducing risk of global spread. *Lancet Infect Dis* 2014 Oct; **14**(10):1001–10.
- Zaki AM, van Boheemen S, Bestebroer TM, et al. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012; **367**:1814–20.
- Zumla A, Hui DS, Perlman S. Middle East respiratory syndrome. *Lancet* 2015 Jun 3. pii: S0140-6736(15)60454-8.
- Hui DS, Memish ZA, Zumla A. Severe acute respiratory syndrome vs. the Middle East respiratory syndrome. *Curr Opin Pulm Med* 2014 May; **20**(3):233–41.
- Source: Korean MOH press release 23 Jun 2015 [trans. Korean subscribers, edited http://www.mw.go.kr/front_new/al/sal0301vw.jsp?PAR_MENU_ID=04&MENU_ID=0403&page=1&CONT_SEQ=323670].
- Hui DS, Perlman S, Zumla A. Spread of MERS to South Korea and China. *Lancet Respir Med* 2015 Jun 4. pii: S2213-2600(15)00238-6. doi: 10.1016/S2213-2600(15)00238-6.
- WHO -IHR Emergency Committee concerning Middle East respiratory syndrome coronavirus <http://www.who.int/mediacentre/news/statements/2015/ihr-ec-mers/en/> -accessed June 24th 2015.
- Memish ZA, Mishra N, Olival KJ, et al. Middle East respiratory syndrome coronavirus in bats, Saudi Arabia. *Emerg Infect Dis* 2013; **19**:1819–23.
- Reusken CB, Messadi L, Feyisa A, Ularanu H, Godeke GJ, Danmarwa A, et al. Geographic distribution of MERS coronavirus among dromedary camels, Africa. *Emerg Infect Dis* 2014 Aug; **20**(8):1370–4.
- Müller MA, Corman VM, Jores J, Meyer B, Younan M, Liljander A, et al. MERS coronavirus neutralizing antibodies in camels, Eastern Africa, 1983–1997. *Emerg Infect Dis* 2014 Dec; **20**(12):2093–5.
- Haagmans BL, Al Dhahiry SH, Reusken CB, et al. Middle East respiratory syndrome coronavirus in dromedary camels: an outbreak investigation. *Lancet Infect Dis* 2014; **14**:140–5.
- WHO Report: Middle East Respiratory Syndrome coronavirus (MERS-CoV) – Saudi Arabia-Disease outbreak news <http://www.who.int/csr/don/04-june-2015-mers-saudi-arabia/en/>.
- Zumla A, Mwaba P, Bates M, Al-Tawfiq JA, Maeurer M, Memish ZA. The Hajj pilgrimage and surveillance for Middle East Respiratory syndrome coronavirus in pilgrims from African countries. *Trop Med Int Health* 2014 Jul; **19**(7):838–40.
- Annan A, Owusu M, Marfo KS, Larbi R, Sarpong FN, Adu-Sarkodie Y, et al. High prevalence of common respiratory viruses and no evidence of Middle East Respiratory Syndrome Coronavirus in Hajj pilgrims returning to Ghana, 2013. *Trop Med Int Health* 2015 Jun; **20**(6):807–12.
- Memish ZA, Zumla A, Alhakeem RF, Assiri A, Turkestani A, Al Harby KD, et al. Hajj: infectious disease surveillance and control. *Lancet* 2014 Jun 14; **383**(9934):2073–82.
- Al-Tawfiq JA, Zumla A, Memish ZA. Travel implications of emerging coronaviruses: SARS and MERS-CoV. *Travel Med Infect Dis* 2014 Sep-Oct; **12**(5):422–8.
- Memish ZA, Almasri M, Turkestani A, Al-Shangiti AM, Yezli S. Etiology of severe community-acquired pneumonia during the 2013 Hajj-part of the MERS-CoV surveillance program. *Int J Infect Dis* 2014 Aug; **25**:186–90.
- Al-Tawfiq JA, Memish ZA. An update on Middle East respiratory syndrome: 2 years later. *Expert Rev Respir Med* 2015 Jun; **9**(3):327–33.
- Soliman T, Cook AR, Coker RJ. Pilgrims and MERS-CoV: what's the risk? Emerging Themes in *Epidemiology* 2015; **12**:1–3.
- Assiri A, McGeer A, Perl TM, Price CS, Al Rabeeah AA, Cummings DA, et al., KSA MERS-CoV Investigation Team. Hospital outbreak of Middle East respiratory syndrome coronavirus. *N Engl J Med* 2013 Aug 1; **369**(5):407–16.
- Drosten C, Muth D, Corman VM, Hussain R, Al Masri M, HajjOmar W, et al. An observational, laboratory-based study of outbreaks of middle East respiratory syndrome coronavirus in Jeddah and Riyadh, kingdom of Saudi Arabia, 2014. *Clin Infect Dis* 2015 Feb 1; **60**(3):369–77.
- Oboho IK, Tomczyk SM, Al-Asmari AM, Banjar AA, Al-Mugti H, Aloraini MS, et al. 2014 MERS-CoV outbreak in Jeddah—a link to health care facilities. *N Engl J Med* 2015 Feb 26; **372**(9):846–54.
- Al-Abdallat MM, Payne DC, Alqasrawi S, Rha B, Tohme RA, Abedi GR, et al., Jordan MERS-CoV Investigation Team. Hospital-associated outbreak of Middle East respiratory syndrome coronavirus: a serologic, epidemiologic, and clinical description. *Clin Infect Dis* 2014 Nov 1; **59**(9):1225–33.
- Petersen E, Hui DS, Perlman S, Zumla A. Middle East Respiratory Syndrome-advancing the public health and research agenda on MERS- lessons from the South Korea outbreak. *Int J Infect Dis* 2015 Jun 10; **36**:54–5.
- Al-Tawfiq JA, Zumla A, Memish ZA. Coronaviruses: severe acute respiratory syndrome coronavirus and Middle East respiratory syndrome coronavirus in travelers. *Curr Opin Infect Dis* 2014 Oct; **27**(5):411–7.
- Infection prevention and control of epidemic- and pandemic-prone acute respiratory infections in health care - WHO Guidelines. Geneva, World Health Organization, 2014. Available at http://apps.who.int/iris/bitstream/10665/112656/1/9789241507134_eng.pdf.
- WHO Interim guidance WHO/MERS/IPC/15.1 Infection prevention and control during health care for probable or confirmed cases of Middle East respiratory syndrome coronavirus (MERS-CoV) infection. Update 4th June 2015. http://apps.who.int/iris/bitstream/10665/174652/1/WHO_MERS_IPC_15.1_eng.pdf?ua=1.
- PHE Report: Infection control advice – Middle East respiratory syndrome coronavirus (MERS-CoV). 28th June 2013 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/361569/MERS-CoV_infection_control.pdf.
- Pavli A, Tsiodras S, Maltezou HC. Middle East respiratory syndrome coronavirus (MERS-CoV): prevention in travelers. *Travel Med Infect Dis* 2014 Nov-Dec; **12**(6 Pt A):602–8. doi: 10.1016/j.tmaid.2014.10.006. Epub 2014 Oct 19.
- Al-Tawfiq JA, Zumla A, Gautret P, Gray GC, Hui DS, Al-Rabeeah AA, et al. Surveillance for emerging respiratory viruses. *Lancet Infect Dis* 2014 Oct; **14**(10):992–1000.
- Kasolo F, Yoti Z, Bakayita N, et al. IDSR as a platform for implementing IHR in African countries. *Biosecur Bioterror* 2013; **11**:163–9.
- Kebede S, Conteh IN, Steffen CA, et al. Establishing a national influenza sentinel surveillance system in a limited resource setting, experience of Sierra Leone. *Health Res Policy Syst* 2013; **11**:22.
- Zumla A, Al-Tawfiq JA, Enne VI, Kidd M, Drosten C, Breuer J, et al. Rapid point of care diagnostic tests for viral and bacterial respiratory tract infections—needs, advances, and future prospects. *Lancet Infect Dis* 2014 Nov; **14**(11):1123–35.
- Zumla A, Memish ZA, Maeurer M, Bates M, Mwaba P, Al-Tawfiq JA, et al. Emerging novel and antimicrobial-resistant respiratory tract infections: new drug development and therapeutic options. *Lancet Infect Dis* 2014 Nov; **14**(11):1136–49.

Alimuddin Zumla^a
 Roxana Rustomjee^b
 Francine Ntoumi^c
 Peter Mwaba^d
 Matthew Bates^e
 Markus Maeurer^d
 David S. Hui^f
 Eskild Petersen^{g,*}

^aDivision of Infection and Immunity, University College London, and NIHR Biomedical Research Centre, UCL Hospitals NHS Foundation Trust, London, United Kingdom

^bSouth Africa Medical Research Council, Cape Town, South Africa

^cFondation Congolaise pour la Recherche Médicale, Brazzaville, Republic of Congo

^dUNZA-UCLMS Research and Training Project, University Teaching Hospital, Lusaka, Zambia

^eTherapeutic Immunology, Departments of Laboratory Medicine and Microbiology, Tumour and Cell Biology, Karolinska Institute, Stockholm, Sweden

^fDivision of Respiratory Medicine and Stanley Ho Center for Emerging Infectious Diseases, The Chinese University of Hong Kong, Prince of Wales Hospital, Hong Kong Special Administrative Region, China

^gDepartment of Infectious Diseases and Clinical Microbiology, Aarhus University Hospital Skejby, Aarhus, Denmark

*Corresponding author.

E-mail addresses: a.zumla@ucl.ac.uk (A. Zumla),
Roxana.Rustomjee@mrc.ac.za (R. Rustomjee),
ftntoumi@fcrm-congo.com (F. Ntoumi),
pbmwaba2000@gmail.com (P. Mwaba),
matthew.bates@ucl.ac.uk (M. Bates),
Markus.Maeurer@ki.se (M. Maeurer),
dschui@cuhk.edu.hk (D.S. Hui),
eskildp@dadlnet.dk (E. Petersen).

Corresponding Editor: Eskild Petersen, Aarhus, Denmark.