



SHORT COMMUNICATION

Natural FCoV infection: cats with FIP exhibit significantly higher viral loads than healthy infected cats

Anja Kipar ^{Drmedvet, DiplECVP^{1,2*}}, Keith Baptiste ^{BVMS, PhD, DiplACVIM²}, Andreas Barth ¹,
Manfred Reinacher ^{Prof Drmedvet, DiplECVP¹}

¹Institut für Veterinär-Pathologie,
Justus-Liebig-Universität Giessen,
Giessen, Germany

²Department of Veterinary
Pathology, Faculty of Veterinary
Science, University of Liverpool,
Crown Street, Liverpool L69 7ZJ,
UK

Natural feline coronavirus (FCoV) infection has been shown to not only induce intestinal infection with viral shedding, but also systemic infection which either remains without clinical signs or leads to feline infectious peritonitis (FIP). As systemic infection is not the key event in the development of FIP, the question arises as to whether a potential difference in viral load might be of importance. Therefore, the purpose of this study was to quantitatively assess feline coronavirus (FCoV) RNA loads in haemolymphatic tissues of healthy, long-term FCoV-infected cats and cats with FIP. In cats that died from FIP, viral loads were significantly higher, indicating a higher rate of viral replication or a reduced capacity for viral clearance in cats developing and/or suffering from FIP.

© 2005 ESFM and AAFP. Published by Elsevier Ltd. All rights reserved.

Date accepted: 12 July 2005

Feline infectious peritonitis (FIP) is a well-known and widely distributed coronavirus (CoV)-induced systemic disease in cats, characterised by fibrinous-granulomatous serositis with protein-rich effusions into body cavities, granulomatous-necrotising phlebitis and periphlebitis and granulomatous inflammatory lesions in several organs (Hayashi et al 1977, Weiss and Scott, 1981a, b, Kipar et al 1998, 2005).

Feline CoV (FCoV) is transmitted via the faecal-oral route and infects primarily enterocytes (Pedersen 1995), but subsequently spreads systemically via a monocyte-associated viraemia (Gunn-Moore et al 1998, Kipar et al 1999, 2005, Meli et al 2004, de Groot-Mijnes et al 2005). FCoV-infected circulating monocytes are responsible for both viral dissemination and, in an activated state, development of vasculitis (Weiss and Scott 1981a, b, Pedersen 1995, Kipar et al 2005). Also, regardless of the development of FIP, FCoV infection is associated with increased numbers of (proliferating) monocytes/macrophages in haemolymphatic tissues (Kipar et al

1999, 2001a). On the other hand, it has been suspected that higher levels of viral replication could be a prerequisite for the occurrence of the virulent viral mutants among the quasispecies cloud generally seen in infected animals as a consequence of the high rate of viral mutation (Vennema et al 1998, Gunn-Moore et al 1999). Considering also the role of monocytes/macrophages as mediators of both viral replication and formation of granulomatous lesions in FIP, the question arises as to whether the long-term outcome of FCoV infection might relate to the degree of viral replication within an infected individual. We, therefore, performed a relative quantitation of FCoV RNA loads in haemolymphatic tissues in naturally FCoV-infected cats with and without FIP.

The study was performed on 15 routinely necropsied cats which had died, with or without euthanasia, due to FIP (group 1). Animals ranged from 5 months to 4 years of age and all exhibited lesions typical for FIP, namely fibrinous to granulomatous peritonitis and pleuritis, often associated with effusion, and/or granulomatous lesions in various organs, lymph nodes and the central nervous system. Brain and spinal cord

*Corresponding author. Tel: +44-151-794-4260 (Secretary ext 4265);
Fax: +44-151-794-4268. E-mail: akipar@liverpool.ac.uk

involvement was characterised by a granulomatous leptomeningitis, often with granulomatous phlebitis and periphlebitis. The latter was also observed occasionally in eyes, renal cortices and lungs. Diagnosis of FIP was confirmed by immunohistological demonstration of FCoV antigen within macrophages in lesions (Kipar et al 1998). Group 2 consisted of 13 apparently healthy specific pathogen-free cats, aged 1 year, who had been housed for 30 weeks with cats developing FIP (Kipar et al 1999). All group 2 cats had shown positive results in tests for CoV antibodies, circulating FCoV-specific immune complexes and monocyte-associated FCoV viraemia (Kipar et al 1999).

Samples from spleen, mesenteric lymph nodes and bone marrow were collected immediately after death from all cats in areas without macroscopic FIP lesions, and frozen at -80°C for RNA extraction. Additional tissue samples were fixed in 10% buffered formalin, routinely embedded in paraffin and tested by immunohistology for the presence of FCoV antigen (Kipar et al 1998, 2005).

For RNA extraction, approximately 100 mg of tissue was homogenised with a tissue homogeniser (PCR Tissue Homogenizing Kit, Süd-Laborbedarf GmbH, Gauting, Germany) and lysed in each 700 μl lysis buffer. Total RNA was extracted with a commercially available kit (RNeasy Mini kit; QIAGEN, Hilden, Germany) according to the manufacturer's protocol and dissolved in 30 μl RNase-free water. A one-tube real-time TaqMan-RT-PCR, detecting a conserved 102 bp fragment of the FCoV 7b gene, using primers and probes already described (Gut et al 1999, Meli et al 2004), was performed, using an automated fluorometer (ABI Prism 7700 Sequence Detection System, PE Applied Biosystems, Weiterstadt, Germany). RNA extracts from *in vitro* FIPV-infected monolayers of feline embryonal cells served as positive controls. Adequate negative controls were included into each TaqMan PCR run. In parallel, cDNA was synthesised and real-time TaqMan-PCR for feline GAPDH was performed according to previously published protocols (Kipar et al 2001b).

Relative quantification of FCoV transcripts was undertaken by the comparative C_T method and is reported as relative transcription or the n -fold differences of FCoV transcripts relative to the calibrator cDNA (fGAPDH) (Kipar et al 2001b). For samples where an FCoV-RNA signal was not observed after 45 cycles, an approximated C_T value was created. In order to get this

value, the lowest C_T for fGAPDH obtained from the sample with the highest expression of GAPDH was subtracted from 45 (number of cycles). The resulting value was rounded to the next higher integer value. For each sample, relative quantitation of FCoV amounts was achieved by calculating the difference between the fGAPDH C_T value and the FCoV C_T value (ΔC_T). These served to both normalise for differences in the amount of total nucleic acid added to each reaction and to compare the amount of FCoV RNA in the different tissue samples, and expressed as $2^{-\Delta C_T}$.

Initially, all data were assessed using descriptive statistics. Data analysis involved comparisons between cats in groups 1 and 2. First, comparisons were made between groups 1 and 2 according to the organ examined (eg, bone marrow, spleen, mesenteric lymph nodes) using a Mann–Whitney test with unpaired ties. For instance, the relative virus quantitation in group 1 cats' bone marrow was compared to relative virus quantitation in group 2 cats' bone marrow. Finally, the relative virus quantitation of each organ within groups was compared using a Kruskal–Wallis analysis of variance. Significance was defined as a P -value <0.05 .

In all cats with FIP (100%) and 11 of the 13 FCoV-infected cats without FIP (85%), FCoV RNA was detected in at least one of the tissues examined. In cats with FIP, test positive samples included spleen in 60% (9/15), mesenteric lymph nodes in 87% (13/15) and bone marrow in 67% (10/15). In five cases (33%), only one organ tested positive, whereas two organs were positive in three cases (20%) and all organs tested positive in the remaining seven cases (47%). In FCoV-infected cats without FIP, test positive samples included spleen in 38% (5/13), mesenteric lymph nodes in 33% (4/12) and bone marrow in 46% (6/13) each. In seven cases, only one organ tested positive, in the remaining four cases, two organs tested positive.

The relative FCoV genome content in haemolymphatic tissues was variable but overall significantly higher in cats with FIP than in FCoV-infected cats without FIP (bone marrow: $P = 0.0273$, mesenteric lymph nodes: $P = 0.0029$, spleen: $P = 0.0037$; Fig 1). There was no significant difference in the relative virus load between the different organs in each group of animals.

FCoV antigen was not detected by immunohistology in tissues from FCoV-infected cats without FIP and in lesion-free areas of tissues from FIP cats.




Fig 1. Relative FCoV loads in haemolymphatic tissues of FCoV-infected cats and cats with FIP (box and whisker plots). (1) = group 1: cats with FIP ($n = 15$), (2) = group 2: FCoV-infected cats without FIP ($n = 13$), BM = bone marrow, LN = lymph nodes.

This study represents the first attempt to quantify the viral load in FCoV-infected cats with and without FIP. We have chosen to examine the haemolymphatic tissues, as these represent a major site for the accumulation of monocytes/macrophages (ie, cells responsible for viral replication) (Stoddart and Scott 1989; Gunn-Moore et al 1998, Kipar et al 2005, Simons et al 2005) and likely best reflect the actual viral load.

All cats with FIP and the vast majority of FCoV-infected cats without FIP (85%) exhibited systemic infection, demonstrated by the highly sensitive real-time-RT-PCR method (Gut et al 1999). Findings confirm other studies, showing FCoV long-term, persistent viraemia and systemic spread regardless of the development of FIP (Gunn-Moore et al 1998, Kipar et al 1999, Meli et al 2004, de Groot-Mijnes et al 2005). However, our study identified generally higher viral loads in the haemolymphatic tissues of cats with FIP than in FCoV-infected cats without disease.

Previous *in vitro* studies provided evidence that FCoV virulence is associated with higher effectiveness of infecting and replicating in peritoneal macrophages (Stoddart and Scott 1989). This is further supported by a recent study which showed by means of a reverse transcriptase-polymerase chain reaction specific for FCoV

mRNA that replication of FCoV in circulating monocytes occurs mainly in cats with FIP (Simons et al 2005). We could show that cats with FIP have higher viral loads in spleen, lymph nodes and bone marrow, the major organs harbouring monocytes/macrophages, the cells mediating FCoV viraemia. Taken together, all these findings indicate that enhanced viral replicative capacity could be a key feature in the development of FIP. This could be characteristic for the virulent variants of FCoV which are predominantly formed within the individual infected host and characterised by deletions in genes encoding non-structural proteins of yet unknown function, which develop during replication (Vennema et al 1998, Kennedy et al 2001). On the other hand, it is possible that reduced capacity for viral clearance in cats developing and/or suffering from FIP, possibly overall or only in monocytes/macrophages, is responsible for the higher viral loads in cats with FIP. Evidence for this was provided by a recent study which observed rise in blood viral RNA levels together with immunosuppression in cats with end-stage FIP (de Groot-Mijnes et al 2005). This supports the idea that the viral loads in infected animals are a consequence of individual host factors, particularly the immune response (Herrewegh et al 1997, de Groot-Mijnes et al 2005).

Our study did not examine viral characteristics, particularly viral strains and viral mutations, in the two groups of cats. Therefore, further genetic studies are required to assess the role of viral variation and quasispecies clouds in individual and groups of cats, the replicative capacity of individual FCoV isolates and their direct impact on the outcome of FCoV infection (Vennema et al 1998, Gunn-Moore et al 1999, Kennedy et al 2001). However, together with other recent studies on the development of FIP lesions and the immunological changes associated with FCoV infection and FIP (de Groot-Mijnes et al 2005, Kipar et al 2005), this study provides another important piece of information confirming that FIP develops as a consequence of an interplay between virulent FCoVs and the immune system of the individual infected cat.

Acknowledgements

We wish to thank Dr W. Leukert for the clinical supervision of the FCoV-infected cats and acknowledge the cooperation of Prof R.M. Bohle and Dr L. Fink, Department of Pathology, University of Giessen, who provided the real-time-PCR facilities. This study was supported by a grant from the Deutsche Forschungsgemeinschaft (KI 617/3).

References

- de Groot-Mijnes JD, van Dun JM, van der Most RG, de Groot RJ (2005) Natural history of a recurrent feline coronavirus infection and the role of cellular immunity in survival and disease. *Journal of Virology* **79**, 1036–1044.
- Gunn-Moore DA, Gruffydd-Jones TJ, Harbour DA (1998) Detection of feline coronaviruses by culture and reverse transcriptase-polymerase chain reaction of blood samples from healthy cats and cats with clinical feline infectious peritonitis. *Veterinary Microbiology* **62**, 193–205.
- Gunn-Moore DA, Gunn-Moore FJ, Gruffydd-Jones TJ, Harbour DA (1999) Detection of FCoV quasispecies using denaturing gradient gel electrophoresis. *Veterinary Microbiology* **69**, 127–130.
- Gut M, Leutenegger CM, Huder JB, Pedersen NC, Lutz H (1999) One-tube fluorogenic reverse transcription-polymerase chain reaction for the quantitation of feline coronaviruses. *Journal of Virological Methods* **77**, 37–46.
- Hayashi T, Goto N, Takahashi R, Fujiwara K (1977) Systemic vascular lesions in feline infectious peritonitis. *Japanese Journal of Veterinary Science* **39**, 365–377.
- Herrewegh AAPM, Mähler M, Hedrich HJ, Haagmans BL, Egberink HF, Horzinek MC, Rottier PJM, de Groot RJ (1997) Persistence and evolution of feline coronavirus in a closed cat-breeding colony. *Virology* **234**, 349–363.
- Kennedy M, Boedeker N, Gibbs P, Kania S (2001) Deletions in the 7a ORF of feline coronavirus associated with an epidemic of feline infectious peritonitis. *Veterinary Microbiology* **81**, 227–234.
- Kipar A, Bellmann S, Kremendahl J, Köhler K, Reinacher M (1998) Cellular composition, coronavirus antigen expression and production of specific antibodies in lesions in feline infectious peritonitis. *Veterinary Immunology and Immunopathology* **65**, 243–257.
- Kipar A, Bellmann S, Gunn-Moore DA, Leukert W, Köhler K, Menger S, Reinacher M (1999) Histopathological alterations of lymphatic tissues in cats without feline infectious peritonitis after long-term exposure to FIP virus. *Veterinary Microbiology* **69**, 131–137.
- Kipar A, Köhler K, Leukert W, Reinacher M (2001a) Evaluation of lymphatic tissue activity in cats with spontaneous feline infectious peritonitis. *Journal of Comparative Pathology* **125**, 182–191.
- Kipar A, Leutenegger CM, Hetzel U, Akens MK, Mislin CN, Reinacher M, Lutz H (2001b) Cytokine mRNA levels in isolated feline monocytes. *Veterinary Immunology and Immunopathology* **78**, 305–315.
- Kipar A, May H, Menger S, Weber M, Leukert W, Reinacher M (2005) Morphological features and development of granulomatous vasculitis in feline infectious peritonitis. *Veterinary Pathology* **42**, 321–330.
- Meli M, Kipar A, Müller C, Jenal K, Gönczi E-E, Borel N, Gunn-Moore D, Chalmers S, Lin F, Reinacher M, Lutz H (2004) High viral loads despite absence of clinical and pathological findings in cats experimentally infected with feline coronavirus (FCoV) type I and in naturally FCoV-infected cats. *Journal of Feline Medicine and Surgery* **6**, 69–81.
- Pedersen NC (1995) An overview of feline enteric coronavirus and infectious peritonitis virus infections. *Feline Practice* **23**, 7–20.
- Simons FA, Vennema H, Rofina JE, Pol JM, Horzinek MC, Rottier PJ, Egberink HF (2005) A mRNA PCR for the diagnosis of feline infectious peritonitis. *Journal of Virological Methods* **124**, 111–116.
- Stoddart CA, Scott FW (1989) Intrinsic resistance of feline peritoneal macrophages to coronavirus infection correlates with in vivo virulence. *Journal of Virology* **63**, 436–440.
- Vennema H, Poland A, Foley J, Pedersen NC (1998) Feline infectious peritonitis viruses arise by mutation from endemic feline enteric coronaviruses. *Virology* **243**, 150–157.
- Weiss RC, Scott FW (1981a) Pathogenesis of feline infectious peritonitis: nature and development of viraemia. *American Journal of Veterinary Research* **4**, 382–390.
- Weiss RC, Scott FW (1981b) Pathogenesis of feline infectious peritonitis: pathologic changes and immunofluorescence. *American Journal of Veterinary Research* **42**, 2036–2048.