



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

2019-nCoV transmission through the ocular surface must not be ignored

Chaolin Huang and colleagues¹ reported the epidemiology, symptoms, and treatment of patients infected by the 2019 novel coronavirus (2019-nCoV) in Wuhan, China. As ophthalmologists, we believe that transmission of 2019-nCoV through the eyes was ignored.

On Jan 22, Guangfa Wang, a member of the national expert panel on pneumonia, reported that he was infected by 2019-nCoV during the inspection in Wuhan.² He wore an N95 mask but did not wear anything to protect his eyes. Several days before the onset of pneumonia, Wang complained of redness of the eyes. Unprotected exposure of the eyes to 2019-nCoV in the Wuhan Fever Clinic might have allowed the virus to infect the body.²

Infectious droplets and body fluids can easily contaminate the human conjunctival epithelium.³ Respiratory viruses are capable of inducing ocular complications in infected patients, which then leads to respiratory infection.⁴ Severe acute respiratory syndrome coronavirus (SARS-CoV) is predominantly transmitted through direct or indirect contact with mucous membranes in the eyes, mouth, or nose.⁵ The fact that exposed mucous membranes and unprotected eyes increased the risk of SARS-CoV transmission⁴ suggests that exposure of unprotected eyes to 2019-nCoV could cause acute respiratory infection.

Thus, Huang and colleagues¹ should have analysed conjunctival scrapings from both confirmed and suspected 2019-nCoV cases during the onset of symptoms. The respiratory tract is probably not the only transmission route for 2019-nCoV, and all ophthalmologists examining suspected cases should wear protective eyewear.

We declare no competing interests.

*Cheng-wei Lu, Xiu-fen Liu,
Zhi-fang Jia
lcwchina800@sina.com

Department of Ophthalmology (C-WL, X-FL) and
Division of Clinical Research (Z-FJ), First Hospital of
Jilin University, Changchun 130021, China

- 1 Huang C, Wang YM, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- 2 Dai X. Peking University Hospital Wang Guangfa disclosed treatment status on Weibo and suspected infection without wearing goggles. Jan 22, 2020. <http://www.bjnews.com.cn/news/2020/01/23/678189.html> (accessed Jan 24, 2020).
- 3 Olofsson S, Kumlin U, Dimock K, Arnberg N. Avian influenza and sialic acid receptors: more than meets the eye? *Lancet Infect Dis* 2005; **5**: 184–88.
- 4 Belser JA, Rota PA, Tumpey TM. Ocular tropism of respiratory viruses. *Microbiol Mol Biol Rev* 2013; **77**: 144–56.
- 5 Peiris JS, Yuen KY, Osterhaus AD, Stohr K. The severe acute respiratory syndrome. *N Engl J Med* 2003; **349**: 2431–41.



Published Online
February 6, 2020
[https://doi.org/10.1016/S0140-6736\(20\)30313-5](https://doi.org/10.1016/S0140-6736(20)30313-5)

Submissions should be made via our electronic submission system at <http://ees.elsevier.com/thelancet/>