

ERRATUM

High water vs. *ad libitum* water intake for autosomal dominant polycystic kidney disease: a randomized controlled feasibility trial

R. El-Damanawi^{1,2}, M. Lee³, T. Harris⁴, L.B. Cowley^{2,5}, S. Bond², H. Pavey², R.N. Sandford⁶, I.B. Wilkinson^{1,2}, F.E. Karet Frankl⁶ and T.F. Hiemstra^{1,2}

From the ¹Division of Experimental Medicine and Immunotherapeutics, Department of Medicine, University of Cambridge, Cambridge, ²Cambridge Clinical Trials Unit, Cambridge University Hospitals NHS Foundation Trust, Cambridge, ³Division of Anaesthesia, Department of Medicine, University of Cambridge, Cambridge, ⁴PKD Charity, 91 Royal College, London, ⁵Patient Led Research Hub, Cambridge NIHR Biomedical Research Centre, Cambridge and ⁶Department of Medical Genetics, University of Cambridge, Hills Road, Cambridge, UK

Address correspondence to Dr T.F. Hiemstra, Cambridge Clinical Trials Unit, Box 401 Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0QQ, UK. email: tfh24@cam.ac.uk

QJM: An International Journal of Medicine 2020; 113:258–65.

doi: 10.1093/qjmed/hcz278

This manuscript has been amended to correct an error in the abstract. The section 'high water (HW) intake targeting urine osmolality (UOsm) ≥ 270 mOsm/kg'

Should have read

'high water (HW) intake targeting urine osmolality (UOsm) ≤ 270 mOsm/kg'

The Publisher apologises for any confusion caused.