



Published in final edited form as:

Cancer Causes Control. 2020 April ; 31(4): 291–302. doi:10.1007/s10552-020-01283-3.

Obesity is an Initiator of Colon Adenomas but not a Promoter of Colorectal Cancer in the Black Women's Health Study

Chiranjeev Dash^{1,2}, Jeffrey Yu³, Sarah Nomura¹, Jiachen Lu¹, Lynn Rosenberg³, Julie R. Palmer³, Lucile L. Adams-Campbell^{1,2}

¹Georgetown Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, DC, USA.

²Office of Minority Health and Health Disparities Research, Georgetown Lombardi Comprehensive Cancer Center, 1000 New Jersey Ave SE, Washington, DC, 20003, USA.

³Slone Epidemiology Center at Boston University, Boston, MA, USA.

Abstract

Purpose—Evidence for the association of anthropometrics with colorectal neoplasms is limited for African Americans.

Methods—We examined anthropometric measures with both colorectal adenoma and colorectal cancer (CRC) risk in the ongoing Black Women's Health Study. In a nested case-control analysis, 954 cases of colorectal adenoma were compared with 3,816 polyp-free controls, matched on age and follow-up time. For the CRC analyses, 413 incident CRC cases were identified over a 16-year follow up (802,783 person-years). Adenoma cases and CRC were verified by medical record review. We used multivariable conditional logistic regression analyses (for adenoma) and Cox proportional hazards analyses (for CRC) that included anthropometric exposures and selected confounders.

Results—Overall body mass index (BMI) and other anthropometric factors were not associated with colorectal adenoma or cancer risk in Black women. However, increased risk of adenoma (but not CRC) was observed among especially related to adenomas in the proximal colon. Among women 50 years of age, risk of proximal adenoma increased 14% (95% CI: 1.00, 1.31), 35% (95% CI: 1.12, 1.63), and 25% (0.93, 1.68) with each standard deviation increase in BMI, waist circumference, and waist to hip ratio, respectively. None of the anthropometric factors were associated with young onset CRC or adenoma risk.

Conclusion—Our results suggest that obesity might be an initiator for colon adenomas but not a promoter for colorectal cancer among Black women.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <https://www.springer.com/aam-terms-v1>

Correspondence to: Lucile Adams-Campbell, Office of Minority Health and Health Disparities Research, Georgetown Lombardi Comprehensive Cancer Center, 1000 New Jersey Ave SE, Washington, D.C. 20003. Phone: 202-687-5367. lla9@georgetown.edu.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Keywords

African American women; colorectal adenoma; colorectal cancer; BMI; obesity

Numerous studies have evaluated the association of obesity, primarily body mass index (BMI), with colorectal cancer (CRC) and adenomas.[1–4] Recent studies, including two meta-analyses, suggest that BMI is more strongly associated with CRC and adenomas in men than women.[4, 5] However, it is possible that BMI might not be the most relevant anthropometric measure of obesity-associated colorectal neoplasia risk. Measures of abdominal adiposity, such as waist circumference (WC) and waist to hip ratio (WHR), might be more strongly associated with colorectal neoplasia risk than body mass index (BMI) among women.

Abdominal adiposity, through its effects on pro-inflammatory, oxidative stress, and metabolic pathways, has been hypothesized to be a better biological measure of obesity-associated CRC than BMI.[6] Additionally, the positive association of abdominal obesity with estradiol, a significant modulator of the estrogen pathway hypothesized to be involved in colorectal carcinogenesis, provides biological support for waist circumference and WHR as more relevant measures of obesity among women.[7]

Anthropometric measures other than BMI and abdominal adiposity have not been well studied as risk factors for CRCs and adenomas. Some studies have suggested adult weight gain as a risk factor for colorectal neoplasms.[8] In addition, given the strong correlation between BMI and WC, it has been suggested that the estimation of independent effects of these 2 anthropometric measures as epidemiologic risk factors is difficult even in statistical models that include both factors.[9] A new measure, a body shape index (ABSI), was recently defined as $WC/(BMI^{2/3} \text{ Height}^{1/2})$. [10] ABSI was derived as a measure of abdominal adiposity that has little correlation with either weight or BMI.

Relatively few studies have investigated multiple anthropometric measures as risk factors for colorectal adenoma and cancer among women and none have been adequately powered for these analyses among African American women. In a previous analysis from the Black Women's Health Study (BWHS), we found associations of both body mass index (BMI) and waist-to-hip ratio (WHR) with colorectal polyps, but the association with polyp location (proximal versus distal) was not assessed.[11] In the present study, we investigated association of BMI, WC, WHR, weight gain in adulthood, and ABSI with risk of colorectal adenoma and cancer among African American women in the BWHS.

METHODS

Study population

The BWHS is a prospective cohort study of African American women from across the United States. In 1995, 59,000 women aged 21 to 69 years enrolled by responding to health questionnaires mailed to subscribers of Essence magazine, members of several African American professional associations, and friends of early respondents.[12] Approximately equal proportions were from the Northeast, South, Midwest, and West. [13] Respondents

completed 14-page questionnaires on demographics, health status, and medical and lifestyle variables. The baseline questionnaire obtained information on adult height, current weight, demographic characteristics, reproductive history, medical history, use of medications, use of cigarettes and alcohol, and usual diet. Since 1995, follow-up questionnaires have been sent every two years to update information on reproductive history and other exposures and identify new occurrences of cancer and other serious illnesses. Follow-up of the baseline cohort has been successful with a follow-up rate of eighty-seven percent of all potential person-years through 2013. Approval for the study was obtained from Boston University Institutional Review Board.

Study design

We used a nested case-control design to investigate the association of anthropometric factors with adenoma and a prospective cohort design to investigate the association of anthropometric factors with CRC risk.

Adenoma case and control ascertainment—Participants were asked about a list of diseases and date of first diagnosis on baseline and follow-up questionnaires. In 1999, “colon or rectal polyps” was added to the list of illnesses for which participants were asked to indicate whether they had received a first diagnosis in the previous two years. Women who reported a colon or rectal polyp were asked for permission to obtain medical records relevant to the colonoscopy. Characteristics of all women who reported a polyp were similar to those of women for whom medical records were obtained and confirmed an adenoma. Mean BMI in 1995, and mean waist to hip ratio in 1995 are 28.9 kg/m² and 0.79 among all women who reported a polyp and 28.5 kg/m² and 0.79 among women with a confirmed adenoma.

Cases in the present analyses were colorectal adenomas confirmed by pathology reports and first identified by self-report of colorectal “polyp” on any of the 1999 through 2011 follow-up questionnaires. There were 954 confirmed adenomas from among the 23,804 women who reported a colonoscopy or sigmoidoscopy during the follow-up period from 1997 to 2011 and had not had a colorectal polyp or any cancer at the start of follow-up in 1997.

A risk set sampling approach was used to select controls from among participants who reported undergoing a colonoscopy or sigmoidoscopy but had not reported a colorectal polyp during or prior (< 10 years) to the follow-up period in which the index case reported an incident adenoma diagnosis. Four controls were randomly selected from the list of eligible controls, matched to cases on age and follow-up period at the time of adenoma diagnosis. Relevant exposure data for the controls were abstracted from the questionnaires prior to the “index period” (year for which the index case reported a polyp). Women with cancer (including colorectal cancer), polyps other than adenomas, and women for whom a medical record for polyp review could not be obtained, were excluded from the analysis.

CRC case ascertainment—Colon and rectal cancer cases (ICD-10 colon cancer: C18.0-C18.9 and C26.0; ICD-10 rectal cancer: C19.9 and C20.9) were identified for follow-up from 1995 through 2011 through self-report on the follow-up questionnaires, through linkage with cancer registries in 24 states in which 95% of participants live, and through

death records. Pathology data were obtained from hospitals or registries for confirmation. To date, of 397 self-reported cases occurring in the BWHS during follow-up for which pathology data were obtained, 394 (99 %) were confirmed as colorectal cancer. Given the accuracy of self-report, all self-reported cases were included in the present analyses unless found to be incorrectly reported based on pathology data.

Assessment of anthropometric factors—In 1995, we collected information on self-reported height (in feet and inches), current weight (in pounds), and weight at age 18 (in pounds). We also asked each participant to measure her waist circumference at the level of the umbilicus (in inches) and hip circumference at its widest location (in inches). Current weight was updated every 2 years by follow-up questionnaire and waist/hip circumferences were updated in 2005. Height (1995) and current weight were used to calculate body mass index (BMI) (kg/m^2); waist circumference was divided by hip circumference to calculate WHR; and adult weight change was calculated by subtracting weight at age 18 from participant-reported current weight. ABSI was calculated using the following formula: $\text{WC} / (\text{BMI}^{2/3} \text{ height}^{1/2})$, where WC and height are in m, and BMI is in kg/m^2 . Self-reported weight (Spearman correlation=0.97) and height (Spearman correlation=0.93) were highly correlated with technician measurements in a BWHS validation study.[14, 15]

Assessment of covariates—Covariates for analysis were selected a priori from the literature. Data on age, education, cigarette smoking, regular (at least 3 days a week) aspirin use, alcohol intake, menopausal status, and postmenopausal hormone therapy were collected on the baseline questionnaire (1995) and updated based on data reported on the follow-up questionnaires. In the 1997 and subsequent questionnaires, participants provided information on the number of hours spent each week on vigorous exercise such as basketball, swimming, running and aerobics. Information on education was obtained in 1995 and information on family history of colorectal cancer in a first-degree relative was obtained in 1999. Women were classified as premenopausal if they were still menstruating and as postmenopausal if they had a natural menopause (no periods for at least a year) or bilateral oophorectomy. Women with hysterectomy but without a bilateral oophorectomy were classified as postmenopausal if they were above the age of 56, and as premenopausal if they were below 43 years of age. Women who did not report menopausal status or had undergone hysterectomy without a bilateral oophorectomy and were age 43–56 were classified as having “unknown” menopause status. Weekly servings of fruits and vegetables, total red meat intake, and total daily energy intake were derived from the 68-item modified version of the National Cancer Institute (NCI)–Block food frequency questionnaire administered to all participants at baseline (1995) and in the 2001 questionnaire.[16] Dietary variables were derived from the food frequency questionnaire administered in 1995 if the index period was prior to 2001 and from the 2001 food frequency questionnaire if the index period was at or after the 2001 follow up. Time-varying covariates were reassigned for every 2 years of follow-up by using the Andersen-Gill data structure.[17] This creates a new record for every follow-up cycle at which the participant is at risk, and assigns covariate values for that specific questionnaire cycle. For adenoma cases in the nested case-control analysis, covariates were based on the questionnaires administered in the cycle prior to when the polyp (later determined to be an adenoma) was reported (index period). For matched

controls, covariates were also based on responses in the questionnaire cycle before the “index period”.

Statistical Analysis—Baseline age-standardized means (continuous variables) and proportions (categorical variables) were calculated across baseline BMI categories for population characteristics. Anthropometric variables were analyzed as continuous (with effect estimates for adenoma and CRC risk per 1 standard deviation increase reported) and categorical variables. Tests for linearity assumption were conducted using restricted cubic spline regression for models with continuous anthropometric variables, and no deviations from linearity were observed for any anthropometric variable. WC, WHR, and ABSI were categorized in quintiles, BMI was categorized using World Health Organization recommended standardized categories (<18.5, 18.5–24.9, 25–29.9, 30 kg/m²), and weight gain since age 18 in 5 categories (<10, 10–14, 15–19, 20–24, 25 pounds). We used conditional logistic regression to estimate age-adjusted and multivariable-adjusted odds ratios (ORs) and 95% confidence intervals (CIs) for risk of colorectal adenoma in association with anthropometric factors. Associations between anthropometrics and colorectal cancer incidence were evaluated using Cox proportional hazards regression (PROC PHREG) using the Andersen-Gill data structure for time-varying exposures and covariates. Person-years were calculated from baseline until the occurrence of colorectal cancer, loss to follow-up, death, or the end of follow-up in 2011. Since colonoscopy screening may alter the natural history and subsequent risk of CRC through removal of preneoplastic adenomas, we conducted sensitivity analysis by excluding all cases of diagnosed adenomas in the CRC-anthropometrics analysis. In the multivariable models, we adjusted for the following potential confounders: age, education, smoking status, alcohol intake, family history of colorectal cancer in first-degree relative, regular aspirin use, menopausal status, vigorous activity, total energy intake, fruit and vegetable intake, and red meat intake. In addition, models with BMI and weight change since age 18 as the primary exposure variables adjusted for BMI at age 18. Models with WC or WHR as the exposure variables did not adjust for BMI because of the high correlation of BMI with these variables. Instead, ABSI was used as a measure of abdominal obesity uncorrelated with BMI. Tests for linear trend in models where anthropometric measures were treated as categorical variables were performed by assigning the median value for each category/quintile and modeling this variable as a continuous variable. To determine whether associations differed by age at diagnosis of adenoma or cancer (<50, 50 years of age) we stratified primary analyses by age at diagnosis. For the CRC analysis (cohort), person-time contributed by the participants before they reached 50 years of age was the denominator for the CRC rates in the “<50 years of age” stratum; and participants who did not develop cancer prior to 50 years of age were censored at age 50 for this stratum. Person-time was similarly calculated for the “50 years of age” stratum. We assessed effect modification of the association between adenoma and anthropometrics by age. Interaction was assessed using the log-likelihood ratio test that compared models with and without the multiplicative interaction terms of anthropometric factor with age category (e.g., BMI*age). In addition, we analyzed adenoma and cancer location within the colorectum - colon (proximal and distal) and rectal adenoma/cancer in separate models to determine site-specific associations of anthropometric factors with

adenoma/cancer risk. All statistical analyses were performed using SAS version 9.2 (SAS Institute Inc., Cary, NC).

RESULTS

The baseline characteristics of the study population are shown in table 1. At baseline, women in the highest categories of BMI were more likely to be postmenopausal and had lower educational attainment, less vigorous physical activity, greater total energy intake and servings per week of red meat, and a higher prevalence of regular aspirin use.

Adenoma

Overall, ORs for colorectal adenoma were not meaningfully different from 1.0 for categories of increasing BMI, waist circumference, and waist to hip ratio. Among women who gained 10 kgs or more since age 18 risk of adenoma was significantly increased compared to those with less than a 10 kg weight gain. Similarly, adenoma risk was higher with increasing ABSI even though the findings were not statistically significant. (Table 2) In models stratified by age (<50, ≥50 years at diagnosis), associations of all anthropometric factors, except ABSI, were stronger in women with an older age of onset than younger women. One standard unit increase in BMI, waist circumference, and weight change since age 18 were associated with a 12%, 17%, and 8% increased risk of colorectal adenoma among older women whereas among women diagnosed before age 50 the corresponding ORs were below 1.0. Among the older women, a weight gain of 25 kg or more since age 18 relative to <10 kgs was associated with a 38% increase in risk of adenoma (95% CI: 1.03–1.86). However, no significant statistical interaction by age was observed for any anthropometric factor. (Table 2).

Analyses of anthropometric factors in relation to colon adenoma risk by site (proximal) are shown in Table 3. Limited numbers of rectal adenomas among older women in the analytic cohort resulted in unstable effect estimates and these results are not presented. Evidence of an association with measures of body size was observed for proximal adenoma risk. Although not shown, these associations were not observed for distal adenomas. One standard deviation increase in BMI was associated with a 14% increased risk of proximal adenoma (OR: 1.14, 95% CI: 1.00, 1.31) in women ≥50 years. Increasing waist circumference and waist to hip ratio were associated with a 35% (95% CI: 1.12, 1.63) and 25% (95% CI: 0.93, 1.68) increased risk of proximal adenomas with one SD increase, respectively. The association of weight gain since age 18 was also associated with proximal adenoma risk with a 25 kg weight gain (relative to <10 kg weight gain) resulting in a 66% increased proximal adenoma risk (95% CI: 1.13, 2.44) but not distal adenoma risk. Increasing ABSI also appeared to be associated with proximal adenoma risk (OR 1.88, 95% CI: 0.99, 3.56 comparing the highest to the lowest quintile).

Colorectal Cancer Risk (CRC)

A total of 57,386 participants were included in the analysis after excluding those who had not returned any follow-up questionnaire or who had prevalent cancer at baseline. In the 16 years of follow-up from 1995 to 2011, 413 incident CRC cases were identified over 802,783 person-years. In multivariate models, none of the anthropometric factors were associated

with CRC risk among either < 50 or ≥ 50 year old women (Table 4). Associations of anthropometric factors with CRC risk did not differ materially across sites (proximal, distal, rectal) and the results were mostly null (data not shown). In sensitivity analyses where all diagnosed cases of adenomas were excluded from the cohort, the primary associations between anthropometric factors and CRC risk remain unchanged (data not shown).

DISCUSSION

Overall BMI and other anthropometric factors were not associated with the risk of either colorectal adenoma or cancer in this cohort study of Black women in the US. BMI, waist circumference, WHR, weight gain since age 18, and ABSI were modestly associated with increased risk of proximal colon adenomas, among women ≥ 50 years old. However, none of the selected anthropometric factors were associated with CRC risk in either young or old onset CRC. Our findings suggest that among African-American women, obesity may be associated with adenoma risk among older but not younger women; and that obesity is not associated with CRC risk.

Although it has been well established in studies largely comprising White participants that BMI modestly increases the risk of colorectal adenomas[1, 2], more so among men than women, studies in African-Americans are limited, and there is a particular lack of data on adenoma risk in African American women. Data on anthropometric measures other than BMI are even more limited. Our findings are similar to those from two prospective colonoscopy-based studies. Sedjo et al. reported BMI and weight-gain but not WC or WHR associated with adenoma risk.[18] Race-stratified results for BMI reported by Sedjo et al. suggested similar results among Whites and African-Americans, but were underpowered for associations among African-Americans. Murphy et al in a colonoscopy-based study reported a moderate increase in adenoma risk associated with increasing WC and WHR but not BMI among African-American men and women.[19] Our results for BMI and adenoma risk among older women are similar to estimates from meta-analyses based largely on white women. [1, 2] Our results for WHR conform to previously reported estimates of WHR-associated adenoma risks[20, 21], but only for proximal adenomas, that suggested WHR as an independent risk factor for adenoma among older women. Multiple studies have reported higher prevalence of proximal adenomas among African-Americans compared to Caucasians and other races,[22–24] but the reasons for this observation remain unclear. Given the high rates of obesity in African-American women compared to other races,[25] our findings that obesity is an independent risk factor for proximal adenomas might partly explain the higher prevalence of proximal adenomas in this population.

Most studies of BMI in relation to CRC risk have been conducted in primarily White populations. Results from these studies are conflicting, with some suggesting multiple anthropometric factors, including BMI, associated with CRC risks[26–28]; others suggesting markers of abdominal obesity but not BMI as CRC risk factors[29, 30]; and, still others reporting no association between anthropometrics and CRC risk[31–35]. Analyses among postmenopausal women in the Women’s Health Initiative[26] (BMI, WC, WHR, and ABSI), Cancer Prevention Study–II cohort[27] (BMI and WC), and the Nurses’ Health Study[28] (BMI, WC, and WHR) reported positive associations of BMI and other anthropometric

factors with CRC risk. However, in a pooled analysis of 11 Australian cohorts, Harding et al. reported modest CRC risk associated with WC, WHR, and ABSI but not BMI[29]; and results from the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort suggested that abdominal obesity measures (WC and WHR) but not BMI were associated with colon cancer risk among women[30]. In a more recent analysis from the EPIC cohort, Steins Bisschop et al. reported that neither BMI nor weight gain were associated with CRC risk among women[32] Other large studies of White women have reported results similar to our own. In an analysis from the Framingham Heart Study, BMI and waist circumference were not associated with colon cancer risk among women.[31] Similarly, Keimling et al. reported null associations between BMI, WHR, and WC and CRC risk among women in the NIH-AARP cohort[33]; and Renehan et al. reported no associations between BMI at age 18 and weight change since age 18 with CRC risk among women in this cohort.[34] Our results for young onset colorectal cancer among Black women are different from those observed in the Nurses' Health Study II. Liu et al. [36] reported a higher risk of CRC in a cohort of primarily White women comparing overweight (BMI 25–29.9) and obese (BMI ≥ 30) women to those with BMIs between 18.5 to 24.9. However, BMI (and other anthropometric factors) were not associated with young onset CRC risk among Black women in our study.

Only one previous study has investigated risks of colorectal adenoma and CRC with anthropometric factors within the same study. In an analysis from the PLCO, in a primarily White population, Kitahara et al. reported that BMI was not associated with either adenoma or CRC risk among women.[37] That study lacked data on proximal adenomas because of the use of sigmoidoscopy for CRC screening in the PLCO and did not report on WC/WHR. Our study is the first among African-Americans to investigate adenoma and CRC risk in the same study with data on both proximal and distal (although not shown) adenomas and cancer and to assess multiple anthropometric exposures. Our results for CRC are similar to those reported by Kitahara et al., [37] but in our study increasing BMI was associated with proximal colorectal adenoma risk among older women. Although CRCs usually arise from adenomatous polyps, most adenomas will not progress to cancer. Our finding that obesity, after adjusting for diet, physical activity, and other CRC risk factors, might be associated with adenoma but not CRC, suggests that obesity might be important for adenoma formation but not factors related to progression (e.g. dysplasia) to cancer among African-American women.

The strengths of our study include the nested design for adenoma analysis within a large prospective cohort of African American women in the United States, adenoma and cancer outcomes verified by medical records, availability of data to derive multiple anthropometric exposures, high cohort retention resulting in updated measures of exposures and covariates, and detailed information on a large number of covariates. In addition, results from our study are generalizable to most African American women in the United States, but not to those with low educational attainment. More than 95% of the BWHS cohort had a high school education or more at enrollment compared with 83% of African American women in the general population.[38]

Our study was limited by the use of self-reported data for anthropometrics. However, in a validation study of 115 BWHS participants from Washington DC area, Spearman

correlations of self-reported anthropometric versus technician-measured data were >0.90 for height and weight; and >0.70 for WC[39]. Participant-reported polyps were verified by medical record review in our study, but the absence of polyps was not verified.

Misclassification in which participants diagnosed with adenomas failed to report polyps and were therefore included in the control group would attenuate the anthropometrics-adenoma association toward null. However, previous studies have shown that self-report of polyps has a high negative predictive value (94%–100%) for adenomas, and it is unlikely that this was a major source of bias in our study.[40, 41] We had smaller sample sizes for analyses for CRC outcomes by age groups, i.e., <50 or ≥ 50 years, and were underpowered for such analyses, especially for rectal cancer. Given the 5 anthropometric factors we examined including analyses stratified by age it is possible a few associations might have been statistically significant simply due to chance (multiple testing) and our findings from proximal adenomas should be validated in future studies. Similarly, we did not have adequate number of events or follow-up time to effectively determine risk of either adenoma or CRC associated with a very low BMI (<18.5 kg/m²). Finally, control for vigorous activity and red meat, fruit, and vegetable intake might not have been adequate to prevent residual confounding by physical activity and diet. In addition, we did not have data on sedentary behavior available at baseline and could not adjust for this potential confounder in our statistical models.

In summary, BMI and other indicators of obesity were not associated with CRC risk or young-onset CRC risk in this large cohort of African American women. However, BMI, waist circumference, WHR, and weight change were associated with moderately increased risk of proximal colorectal adenomas among African American women above the age of 50. More studies in minority populations are needed to firmly establish the role of obesity in CRC risk.

ACKNOWLEDGEMENTS

Data on colorectal cancer pathology were obtained from several state cancer registries (AZ, CA, CO, CT, DE, DC, FL, GA, IL, IN, KY, LA, MD, MA, MI, NJ, NY, NC, OK, PA, SC, TN, TX, VA), and results reported do not necessarily represent their views. The authors are grateful to the participants and staff of the BWHS.

FUNDING

This work was supported by National Cancer Institute grants R01 CA058420 (L. Rosenberg), UO1 CA164974 (L. Rosenberg). CD is supported by National Cancer Institute training grant K07 CA197112 (C.Dash). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Cancer Institute or the National Institutes of Health.

References

1. Ben Q, An W, Jiang Y, et al. (2012) Body mass index increases risk for colorectal adenomas based on meta-analysis. *Gastroenterology* 142: 762–772. [PubMed: 22245665]
2. Okabayashi K, Ashrafian H, Hasegawa H, et al. (2012) Body mass index category as a risk factor for colorectal adenomas: a systematic review and meta-analysis. *Am J Gastroenterol* 107: 1175–85; quiz 1186. [PubMed: 22733302]
3. Wang J, Yang DL, Chen ZZ, Gou BF (2016) Associations of body mass index with cancer incidence among populations, genders, and menopausal status: A systematic review and meta-analysis. *Cancer Epidemiol* 42.
4. Ma Y, Yang Y, Wang F, et al. (2013) Obesity and risk of colorectal cancer: a systematic review of prospective studies. *PLoS One* 8.

5. Ben Q, An W, Jiang Y, et al. (2012) Body mass index increases risk for colorectal adenomas based on meta-analysis. *Gastroenterology* 142: :762–72. [PubMed: 22245665]
6. Martinez-Useros J, Garcia-Foncillas J (2016) Obesity and colorectal cancer: molecular features of adipose tissue.. *J Transl Med* 22: 21.
7. Chen J, Iverson D (2012) Estrogen in obesity-associated colon cancer: friend or foe? Protecting postmenopausal women but promoting late-stage colon cancer.. *Cancer Causes Control* 23: 1767. [PubMed: 23011535]
8. Karahalios A, English DR, Simpson JA (2015) Weight change and risk of colorectal cancer: a systematic review and meta-analysis.. *Am J Epidemiol* 181: 832. [PubMed: 25888582]
9. World Health Organization (2011) Waist circumference and waist-hip ratio: report of a WHO expert consultation. 8–11.
10. Krakauer NY, Krakauer JC (2012) A new body shape index predicts mortality hazard independently of body mass index. *PLoS One* 7: e39504. [PubMed: 22815707]
11. Wise LA, Rosenberg L, Palmer JR, Adams-Campbell LL (2008) Anthropometric risk factors for colorectal polyps in African-American women.. *Obesity (Silver Spring)* 16: 859. [PubMed: 18239567]
12. Fleming DA, Sheppard VB, Mangan PA, et al. (2006) Caregiving at the end of life: Perceptions of health care quality and quality of life among patients and caregivers. *J Pain Symptom Manage* 31: 407–420. [PubMed: 16716871]
13. Press R, Carrasquillo O, Sciacca RR, Giardina EG (2008) Racial/ethnic disparities in time to follow-up after an abnormal mammogram. *J Womens Health (Larchmt)* 17: 923–930. [PubMed: 18554094]
14. Carter-Nolan PL, Adams-Campbell LL, Makambi K, Lewis S, Palmer JR, Rosenberg L (2006) Validation of physical activity instruments: Black Women’s Health Study. *Ethn Dis* 16: 943–947. [PubMed: 17061751]
15. Wise LA, Palmer JR, Spiegelman D, et al. (2005) Influence of body size and body fat distribution on risk of uterine leiomyomata in U.S. black women. *Epidemiology* 16: 346–354. [PubMed: 15824551]
16. Smith RA, Manassaram-Baptiste D, Brooks D, et al. (2014) Cancer screening in the United States, 2014: a review of current American Cancer Society guidelines and current issues in cancer screening. *CA Cancer J Clin* 64: 30–51. [PubMed: 24408568]
17. Ramirez AG, Perez-Stable EJ, Penedo FJ, et al. (2013) Navigating Latinas with breast screen abnormalities to diagnosis: the Six Cities Study. *Cancer* 119: 1298–1305. [PubMed: 23233265]
18. Sedjo RL, Byers T, Levin TR, et al. (2007) Change in body size and the risk of colorectal adenomas. *Cancer Epidemiol Biomarkers Prev* 16: 526–531. [PubMed: 17372248]
19. Murphy CC, Martin CF, Sandler RS (2015) Racial differences in obesity measures and risk of colorectal adenomas in a large screening population. *Nutr Cancer* 67: 98–104. [PubMed: 25425186]
20. Hong S, Cai Q, Chen D, Zhu W, Huang W, Li Z (2012) Abdominal obesity and the risk of colorectal adenoma: a meta-analysis of observational studies. *Eur J Cancer Prev* 21: 523–531. [PubMed: 22343656]
21. Thompson CL, Berger NA, Chak A, Li L (2012) Racial differences in measures of obesity and risk of colon adenoma. *Obesity (Silver Spring)* 20: 673–677. [PubMed: 21996657]
22. Thornton JG, Morris AM, Thornton JD, Flowers CR, McCashland TM (2007) Racial variation in colorectal polyp and tumor location. *J Natl Med Assoc* 99: 723–728. [PubMed: 17668638]
23. Ozick LA, Jacob L, Donelson SS, Agarwal SK, Freeman HP (1995) Distribution of adenomatous polyps in African-Americans. *Am J Gastroenterol* 90: 758–760. [PubMed: 7733083]
24. Jackson CS, Vega KJ (2015) Higher prevalence of proximal colon polyps and villous histology in African-Americans undergoing colonoscopy at a single equal access center. *J Gastrointest Oncol* 6: 638–643. [PubMed: 26697195]
25. Hales CM, Carroll MD, Fryar CD, Ogden CL (2017) Prevalence of obesity among adults and youth: United States, 2015–2016. 288.
26. Kabat GC, Ginsberg M, Sparano JA, Rohan TE (2016) Risk of Recurrence and Mortality in a Multi-Ethnic Breast Cancer Population. *J Racial Ethn Health Disparities*.

27. Goldman M, Druet P, Gleichmann E (1991) TH2 cells in systemic autoimmunity: insights from allogeneic diseases and chemically-induced autoimmunity. *Immunol Today* 12: 223–227. [PubMed: 1832281]
28. Giovannucci E, Ascherio A, Rimm EB, Colditz GA, Stampfer MJ, Willett WC (1995) Physical activity, obesity, and risk for colon cancer and adenoma in men. *Ann Intern Med* 122: 327–334. [PubMed: 7847643]
29. Harding JL, Shaw JE, Anstey KJ, et al. (2015) Comparison of anthropometric measures as predictors of cancer incidence: A pooled collaborative analysis of 11 Australian cohorts. *Int J Cancer* 137: 1699–1708. [PubMed: 25810218]
30. Pischon T, Lahmann PH, Boeing H, et al. (2006) Body size and risk of colon and rectal cancer in the European Prospective Investigation Into Cancer and Nutrition (EPIC). *J Natl Cancer Inst* 98: 920–931. [PubMed: 16818856]
31. Moore LL, Bradlee ML, Singer MR, et al. (2004) BMI and waist circumference as predictors of lifetime colon cancer risk in Framingham Study adults. *Int J Obes Relat Metab Disord* 28: 559–567. [PubMed: 14770200]
32. Steins Bisschop CN, van Gils CH, Emaus MJ, et al. (2014) Weight change later in life and colon and rectal cancer risk in participants in the EPIC-PANACEA study. *Am J Clin Nutr* 99: 139–147. [PubMed: 24225355]
33. Keimling M, Renehan AG, Behrens G, et al. (2013) Comparison of associations of body mass index, abdominal adiposity, and risk of colorectal cancer in a large prospective cohort study. *Cancer Epidemiol Biomarkers Prev* 22: 1383–1394. [PubMed: 23720402]
34. Renehan AG, Flood A, Adams KF, et al. (2012) Body mass index at different adult ages, weight change, and colorectal cancer risk in the National Institutes of Health-AARP Cohort. *Am J Epidemiol* 176: 1130–1140. [PubMed: 23186750]
35. Akinyemiju T, Wiener H, Pisu M (2017) Cancer-related risk factors and incidence of major cancers by race, gender and region; analysis of the NIH-AARP diet and health study. *BMC Cancer* 17: 597–017-3557–1.
36. Liu PH, Wu K, Ng K, et al. (2019) Association of Obesity With Risk of Early-Onset Colorectal Cancer Among Women. *JAMA Oncol* 5: 37–44. [PubMed: 30326010]
37. Kitahara CM, Berndt SI, de Gonzalez AB, et al. (2013) Prospective investigation of body mass index, colorectal adenoma, and colorectal cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *J Clin Oncol* 31: 2450–2459. [PubMed: 23715565]
38. Dash C, Palmer JR, Boggs DA, Rosenberg L, Adams-Campbell LL (2014) Type 2 diabetes and the risk of colorectal adenomas: Black Women’s Health Study. *Am J Epidemiol* 179: 112–119. [PubMed: 24091887]
39. Wise LA, Rosenberg L, Palmer JR, Adams-Campbell LL (2008) Anthropometric risk factors for colorectal polyps in African-American women. *Obesity (Silver Spring)* 16: 859–868. [PubMed: 18239567]
40. Giovannucci E, Rimm EB, Stampfer MJ, Colditz GA, Ascherio A, Willett WC (1994) Aspirin use and the risk for colorectal cancer and adenoma in male health professionals. *Ann Intern Med* 121: 241–246. [PubMed: 8037405]
41. Madlensky L, Daftary D, Burnett T, et al. (2007) Accuracy of colorectal polyp self-reports: findings from the colon cancer family registry. *Cancer Epidemiol Biomarkers Prev* 16: 1898–1901. [PubMed: 17726139]

Table 1.

Baseline Characteristics in the Black Women's Health Study Cohort according to BMI, 1995

Variable	<i>BMI < 18.5 (N=932)</i>		<i>BMI 18.5–24.9 (N=20917)</i>		<i>BMI 25–29.9 (N=17854)</i>		<i>BMI 30 (N=17144)</i>	
	<i>Mean (SD)</i>	<i>N(%)</i>	<i>Mean (SD)</i>	<i>N(%)</i>	<i>Mean (SD)</i>	<i>N(%)</i>	<i>Mean (SD)</i>	<i>N(%)</i>
Age, years; mean (SD)	31.76 (8.70)		36.19 (10.01)		40.36 (10.83)		40.52 (10.57)	
Education								
12 years		134 (14.41)		2972 (14.23)		3621 (20.32)		4084 (23.87)
13–15 years		364 (39.14)		7193 (34.45)		6458 (36.24)		6468 (37.80)
16 years		432 (46.45)		10715 (51.32)		7743 (43.45)		6559 (38.33)
Current smokers		161 (17.27)		3182 (15.24)		3135 (17.59)		2695 (15.74)
Alcohol (< 7 drinks/week)		47 (5.06)		1138 (5.48)		1100 (6.21)		947 (5.57)
Family history of colorectal cancer in first degree relative		34 (3.65)		1036 (4.95)		1027 (5.75)		1056 (6.16)
History of colorectal cancer screening		34 (3.65)		1269 (6.07)		1672 (9.36)		1533 (8.94)
Regular aspirin use		48 (5.15)		1424 (6.81)		1757 (9.84)		2082 (12.14)
Vigorous physical activity (< 5 hours/week)		97 (10.71)		3706 (18.29)		2342 (13.71)		1408 (8.57)
Post-menopausal		59 (6.43)		2322 (11.63)		3597 (21.74)		3636 (22.93)
Postmenopausal hormone therapy (current)		28 (3.06)		1560 (7.59)		2071 (11.84)		1889 (11.26)
Total energy intake (kcal); mean (SD)	1530.73 (678.32)		1414.27 (606.93)		1447.86 (607.60)		1571.56 (647.91)	
Red meat servings/week; mean (SD)	4.61 (5.14)		3.49 (4.11)		3.77 (4.17)		4.61 (4.86)	
Fruit and vegetable servings/week; mean (SD)	12.50 (13.66)		14.43 (13.27)		15.58 (13.69)		15.11 (13.61)	

Abbreviations: BMI, body mass index; SD, standard deviation

Table 2.

Associations of anthropometric measures with incident colorectal adenoma by age in the Black Women's Health Study (in a nested matched case-control study)¹, 1997–2011

Anthropometrics	All Participants		<50 years		50 years	
	# Cases/ controls	Multivariate OR ² (95% CI)	# Cases/ controls	Multivariate OR ² (95% CI)	# Cases/ controls	Multivariate OR ² (95% CI)
BMI, kg/m²						
Categories ³						
<18.5	3 / 12	0.56 (0.15, 2.09)	0 / 2	-	3 / 10	0.71 (0.18, 2.77)
18.5–24.9	209 / 617	1.00	74 / 187	1.00	135 / 430	1.00
25–29.9	332 / 993	1.01 (0.82, 1.24)	78 / 244	0.85 (0.58, 1.26)	254 / 749	1.09 (0.85, 1.40)
30	402 / 1200	1.06 (0.85, 1.32)	108 / 340	0.91 (0.60, 1.37)	294 / 860	1.15 (0.88, 1.50)
<i>P_{trend}</i> ⁴		0.70		0.60		0.39
Continuous						
1 SD (6.64 kg/m ²) increase ⁵	-	1.05 (0.96, 1.16)	-	0.91 (0.79, 1.05)	-	1.12 (0.99, 1.24)
<i>P, interaction by age</i> ⁶						0.08
Waist Circumference, cm						
Quintiles ³						
<76.2	164 / 474	1.00	56 / 160	1.00	108 / 314	1.00
76.2–88.8	150 / 481	0.91 (0.70, 1.19)	42 / 128	0.97 (0.60, 1.58)	108 / 353	0.92 (0.67, 1.26)
88.9–99.0	147 / 425	1.00 (0.74, 1.36)	42 / 120	0.92 (0.52, 1.63)	105 / 305	1.06 (0.67, 1.26)
99.1–109.1	143 / 453	0.96 (0.69, 1.34)	32 / 115	0.75 (0.39, 1.42)	111 / 338	1.09 (0.73, 1.61)
109.2	171 / 411	1.20 (0.84, 1.71)	36 / 107	0.89 (0.45, 1.75)	135 / 304	1.38 (0.90, 2.10)
<i>P_{trend}</i> ⁴		0.36		0.57		0.13
Continuous						
1 SD (18.23 cm) increase ⁵	-	1.09 (0.97, 1.23)	-	0.94 (0.75, 1.17)	-	1.17 (1.01, 1.35)
<i>P, interaction by age</i> ⁶						0.11
Waist to Hip Ratio						
Quintiles ³						
<0.75	142 / 471	1.00	40 / 139	1.00	102 / 332	1.00
0.75–0.82	167 / 432	1.30 (0.99, 1.71)	48 / 113	1.83 (1.09, 3.11)	119 / 319	1.21 (0.88, 1.68)
0.83–1.07	138 / 433	1.03 (0.74, 1.34)	33 / 128	1.08 (0.60, 1.96)	105 / 305	1.05 (0.72, 1.52)
1.08–1.23	168 / 456	1.18 (0.65, 1.70)	44 / 113	2.42 (0.76, 7.72)	124 / 343	0.99 (0.54, 1.81)
1.24	157 / 432	1.19 (0.58, 1.54)	41 / 123	2.20 (0.65, 7.43)	116 / 309	1.03 (0.56, 1.88)
<i>P_{trend}</i> ⁴		0.54		0.41		0.51
Continuous						
1 SD (0.25) increase ⁵	-	1.02 (0.84, 1.24)	-	0.91 (0.61, 1.34)	-	1.06 (0.84, 1.34)

Anthropometries	All Participants		<50 years		50 years	
	# Cases/ controls	Multivariate OR ² (95% CI)	# Cases/ controls	Multivariate OR ² (95% CI)	# Cases/ controls	Multivariate OR ² (95% CI)
<i>P</i> , interaction by age ⁶						0.50
Weight change since age 18, kg						
Categories ³						
<10	128 / 450	1.00	41 / 147	1.00	87 / 303	1.00
10–14	121 / 327	1.37 (1.02, 1.85)	41 / 104	1.35 (0.79, 2.29)	80 / 223	1.41(0.98, 2.04)
15–19	121 / 397	1.11 (0.83, 1.49)	36 / 91	1.34 (0.79, 2.29)	85 / 306	1.07 (0.75, 1.52)
20–24	176 / 411	1.58 (1.20, 2.09)	46 / 104	1.60 (0.96, 2.67)	130 / 307	1.65 (1.18, 2.31)
25	397 / 1,217	1.26 (1.00, 1.60)	96 / 323	1.05 (0.68, 1.63)	301 / 894	1.38 (1.03, 1.86)
<i>P</i> _{trend} ⁴		0.11		0.93		0.04
Continuous						
1 SD (15.41 kg) increase ⁵	-	1.05 (0.97, 1.14)	-	0.98 (0.86, 1.16)	-	1.08 (0.98, 1.18)
<i>P</i> , interaction by age ⁶						0.43
Body Shape Index (ABSI)						
Quintiles ³						
<0.065	132 / 446	1.00	47 / 145	1.00	85 / 301	1.00
0.065–0.071	156 / 445	1.14 (0.86, 1.50)	46 / 136	0.92 (0.55, 1.54)	110 / 309	1.45 (0.89, 2.36)
0.072–0.080	156 / 446	1.16 (1.86, 1.58)	43 / 135	1.02 (0.57, 1.83)	113 / 311	0.90 (0.65, 1.25)
0.081–0.089	154 / 445	1.26 (0.84, 1.89)	35 / 107	1.13 (0.50, 2.56)	119 / 338	0.94 (0.71, 1.26)
0.090	173 / 445	1.44 (0.95, 2.18)	36 / 103	1.40 (0.61, 3.23)	137 / 342	0.60 (0.29, 1.23)
<i>P</i> _{trend} ⁴		0.09		0.47		0.14
Continuous						
1 SD (0.01) increase ⁵	-	1.07 (0.93, 1.23)	-	1.06 (0.80, 1.42)	-	1.06 (0.90, 1.26)
<i>P</i> , interaction by age ⁶						0.93

Abbreviations: OR, odds ratio; CI, confidence interval

¹Cases and controls were matched on age and follow-up time

²Adjusted for age, education, smoking, alcohol intake, family history of CRC in a first-degree relative, NSAID use, total energy intake, red meat intake, fruit and vegetable intake, menopausal status, and physical activity.

³Based on a conditional logistic regression model with anthropometric exposures modeled as categorical variables

⁴*P*_{trend} assessed by χ^2 test for linear trend

⁵Based on a conditional logistic regression model with anthropometric exposures modeled as continuous variables

⁶Calculated using the likelihood ratio test comparing the fit of a model including the cross-product term between the anthropometric variable (e.g., BMI) and age category to a model without the cross-product term (e.g., BMI*age category)

Table 3.

Associations of anthropometric measures with incident colon adenoma by site among 50 year old women in the Black Women's Health Study (in a nested matched case-control study¹), 1997–2011

Anthropometrics	All Colon adenoma		Proximal colon adenoma	
	# Cases /controls	Multivariate OR ² (95% CI)	# Cases /controls	Multivariate OR ² (95% CI)
BMI, kg/m²				
Categories ³				
<18.5	2 / 9	0.50 (0.10, 2.45)	0 / 9	-
18.5–24.9	123 / 385	1.00	88 / 385	1.00
25–29.9	227 / 680	1.04 (0.80, 1.36)	156 / 680	1.08 (0.79, 1.47)
30	273 / 788	1.14 (0.86, 1.51)	197 / 788	1.13 (0.81, 1.58)
<i>P</i> _{trend} ⁴		0.45		0.76
Continuous				
1 SD (6.64 kg/m ²) increase ⁵	-	1.08 (0.96, 1.21)	-	1.14 (1.00, 1.31)
Waist Circumference, cm				
Quintiles ³				
<76.2	95 / 281	1.00	58 / 182	1.00
76.2–88.8	98 / 318	0.94 (0.67, 1.31)	66 / 225	0.98 (0.64, 1.49)
88.9–99.0	94 / 269	1.11 (0.75, 1.63)	70 / 184	1.39 (0.86, 2.26)
99.1–109.1	103 / 312	1.12 (0.74, 1.7)	68 / 228	1.17 (0.7, 1.98)
109.2	126 / 284	1.41 (0.9, 2.21)	100 / 206	1.92 (1.11, 3.33)
<i>P</i> _{trend} ⁴		0.13		0.03
Continuous				
1 SD (18.23 cm) increase ⁵	-	1.17 (1.00, 1.36)	-	1.35 (1.12, 1.63)
Waist to Hip Ratio				
Quintiles ³				
<0.75	92 / 296	1.00	51 / 195	1.00
0.75–0.82	107 / 283	1.2 (0.85, 1.7)	76 / 187	1.57 (1.02, 2.44)
0.83–1.07	93 / 272	1.01 (0.68, 1.49)	63 / 183	1.32 (0.8, 2.17)
1.08–1.23	114 / 319	0.97 (0.52, 1.83)	87 / 248	1.36 (0.63, 2.95)
1.24	109 / 288	1.06 (0.57, 2)	84 / 208	1.66 (0.76, 3.62)
<i>P</i> _{trend} ⁴		0.71		0.12
Continuous				
1 SD (0.25) increase ⁵	-	1.08 (0.85, 1.37)	-	1.25 (0.93, 1.68)
Weight change since age 18, kg				
Categories ³				
<10	79 / 275	1.00	50 / 191	1.00
10–14	74 / 198	1.45 (0.99, 2.13)	47 / 140	1.44 (0.89, 2.31)

Anthropometrics	All Colon adenoma		Proximal colon adenoma	
	# Cases /controls	Multivariate OR ² (95% CI)	# Cases /controls	Multivariate OR ² (95% CI)
15–19	75 / 281	1.01 (0.7, 1.47)	57 / 197	1.22 (0.78, 1.91)
20–24	116 / 280	1.61 (1.13, 2.3)	79 / 201	1.78 (1.15, 2.75)
25	278 / 814	1.39 (1.02, 1.89)	206 / 574	1.66 (1.13, 2.44)
<i>P</i> _{trend} ⁴		0.052		0.008
Continuous				
1 SD (15.41 kg) increase ⁵	-	1.07 (0.97, 1.18)	-	1.12 (1.00, 1.26)
Body Shape Index (ABSI)				
Quintiles ³				
<0.065	75 / 265	1.00	47 / 179	1.00
0.065–0.071	101 / 282	1.24 (0.87, 1.75)	66 / 196	1.3 (0.84, 2.01)
0.072–0.080	101 / 279	1.28 (0.87, 1.89)	73 / 189	1.57 (0.96, 2.55)
0.081–0.089	112 / 311	1.36 (0.83, 2.24)	81 / 223	1.59 (0.86, 2.95)
0.090	125 / 314	1.52 (0.9, 2.56)	93 / 229	1.88 (0.99, 3.56)
<i>P</i> _{trend} ⁴		0.12		0.051
Continuous				
1 SD (0.01) increase ⁵	-	1.06 (0.89, 1.27)	-	1.14 (0.91, 1.41)

Abbreviations: CRC, colorectal cancer; OR, odds ratio; CI, confidence interval

¹Cases and controls were matched on age and follow-up time

²Adjusted for age, education, smoking, alcohol intake, family history of CRC in a first-degree relative, NSAID use, total energy intake, red meat intake, fruit and vegetable intake, menopausal status, and physical activity.

³Based on a conditional logistic regression model with anthropometric exposures modeled as categorical variables

⁴*P*_{trend} assessed by χ^2 test for linear trend

⁵Based on a conditional logistic regression model with anthropometric exposures modeled as continuous variables

Table 4.

Associations of anthropometric measures with incident colorectal cancer in the Black Women's Health Study 1995–2011

Anthropometrics	All Participants		<50 years		50 years	
	# Cases / Person years	Multivariate RR <i>I</i> (95% CI)	# Cases / Person years	Multivariate RR <i>I</i> (95% CI)	# Cases / Person years	Multivariate RR <i>I</i> (95% CI)
BMI, kg/m²						
Categories ²						
<18.5	3 / 7466	1.50 (0.47, 4.77)	0 / 6338	-	3 / 1128	2.18 (0.68, 7.00)
18.5–24.9	89 / 221816	1.00	26 / 167945	1.00	63 / 53870	1.00
25–29.9	147 / 260604	1.02 (0.77, 1.35)	46 / 163148	1.46 (0.89, 2.38)	101 / 97456	0.84 (0.61, 1.16)
30	172 / 308962	0.99 (0.73, 1.33)	41 / 190988	0.97 (0.55, 1.71)	131 / 117974	0.90 (0.65, 1.26)
<i>P</i> _{trend} ³		0.98		0.65		0.89
Continuous						
1 SD (6.64 kg/m ²) increase ⁴	-	1.03 (0.90, 1.17)	-	0.88 (0.68, 1.12)	-	1.04 (0.90, 1.21)
<i>P</i> , interaction by age ⁵						0.39
Waist Circumference, cm						
Quintiles ²						
<73	53/155,688	1.00	21 / 123740	1.00	32 / 31948	1.00
73–80	49/108,504	1.07 (0.71, 1.61)	14 / 78883	0.92 (0.47, 1.82)	35 / 29621	1.11 (0.69, 1.8)
81–90	94/149,791	1.24 (0.86, 1.78)	21 / 97279	1.04 (0.56, 1.91)	73 / 52513	1.25 (0.82, 1.9)
91–103	88/156,263	1.13 (0.78, 1.64)	26 / 93404	1.28 (0.7, 2.35)	62 / 62859	1.03 (0.67, 1.6)
104	75/132,762	1.12 (0.75, 1.70)	12 / 64854	0.82 (0.37, 1.79)	63 / 67908	1.13 (0.71, 1.81)
<i>P</i> _{trend} ³		0.55		0.85		0.81
Continuous						
1 SD (18.23 cm) increase ⁴	-	1.04 (0.91, 1.18)	-	0.96 (0.75, 1.23)	-	1.02 (0.89, 1.18)
<i>P</i> , interaction by age ⁵						0.71
Waist to Hip Ratio						
Quintiles ²						
<0.72	80/136,281	1.00	24 / 97157	1.00	56 / 39124	1.00
0.73–0.78	54/138,724	0.56 (0.39, 0.81)	11 / 96567	0.45 (0.22, 0.92)	43 / 42157	0.68 (0.46, 1.01)
0.79–0.85	78/135,689	0.87 (0.63, 1.20)	20 / 93631	0.83 (0.46, 1.51)	58 / 42058	0.91 (0.63, 1.32)
0.86–1.10	71/139,970	0.77 (0.55, 1.09)	23 / 93372	0.96 (0.53, 1.75)	48 / 46598	0.77 (0.52, 1.14)
1.11	71/136,856	0.72 (0.47, 1.08)	14 / 65698	0.7 (0.3, 1.64)	57 / 71158	0.75 (0.47, 1.19)
<i>P</i> _{trend} ³		0.35		0.99		0.37

Anthropometrics	All Participants		<50 years		50 years	
	# Cases / Person years	Multivariate RR <i>I</i> (95% CI)	# Cases / Person years	Multivariate RR <i>I</i> (95% CI)	# Cases / Person years	Multivariate RR <i>I</i> (95% CI)
Continuous						
1 SD (0.25) increase ⁴	-	0.91 (0.79, 1.05)	-	0.91 (0.68, 1.23)	-	0.92 (0.79, 1.08)
<i>P</i> , interaction by age ⁵						0.16
Weight change since age 18, kg						
Categories ²						
<10	62/183,212	1.00	27 / 146155	1.00	35 / 37057	1.00
10–14	50/108338	1.11 (0.75, 1.65)	8 / 77856	0.47 (0.21, 1.03)	42 / 30482	1.51 (0.96, 2.38)
15–19	71/110,772	1.37 (0.95, 1.96)	26 / 72932	1.48 (0.85, 2.56)	45 / 37840	1.31 (0.83, 2.05)
20–24	61/115,724	0.96 (0.65, 1.41)	16 / 72117	0.86 (0.46, 1.61)	45 / 43607	1.16 (0.74, 1.82)
25	159/275,918	1.11 (0.81, 1.53)	34 / 156667	0.77 (0.45, 1.31)	125 / 119252	1.25 (0.84, 1.85)
<i>P</i> _{trend} ³		0.88		0.29		0.80
Continuous						
1 SD (15.41 kg) increase ⁴	-	1.02 (0.92, 1.14)	-	0.89 (0.73, 1.09)	-	1.04 (0.92, 1.18)
<i>P</i> , interaction by age ⁵						0.61
Body Shape Index (ABSI)						
Quintiles ²						
<0.065	73/139,890	1.00	18 / 97488	1.00	55 / 42402	1.00
0.065–0.068	67/139,682	1.02 (0.72, 1.43)	23 / 102391	1.26 (0.68, 2.33)	44 / 37291	0.91 (0.61, 1.35)
0.069–0.073	64/139,893	0.87 (0.61, 1.25)	17 / 100549	0.93 (0.48, 1.8)	47 / 39344	0.92 (0.62, 1.36)
0.074–0.082	77/139,927	0.94 (0.67, 1.33)	23 / 89468	1.32 (0.7, 2.47)	54 / 50459	0.85 (0.58, 1.24)
0.083	76/139,831	0.84 (0.57, 1.25)	13 / 66162	0.89 (0.39, 2.02)	63 / 73669	0.84 (0.55, 1.29)
<i>P</i> _{trend} ³		0.36		0.99		0.37
Continuous						
1 SD (0.01) increase ⁴	-	0.92 (0.81, 1.04)	-	0.96 (0.74, 1.23)	-	0.92 (0.80, 1.05)
<i>P</i> , interaction by age ⁵						0.23

Abbreviations: CRC, colorectal cancer; NSAID, nonsteroidal anti-inflammatory drug; HRT, hormone replacement therapy; RR, relative risk; CI, confidence interval

¹Adjusted for age, education, smoking, alcohol intake, family history of CRC in a first-degree relative, CRC screening, NSAID use, total energy intake, red meat intake, fruit and vegetable intake, menopausal status, and physical activity.

²Based on a Cox proportional regression model with anthropometric exposures modeled as categorical variables

³*P*_{trend} assessed by χ^2 test for linear trend

⁴Based on a Cox proportional regression model with anthropometric exposures modeled as continuous variables

⁵Calculated using the likelihood ratio test comparing the fit of a model including the cross-product term between the anthropometric variable (e.g., BMI) and age category to a model without this term (e.g., BMI*age category)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript