



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ELSEVIER

Available online at www.sciencedirect.com

Journal of Hospital Infection

journal homepage: www.elsevier.com/locate/jhin

Practice points

Minimizing intra-hospital transmission of COVID-19: the role of social distancing

L.E. Wee^{a,b,*}, E.P. Conceicao^c, X.Y.J. Sim^{b,c}, M.K. Aung^c, K.Y. Tan^c,
H.M. Wong^{b,c}, L. Wijaya^b, B.H. Tan^b, M.L. Ling^c, I. Venkatachalam^{b,c}

^a Singhealth Infectious Diseases Residency, Singapore

^b Department of Infectious Diseases, Singapore General Hospital, Singapore

^c Department of Infection Prevention and Epidemiology, Singapore General Hospital, Singapore

ARTICLE INFO

Article history:

Received 6 April 2020

Accepted 8 April 2020

Available online 12 April 2020

Keywords:

Social distancing

COVID-19

Hospital

Nosocomial



In the ongoing COVID-19 pandemic caused by the novel coronavirus, SARS CoV-2, early isolation of hospitalized inpatients with suspected COVID-19 is important to reduce the likelihood of nosocomial spread. However, patients with COVID-19 may present with respiratory syndromes indistinguishable from those caused by common respiratory viruses [1]. This poses a challenge for early isolation and containment, especially during significant ongoing community transmission. While isolation ward beds are prioritized for suspected COVID-19 cases, unsuspected cases of COVID-19 without suspicious contact or travel history may initially be nursed outside of dedicated isolation wards prior to detection. Given that

patients outside the isolation ward may not be subject to movement restrictions and share common facilities, social mingling represents a potential route for nosocomial spread, especially as COVID-19 cases may present with mild symptoms and remain relatively well [1]. While social distancing has been identified as crucial for containment in the community [2], social distancing within hospitals is equally vital in reducing nosocomial spread, especially in hospitals where the majority of patients are nursed in multi-bedded cohort rooms, rather than in single-occupancy rooms.

In Singapore, a globalized Asian city-state, the first imported case of COVID-19 was reported at the end of January 2020; followed by the first case of local transmission in early February 2020 [3]. At our institution, the Singapore General Hospital (SGH), the isolation ward was reserved for confirmed/suspected cases of COVID-19. However, given rising numbers of locally transmitted cases, from 4th February, our institution placed individuals admitting with respiratory symptoms but without suspicious contact or travel history in respiratory surveillance wards (RSWs) where COVID-19 was first excluded and healthcare workers (HCWs) used full personal protective equipment (PPE) including N95 masks, disposable gowns, gloves and face-shields. Despite this resource-intensive containment effort, it was recognized that some cases of COVID-19 with mild symptoms might be initially admitted to the general ward. Our institution therefore emphasized hospital-wide social distancing measures. For patients admitting to the RSW, as the risk of a potentially unsuspected case of COVID-19 was higher, patients were advised to avoid mingling and to wear surgical masks at all times; no visitors were allowed. Additionally, infrastructural modifications were instituted to facilitate social distancing. In the RSWs, patients were nursed in cohort rooms with three patients to a room, spaced at least ~2 m apart, and partitions were placed between patient beds (Figure 1). In the general ward, shared communal

* Corresponding author. Address: Singhealth Infectious Diseases Residency, Singapore. Tel.: +65 96777651.

E-mail address: ian.wee@mohh.com.sg (L.E. Wee).




Figure 1. Comparison of ward layout and social distancing measures employed in a general ward and a respiratory surveillance ward, during the COVID-19 outbreak.

facilities (e.g., day rooms) were closed during the duration of the ongoing COVID-19 outbreak, and patients were limited to one visitor at any time. HCWs in the general ward wore surgical masks. Hospital-wide, in common areas such as waiting areas, pharmacies, food and retail outlets, patients were directed to keep 1 m apart from one another, using visual cues (e.g., floor markings and markings on seats) to guide waiting and queuing in both seated and standing areas.

Over a three-month period from 4th January to 4th April 2020, a total of 75 confirmed cases of COVID-19 were diagnosed in our institution. While the majority of cases (84.0%, 63/75) were admitted to isolation wards, 12 cases of COVID-19 were initially admitted outside of the isolation ward. Of these, the majority (91.6%, 11/12) were admitted to the RSW. One patient was initially admitted to the general ward and nursed in a cohorted cubicle with five other patients, as respiratory symptoms were initially mild. The patient was transferred to an RSW 19 h after admission, where the diagnosis of COVID-19 was made. At diagnosis, the cycle threshold (Ct) value for SARS-CoV-2 on reverse transcription-polymerase chain reaction (RT-PCR) testing of oropharyngeal swab samples was 18, an inverse surrogate for high viral load and potential infectivity; this was in keeping with data suggesting peak viral shedding in the first week of symptoms [4]. A total of 18 patients in the general ward and two patients in the RSW had shared a room or common toilet with the index case; all were deemed to be exposed (Figure 1), given potential contamination of the shared air and surface environment from droplet and fomite spread [5]. A total of eight HCWs in the general ward had cared for the patient while wearing surgical masks. However, none of the

exposed patients or HCWs developed COVID-19 within the estimated incubation period [6], despite being closely followed up for 14 days. Of note, the patient had complied with social distancing measures and had not interacted with any of the other exposed patients. At the patient’s initiative, he had worn a mask throughout the admission as an added precaution to minimize infection.

Minimizing nosocomial transmission of COVID-19 remains a challenge, given the wide spectrum of respiratory syndromes and mild respiratory symptoms at presentation [1]. Influencing patient behaviour to reduce the risk of patient-to-patient spread remains crucial. Social distancing between inpatients is important during an ongoing outbreak and should be reinforced in higher-risk areas.

Acknowledgements

The authors thank their colleagues for the unstinting efforts against COVID-19.

Conflict of interest statement

None declared.

Funding sources

None.

References

[1] Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*

2020. <https://doi.org/10.1056/NEJMoa2002032> [Epub ahead of print].
- [2] Wilder-Smith A, Freedman DO. Isolation, quarantine, social distancing and community containment: pivotal role for old-style public health measures in the novel coronavirus (2019-nCoV) outbreak. *J Travel Med* 2020;27. pii: taaa020.
- [3] Wong JEL, Leo YS, Tan CC. COVID-19 in Singapore – current experience: critical global issues that require attention and action. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.2467> [Epub ahead of print].
- [4] Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020. <https://doi.org/10.1038/s41586-020-2196-x> [Epub ahead of print].
- [5] Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from a symptomatic patient. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.3227> [Epub ahead of print].
- [6] Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, et al. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann Intern Med* 2020. <https://doi.org/10.7326/M20-0504> [Epub ahead of print].