

Total Synthesis

International Edition: DOI: 10.1002/anie.201914620
German Edition: DOI: 10.1002/ange.201914620A Convergent Total Synthesis of the Death Cap Toxin α -AmanitinMary-Ann J. Siegert[†], Caroline H. Knittel[†], and Roderich D. Süssmuth^{*}

In memory of Gerhard Jas

Abstract: The toxic bicyclic octapeptide α -amanitin is mostly found in different species of the mushroom genus *Amanita*, with the death cap (*Amanita phalloides*) as one of the most prominent members. Due to its high selective inhibition of RNA polymerase II, which is directly linked to its high toxicity, particularly to hepatocytes, α -amanitin received an increased attention as a toxin-component of antibody-drug conjugates (ADC) in cancer research. Furthermore, the isolation of α -amanitin from mushrooms as the sole source severely restricts compound supply as well as further investigations, as structure–activity relationship (SAR) studies. Based on a straightforward access to the non-proteinogenic amino acid dihydroxyisoleucine, we herein present a robust total synthesis of α -amanitin providing options for production at larger scale as well as future structural diversifications.

Amatoxins are ribosomally synthesized toxic bicyclic octapeptides mainly found in different species of the fungus genus *Amanita*, most notably *Amanita phalloides*.^[1–4] These compounds are selective inhibitors of eukaryotic RNA polymerase II [α -amanitin, $K_d = 10^{-9}$ M], which ultimately leads to apoptosis.^[4,5] The fungal toxins were first isolated by Heinrich Wieland and Rudolf Hallermayer in 1941 and their structure was elucidated in the 1950s and 1960s (Wieland et al.).^[6,7] All amatoxins contain eight L-configured amino acids and a tryptathionine linkage between the side chains of tryptophan and cysteine, with an (*R*)-configured sulfoxide moiety.^[1,3] However, the amanitins differ in their hydroxylation pattern of the amino acid side chains, which modulates their toxicity. The effect of the differently hydroxylated amino acids on the amatoxin toxicity was investigated in various studies.^[8–11]

In the past 30 years several approaches were made to synthesize amatoxin derivatives.^[8,12–15] Zanotti et al. (1987) described the first synthesis of an amaninamide derivative

starting from the linear octapeptide followed by thioether formation using the Savige-Fontana reaction and subsequent macrocyclization.^[8] The Savige-Fontana methodology is based on an N-terminal Hpi (3*a*-hydroxypyrrolo[2,3-*b*]indole) moiety and a trityl side-chain protected cysteine residue. Simultaneous deprotection of the cysteine residue and activation of the Hpi moiety with trifluoroacetic acid (Tfa) leads to the formation of the thioether.^[14,16] Other routes to tryptathionine involved the reaction of tryptophan with a cysteine-sulfonyl iodide or chloride. Until now, these methodologies were only used for the synthesis of phalloidin and its derivatives.^[17–19]

The most intensively studied amatoxin is α -amanitin (**1**) (Figure 1). It contains the hydroxylated amino acids 6-hydroxytryptophan (Htp), *trans*-4-hydroxyproline (Hyp) and




Figure 1. Structure of α -Amanitin (**1**). The numbers in blue, turquoise and red refer to the number of amino acids of the building blocks (depicted blue, turquoise and red) employed in this total synthesis strategy [5+1+2].

(3*R*,4*R*)-4,5-dihydroxyisoleucine (Dhil).^[1,2] Due to its small molecular size, good solubility in aqueous buffers and its inhibition of RNA polymerase II, it received an increased attention as an antibody-drug conjugate (ADC) in cancer research in the past years.^[14,20,21] In a preclinical study in 2011 conducted by the German Cancer Research Center in Heidelberg (Germany) naturally occurring α -amanitin was conjugated with the chimerized anti-EpCAM monoclonal antibody chiHEA125 to form the antibody-drug conjugate chiHEA125-Ama.^[22] This therapeutic ADC targets the human epithelial adhesion molecule (EpCAM), which is overexpressed in the majority of cancer cells. The study

[*] M.-A. J. Siegert,^[†] C. H. Knittel,^[†] Prof. Dr. R. D. Süssmuth
Institute of Chemistry, Technische Universität Berlin
Straße des 17. Juni 124, 10623 Berlin (Germany)
E-mail: suessmuth@chem.tu-berlin.de

[†] These authors contributed equally to this work.


Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.201914620>.

© 2019 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial, and no modifications or adaptations are made.

showed that chiHEA125-Ama has a potent antitumor activity against pancreatic cancer cells. Compared with chiHEA125 two injections of chiHEA125-Ama with a dosage of 50 or 100 $\mu\text{g kg}^{-1}$ (in relation to α -amanitin), showed significant tumor regression in mice.^[22] Based on its promising properties for drug development there is an increasing demand for α -amanitin, which cannot be satisfied by biotechnological methods or from nature. Only last year the first total synthesis of α -amanitin has been accomplished by Perrin and co-workers, applying a sophisticated Savige-Fontana methodology as a key step for the formation of the tryptathionine bridge.^[23]

Herein, we report an alternative and robust total synthesis of α -amanitin with the aim to be convergent and scalable in solution phase synthesis. Therefore, we applied a peptide fragment based methodology and developed straightforward syntheses to the central Dhil and Htp building blocks.


Scheme 1 shows the general retrosynthetic strategy for the total synthesis of α -amanitin (**1**) referred to as [5+1+2]-strategy. Sulfide oxidation of the tryptathionine linkage was envisioned to be introduced close to the final stages of the synthesis route, when the bicyclic structure was established by macrocyclization between Dhil and Hyp of monocyclic octapeptide **2**. This amanitin precursor should be accessible by stepwise couplings of the monocyclic building block **5** with a Dhil derivative **6** and an Asn-Hyp dipeptide **4**. It was important for reasons of scalability that both, the Dhil derivative **6** and the monocyclic tryptathionine-containing



Scheme 1. Retrosynthetic analysis of α -amanitin (**1**). Final assembly of α -amanitin from three building blocks and enantioselective synthesis of the hydroxylated amino acid building blocks.

building block **5** could be synthesized on a multi-gram scale in solution phase. In our synthesis strategy, **5** is synthesized by preformation of the pseudo-orthogonally protected tryptathionine linkage **7** between a 6-hydroxytryptophan (Htp) derivative (**8**) and cysteine.

The synthesis of 6-hydroxytryptophan was not trivial and finally we decided for a dynamic kinetic resolution (DKR) route employing a chiral tridentate ligand.^[24,25] Starting with 6-benzyloxyindol (**9**), alkylation using L-serine, Ac₂O and AcOH (Scheme 2) was performed, which rendered racemic tryptophan **13**.^[26] After removal of the acetyl group the




Scheme 2. Synthesis of *N*-Teoc-6-benzyloxy-L-tryptophan (**8**) by dynamic kinetic resolution: a) L-serine, Ac₂O, AcOH, 75 °C, 2 h, 82%; b) 40% NaOH, MeOH/H₂O, c) chiral ligand **16**, Ni(NO₃)₂·6H₂O, MeOH, reflux, 16 h, 84% over two steps, d) 6 M HCl, MeOH, 70 °C, 2 h, e) TeocOSu, Et₃N, DMF, 60 °C, 4 d, quant.

racemate was treated with the chiral ligand **16** and Ni(NO₃)₂·6H₂O under basic conditions, which led to nickel(II) complex **14** with a diastereomeric ratio (*dr*) of 99% (see Supporting Information). Disassembly of the nickel(II) complex **14** under acidic conditions resulted in enantiomerically pure 6-benzyloxy-L-tryptophan (**15**). Finally, the free amino acid was trapped with TeocOSu, which provided *N*-Teoc-6-benzyloxy-L-tryptophan (**8**).

With 6-hydroxytryptophan derivative **8** in hand, the tryptathionine linkage was readily established by treatment of a fully protected L-cysteine derivative (**18**) with sulfur chloride (SO₂Cl₂) (Scheme 3).^[17] In the course of the reaction, cleavage of the disulfide afforded the highly reactive sulfenyl chloride monomer as an intermediate, which underwent a S_EAr reaction with Htp (**8**). We realized that protection of the C-terminus of **8** was not necessary for quantitative tryptathionine formation and that slow titration with the sulfenyl chloride solution prevents cysteine double substitution at Htp. Coupling of OSu-ester preactivated tryptathionine **7** with a H-Gly-Ile-Gly-OH tripeptide (see Supporting Information) afforded linear pentapeptide **19**.


After Boc-deprotection with *para*-toluenesulfonic acid (PTSA), macrocyclization with propanephosphonic acid anhydride (T3P) was performed. Both Teoc and *t*Bu protecting groups could be simultaneously cleaved under strong



Scheme 3. Synthesis of the C- and N-terminally deprotected monocyclic tryptathionine peptide **5**: a) Boc_2O , NaHCO_3 , dioxane/ H_2O , r.t., 16 h, 99%; b) SO_2Cl_2 , CHCl_3 , r.t., 1 h; c) **8**, NaHCO_3 , CHCl_3 , 0°C to r.t., quant.; d) H-Gly-Ile-Gly-OH, $\text{CO}(\text{OSu})_2$, collidine, acetonitrile/ H_2O , r.t., 2 h; e) PTSA, THF, r.t., 2 h, 70% over 2 steps; f) T3P, DIPEA, DMF, r.t., 16 h, 70%, g) TFA/ H_2O (7:3), r.t., 2 h, quant.

acidic conditions using TFA, rendering monocyclic building block **5** with no decomposition of the thioether observed.

A particular synthetic challenge was the (3*R*,4*R*)-4,5-*L*-dihydroxyisoleucine derivative **6**, which was finally accomplished in seven steps starting from glycine *tert*-butyl ester (**23**). Quantitative trifluoroacetylation yielded Tfa-Gly-*O**t*Bu (**12**) followed by ruthenium-catalyzed asymmetric allylic alkylation with **22**, in analogy to a method of Kazmaier et al.^[27,28] While the original protocol of asymmetric alkylation employed (*S*)-3-(benzyloxy)-1-butene as alkylating agent, we used a chiral allylic carbonate (**22**), which was synthesized by Sharpless epoxidation of (*E*)-crotyl alcohol **20** followed by reductive elimination promoted by a zinc-copper couple.^[29,30] The chirality transfer promoted by allylic carbonate **22** mainly led to the *anti*-directed formation of a fully protected 4,5-didehydroisoleucine derivative **11** with a diastereomeric ratio (*dr* = 86:14) and an enantiomeric excess (*ee* = 98%), superior to initial original reaction conditions (see Supporting Information). The amino group was deprotected by reduction of the trifluoroacetyl group with sodium borohydride^[31] and refurbished with the Fmoc-protecting group to give **25**. The C-terminal *t*Bu protecting group proved to be essential preventing lactone formation, the main threat during dihydroxylation of the fully protected didehydroisoleucine derivative **25**. The AD reaction was performed by a simple Upjohn dihydroxylation with potassium osmate and NMO. The stereoselectivity of the reaction was controllable solely by the applied solvent mixture. While dihydroxylation in a biphasic system of $\text{H}_2\text{O}/\text{CHCl}_3$ led to the formation of mainly the desired (2*S*,3*R*,4*R*)-diastereomer **10** in a 2.5:1 ratio, the undesired (2*S*,3*R*,4*S*)-diastereomer was mainly formed in a mixture of $\text{H}_2\text{O}/t\text{BuOH}$. Surprisingly, adding (DHQD)₂PHAL ligand to the reaction mixture led to an impairment of the stereoselectivity towards the wrong enantiomer. Finally, the hydroxy groups of the side chain of **10** were protected with the *tert*-butyldimethylsilyl (TBS) protecting group, allowing for the mild orthogonal cleavage of




Scheme 4. Synthesis of side chain-protected Fmoc-(3*R*,4*R*)-4,5-*L*-dihydroxyisoleucine (**6**) a) (+)-DIPT, $\text{Ti}(\text{O}i\text{Pr})_4$, TBHP, DCM, -20°C , 4 h; b) TsCl, Et_3N , DMAP, DCM, -10°C , 30 h, 67%; c) NaI, Zn(Cu), THF, 70°C , 2 h; d) Boc_2O , NaH, THF, 0°C to r.t., 16 h, 75%; e) ethyl trifluoroacetate, NET_3 , MeOH, r.t., 16 h, quant.; f) LHMDS, ZnCl_2 , PPh_3 , [(*p*-cymene) RuCl_2]₂, **22**, THF, -72°C to r.t., 16 h, 88%; g) NaBH_4 , MeOH, r.t., 1 h; h) FmocOSu, Et_3N , dioxane, r.t., 4 h, 82%; i) $\text{K}_2\text{OsO}_4 \cdot \text{H}_2\text{O}$, NMO, $\text{CHCl}_3/\text{H}_2\text{O}$, r.t., 6 h, 40%; j) TBSCl, pyridine/DMF(1:9), r.t., 24 h, 95%; k) TMSOTf, 2,6-lutidine, 0°C to r.t., 2 h, 90%.

*t*Bu in nearly quantitative yield in presence of an excess of TMSOTf, which resulted in the formation of Dhil building block **6** (Scheme 4).

Parallel to the synthesis of Dhil and the monocyclic tryptathionine building block, C-terminally 9-fluorenylmethyl ester (Fm)-protected dipeptide **4** was synthesized in solution phase (see Supporting Information) serving as the third building block for the [5+1+2]-assembly. First, Dhil derivative **6** was coupled to the N- and C-terminally unprotected monocyclic pentapeptide **5** applying a protocol activating the carboxylic function of Dhil as an active ester and concomitantly increasing the nucleophilicity of the amino group of **5** by silylation with *N*-methyl-*N*-trimethylsilylacetylamine (MSA) (Scheme 5).^[32,33] This way no oligomerized side-products were formed. The resulting monocyclic hexapeptide **3** could then be coupled to the previously synthesized C-terminally Fm-protected dipeptide **4** in a one pot reaction. The final cyclization was performed after simultaneous Fmoc- and Fm-cleavage, followed by TBS deprotection with TBAF yielding monocyclic octapeptide **2**. After bicyclization using HATU to **26**, the benzyl protecting group was cleaved from the Htp side-chain of **26** applying a hard acid and soft nucleophile system.^[34] Finally, the oxidation to the sulfoxide was performed using *m*CPBA according to Perrin et al.^[23] affording the target molecule α -amanitin (**1**). The oxidation step resulted to be necessary after benzyl-deprotection and not vice versa, because of sulfoxide instability under strongly acidic conditions.

The final characterization of synthetic α -amanitin was performed by CD spectroscopy. The recorded spectrum coincided well with that of the natural sample (Figure 2). The ¹H- and 2D-NMR-data and the co-injection on RP-C18 chromatography (see Supporting Information) of the synthetic α -amanitin were in accordance with the data of the natural one, proving undoubtedly that the α -amanitin synthesis was successful.

In conclusion, our herein presented synthetic strategy towards α -amanitin is clearly different from the ground



Scheme 5. Fragment couplings and cyclizations to the target molecule α -Amanitin (1): a) MSA, COMU, DIPEA, DMA, 0°C to r.t. 3 h; b) (a and b: onepot), H-Asn-Hyp-OFm-HCl (4), HATU, DIPEA, DMF, 0°C to r.t., 2 h; c) Et₂NH, DMF, 1 h, r.t.; d) 1 M TBAF, THF, r.t., 2 h, 77% over four steps; e) HATU, DIPEA, DMF, r.t., 16 h, 68%, f) BF₃·OEt₂, EtSH, r.t., 2 h; g) mCPBA, iPrOH/EtOH (2:1), r.t., 30 min, 35% over two steps.




Figure 2. CD spectra of natural and synthetic α -amanitin (1).

breaking work of Perrin et al.^[23] in several aspects: with regard to the amino acid building blocks, we developed an independent access to 6-OH-Htp and Dhil, both non-trivial enterprises. For the latter amino acid, we could rely on methodologies developed by Kazmaier et al. which we could engineer into a seven-step synthesis with high enantiomeric excess, which so far is the shortest access to this type of amino acid. Unlike the work of Perrin et al., we employed a [5+1+2]-strategy under preformation of the thioether bond thus rendering a (pseudo-orthogonally) protected

tryptathionine. Finally, the synthesis constitutes the first α -amanitin synthesis fully performed in solution phase. We believe that the convergent and robust synthesis, including the industrially exploitable syntheses of the amino acids Htp and Dhil, will be valuable for future therapeutic purposes in cancer therapy, when larger amounts of α -amanitin are required.

Acknowledgements

We thank Dr. Christian Ewers and Dr. Gerhard Jas for support of the project. Furthermore, the work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation RTG 2473 “Bioactive Peptides”, project number 392923329).

Conflict of interest

The authors declare no conflict of interest.

Keywords: amatoxins · asymmetric synthesis · total synthesis · tryptathionine · α -amanitin

How to cite: *Angew. Chem. Int. Ed.* **2020**, *59*, 5500–5504
Angew. Chem. **2020**, *132*, 5542–5546

- [1] T. Wieland, H. Faulstich, *Crit. Rev. Biochem.* **1978**, *5*, 185–260.
- [2] T. Wieland, *Int. J. Pept. Protein Res.* **1983**, *22*, 257–276.
- [3] T. Wienland, H. Faulstich, *Experientia* **1991**, *47*, 1186–1193.
- [4] J. D. Walton, H. E. Hallen-Adams, H. Luo, *Biopolymers* **2010**, *94*, 659–664.
- [5] M. Cochet-Meilhac, P. Chambon, *Biochim. Biophys. Acta Nucleic Acids Protein Synth.* **1974**, *353*, 160–184.
- [6] T. Wieland, G. Schmidt, *Justus Liebigs Ann. Chem.* **1952**, *577*, 215–233.
- [7] T. Wieland, U. Gebert, *Justus Liebigs Ann. Chem.* **1966**, *700*, 157–173.
- [8] G. Zanotti, C. Möhringer, T. Wieland, *Int. J. Peptide Protein Res.* **1987**, *30*, 450–459.
- [9] G. Zanotti, G. Petersen, *Int. J. Pept. Res.* **1992**, *551*–558.
- [10] G. Shoham, W. N. Lipscomb, T. Wieland, *J. Am. Chem. Soc.* **1989**, *111*, 4791–4809.
- [11] A. Buku, T. Wieland, H. Bodenmüller, H. Faulstich, *Experientia* **1980**, *36*, 33–34.
- [12] G. Zanotti, G. D'auria, L. Paolillo, E. Trivellone, *Int. J. Pept. Protein Res.* **1988**, *32*, 9–20.
- [13] G. Zanotti, T. Wieland, G. Dauria, L. Paolillo, E. Trivellone, *Int. J. Pept. Protein Res.* **1990**, *35*, 263–270.
- [14] J. P. May, D. M. Perrin, *Chem. Eur. J.* **2008**, *14*, 3404–3409.
- [15] L. Zhao, J. P. May, A. Blanc, D. J. Dietrich, A. Loonchanta, K. Matinkhoo, A. Pryyma, D. M. Perrin, *ChemBioChem* **2015**, *16*, 1420–1425.
- [16] J. P. May, P. Fournier, J. Pellicelli, B. O. Patrick, D. M. Perrin, *J. Org. Chem.* **2005**, *70*, 8424–8430.
- [17] M. O. Anderson, A. A. Shelat, R. K. Guy, *J. Org. Chem.* **2005**, *70*, 4578–4584.
- [18] L. A. Schuresko, R. S. Lokey, *Angew. Chem. Int. Ed.* **2007**, *46*, 3547–3549; *Angew. Chem.* **2007**, *119*, 3617–3619.
- [19] G. Yao, J.-O. Joswig, B. G. Keller, R. D. Süssmuth, *Chem. Eur. J.* **2019**, *25*, 8030–8034.
- [20] A. Pahl, C. Lutz, T. Hechler, *Drug Discovery Today Technol.* **2018**, *30*, 85–89.

- [21] J. Anderl, H. Faulstich, T. Hechler, M. Kulke, in *Antib.-Drug Conjug.* (Ed.: L. Ducry), Humana Press, Totowa, NJ, **2013**, pp. 51–70.
- [22] G. Moldenhauer, A. V. Salnikov, S. Lüttgau, I. Herr, J. Anderl, H. Faulstich, *J. Natl. Cancer Inst.* **2012**, *104*, 622–634.
- [23] K. Matinkhoo, A. Pryma, M. Todorovic, B. O. Patrick, D. M. Perrin, *J. Am. Chem. Soc.* **2018**, *140*, 6513–6517.
- [24] S. Zhou, J. Wang, X. Chen, J. L. Aceña, V. A. Soloshonok, H. Liu, *Angew. Chem. Int. Ed.* **2014**, *53*, 7883–7886; *Angew. Chem.* **2014**, *126*, 8017–8020.
- [25] Y. Nian, J. Wang, S. Zhou, S. Wang, H. Moriwaki, A. Kawashima, V. A. Soloshonok, H. Liu, *Angew. Chem. Int. Ed.* **2015**, *54*, 12918–12922; *Angew. Chem.* **2015**, *127*, 13110–13114.
- [26] G. Blaser, J. M. Sanderson, A. S. Batsanov, J. A. K. Howard, *Tetrahedron Lett.* **2008**, *49*, 2795–2798.
- [27] A. Bayer, U. Kazmaier, *Chem. Eur. J.* **2014**, *20*, 10484–10491.
- [28] A. Bayer, U. Kazmaier, *Org. Lett.* **2010**, *12*, 4960–4963.
- [29] E. Balmer, A. Germain, W. P. Jackson, B. Lygo, *J. Chem. Soc. Perkin Trans. 1* **1993**, 399–400.
- [30] Y. Gao, J. M. Klunder, R. M. Hanson, H. Masamune, S. Y. Ko, K. B. Sharpless, *J. Am. Chem. Soc.* **1987**, *109*, 5765–5780.
- [31] F. Weygand, E. Frauendorfer, *Chem. Ber.* **1970**, *103*, 2437–2449.
- [32] V. V. Suresh Babu, G.-R. Vasanthakumar, S. J. Tantry, *Tetrahedron Lett.* **2005**, *46*, 4099–4102.
- [33] *Process for the Manufacture of Persilylated Peptides*, R. Callens, T. Delplanche, EP2060580A1, **2009**.
- [34] K. Fuji, K. Ichikawa, M. Node, E. Fujita, *J. Org. Chem.* **1979**, *44*, 1661–1664.

Manuscript received: November 15, 2019

Accepted manuscript online: December 17, 2019

Version of record online: February 6, 2020