



Published in final edited form as:

Subst Abus. 2019 ; 40(1): 43–51. doi:10.1080/08897077.2018.1475319.

Motivational interviewing training of substance use treatment professionals: A systematic review

Michael B. Madson, PhD^a, Margo C. Villarosa-Hurlocker, PhD^{a,b}, Julie A. Schumacher, PhD^{c,d}, Daniel C. Williams, PhD^c, Jami M. Gauthier, PhD^c

^aDepartment of Psychology, The University of Southern Mississippi, Hattiesburg, Mississippi, USA;

^bCenter on Alcoholism, Substance Abuse, and Addictions, Albuquerque, New Mexico, USA;

^cUniversity of Mississippi Medical Center, Jackson, Mississippi, USA;

^dPrevention Research Institute, Lexington, Kentucky, USA

Abstract

Background: Through evaluations of training programs, systematic reviews, and meta-analyses, advances in identifying best practices for disseminating motivational interviewing (MI) have emerged. To advance this work further, inclusion of thorough descriptions of the following is needed in research publications: study (design, trainee characteristics, setting characteristics), training and coaching methods (if applicable), trainer qualifications, and evaluation of MI skills.

Methods: The purpose of this study was to systematically evaluate the research on MI training of substance use treatment professionals for the inclusion of such descriptions. Twenty-five studies were reviewed using a scoring rubric developed by the authors.

Results: Just over two thirds of the studies (68%) were randomized controlled trials of MI training. The majority of studies provided information about (a) trainee characteristics (professional background =76%, education =60%, experience =56%); (b) setting characteristics training (80%); (c) training methods (format =96%, length =92%); (d) coaching (76%); and (e) evaluation of MI skills (92%).

Conclusion: Findings suggest advancements in MI training studies since previous reviews, especially in regards to the inclusion of feedback and coaching. However, this review also found that inconsistencies in methods and reporting of training characteristics, as well as limited follow-up assessment of trainees' skill, continue to limit knowledge of effective training methods.

Keywords

Motivational interviewing; systematic review; training

CONTACT Michael B. Madson, PhD Michael.madson@usm.edu Department of Psychology, The University of Southern Mississippi, 118 College Drive #5025, Hattiesburg, MS 39406-5025, USA.

Author contributions

Drs. Madson and Villarosa-Hurlocker conceptualized the study contributed data collection, analysis and manuscript development and approval of the final manuscript. Dr. Schumacher contributed to study methodology, manuscript development and revision, and approved the final document. Dr. Williams provided critical feedback on the manuscript and approved the final version. Dr. Gauthier contributed to data analysis, manuscript development and approved the final version.

Introduction

There has been a growing literature focused on motivational interviewing (MI) training, with an emphasis on the evaluation of different methods. The description of MI as a simple concept comprising a complex skill set¹ underscores the importance of training guidelines for effective treatment delivery and quality implementation. Early research evaluating MI training methods showed significant changes in clinician behaviors from training,² yet those changes were not reflected in client outcomes. In 2009, Madson et al. systematically reviewed the MI training literature, finding that although MI trainings were generally effective, they often relied on workshop/seminar formats, which incorporated limited experiential activities, and there were very little coaching or feedback to build MI competency.³ Based on their findings, Madson and colleagues recommended that best practices for reporting MI training and evaluation should include standardized training procedures, an outline of effective ingredients, incorporation of evidence-based measures of competency, and a method for linking training to client outcomes.³ Subsequently, 2 systematic reviews have supported these recommendations.^{4,5} Toward that end, the purpose of the current study was to systematically review published research on MI training with substance use (SU) treatment professionals.

Potential benchmarks for best practices related to the duration and frequency of MI training recently emerged from 2 meta-analytic studies. First, de Roten et al. identified 12–16 hours as an adequate dose of MI workshop training for providers to achieve gains in MI skill, noting that coaching and feedback further enhances skill.⁶ Schwalbe et al. showed that MI training effects were best maintained through at least 3–4 postworkshop contacts, totaling a minimum of 5 hours of coaching from an expert MI trainer over a 6-month period.⁷ Evidence for best practices are starting to emerge, most notably concerning the value of extended training, including feedback and coaching. However, each review further emphasized the need for methodological quality and consistency in MI training studies to further advance best practices.

In the current review, we focus on key elements that should be included in reports of MI training studies to strengthen future research.^{8–12} The review is organized around 3 broad categories: study methodology, training methods, and evaluation. With regard to study methodology, research design provides important information about the strength of conclusions that can be drawn from a particular study.¹³ Also of import with regard to MI training for SU are the characteristics of trainees and training settings, as these are often highly variable and significantly influence the generalizability of findings.^{9,14,15} With regard to training methods, evidence-based training protocols for MI or other interventions are often not readily available, with a few notable exceptions such as the MIA: STEP (Motivational Interviewing Assessment: Supervisory Tools for Enhancing Proficiency) protocol.¹⁶ As such, thorough descriptions of the training methods will better inform training implementation outside the research context. Descriptions should include specific information about the training methods (e.g., workshop, experiential activities, etc.), how those methods were selected, and trainer qualifications.¹⁰ Given longstanding evidence for the importance of coaching and feedback,^{6,7} information on feedback/coaching methods

(length, number of sessions, method of delivery) is also important. With regard to evaluation, it is important to note whether researchers evaluated competency/skill development with evidence-based assessment tools (e.g., observational, client-completed measures).¹⁷

Given the broad implementation of MI training world-wide and the continual evolution of the MI training literature, it is imperative to systematically evaluate the quality and limitations of the existing literature in an ongoing fashion. This ensures that future research continues to improve in quality and scope. The current study focuses on the quality of methods and descriptions in research on MI training. The current study differs from prior studies, which examined MI across several professional disciplines, health care settings, and target behaviors. There is established evidence for MI efficacy in treating SU disorders; therefore, SU providers are an important focal group for evaluating and refining guidelines for effective MI training, which can eventually be translated to other target behaviors. Thus, the purpose of this study was to advance the MI training research with SU providers by evaluating MI training through updated training quality recommendations. Specifically, the review examines the literature published since a review published by Madson et al.,³ with a focus on study methodology, training methods, and evaluation.

Review procedure

We conducted a systematic review following Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) standards¹⁸ by searching the PsycINFO, PsycARTICLES, MEDLINE, Academic Search Premier, ERIC, and SocIndex databases to identify articles from 2008 to January 2018. Key terms used include Motivational Interviewing, Motivational Enhancement Therapy, Brief Motivational Intervention, Training, Workshop, Implementation, Dissemination, Substance Abuse, SU, Addiction, Alcohol, and Drug. We also reviewed the bibliography page on the MI Web site (<http://motivationalinterview.org/library/biblio.html>) to identify additional articles for the review. Our review resulted in 136 articles, all of which outlined use of MI training methods with providers who were treating patients with SU issues. Although the screening, brief intervention, and referral to treatment (SBIRT) protocols incorporate varying degrees and elements of MI, we decided to exclude such articles because MI is not the sole focus of SBIRT; rather, it is embedded within a broader training protocols that may vary substantially in adherence to practices and principles of MI. We also did not want to confuse MI for SBIRT. Additional articles were excluded if the methodology included survey research or if the treatment targets were not substance abuse. As a result, 25 articles were included in the current review. We chose to include articles that were secondary analyses of previous studies and methods/protocol papers because they help contribute to the training literature. These 3 papers are identified in Table 1. Figure 1 outlines the study inclusion procedures following PRISMA guidelines.¹⁸

The first 2 authors independently evaluated the included articles using a coding form developed by the second author. This form was based on recommendations for evaluating the methodological quality of MI training¹⁰ and was approved by the first 3 authors. Descriptors on the coding form were categorized under study methodology, training

methods, and training evaluation. Study methodology descriptors included the study design, setting characteristics, and trainee characteristics. The training methods included descriptors of MI training method and trainer qualifications. Specific MI training method descriptors included MI workshop type, length and duration of training, feedback and coaching of MI using a valid fidelity measure, and amount and duration of feedback and coaching sessions. Descriptors of training evaluation included trainee outcomes and client outcomes.

Decision rules were also included to determine whether authors provided a thorough description of MI training methods and evaluation. If authors described the type of MI training and specified the number or length of training sessions, we determined that they provided a description of training methods. Further, if authors used a fidelity measure but did not provide feedback, then we determined that feedback and coaching methods were not adequately described. Regarding training evaluation, we determined that authors used a fidelity measure to evaluate trainees' MI competency and proficiency if a combination of summary scores, global scores, and behavioral counts was reported. We also determined that the assessment of continued use of MI skills was appropriately described if authors reported trainee scores after 1, 3, 6, or 12 months of formal MI training.

A detailed account of the study findings is presented in Tables 1, 2, and 3. Specifically, study methodology findings are presented in Table 1. Details of training methods are presented in Table 2, and findings for training evaluation are presented in Table 3. Highlights of each evaluation component are described below.

Results

Study design and sample size

All studies provided sufficient information to code study design. Just over two thirds of the studies ($n = 17$, 68%) were randomized controlled trials of MI training in which participants were randomly assigned either to receive various forms of MI training or to receive MI training or a comparison condition. The remaining studies utilized opentrial designs in which all participants received MI training. Almost all ($n = 24$, 96%) were studies of training in MI as a stand-alone intervention. The remaining study³⁶ provided training in brief behavior change counseling. Eight studies (32%) provided training in MI as part of a larger treatment study in which MI was one of the interventions of interest. All studies with the exception of Baer et al.²⁰ provided information about the number of individuals trained ($n = 24$, 96%). Sample size varied across studies, ranging from 2 to 576 with a mean of 86.25 (SD = 120.44).^{25,34}

Trainee professional backgrounds

Nineteen studies provided specific information about the professional backgrounds of training participants (76%). The most well-represented professions were nursing ($n = 10$, 40%), social work ($n = 8$, 36%), and psychology ($n = 7$, 28%). Alcohol and drug counselors ($n = 7$, 28%) and general medical practitioners ($n = 6$, 24%) were also included in several studies. Two studies (8%) included psychiatrists, and 2 included counselors (8%). Pharmacists and criminal justice employees were included in 1 study each (4%). Nine

studies (36%) included trainees from other backgrounds, such as case managers,³⁷ supervisory staff at community treatment facilities,¹⁵ and other professionals in patient care roles (e.g., nonspecified clinicians,²³ physician's assistants²⁴).

Trainees' and trainers' educational backgrounds and years of experience

In addition to assessing professional background, we identified trainees' education levels and years of clinical experience. Education level was reported in 60% of studies ($n = 15$), and years of clinical experience was reported in 56% ($n = 14$). Fifty-two percent of studies ($n = 13$) included trainees with advanced degrees (i.e., master's or greater). Specifically, 40% of studies ($n = 10$) included trainees who had a master's degree, 28% ($n = 7$) included trainees who had a nonmedical doctoral degree (e.g., PhD, PsyD, DSW), and 8% ($n = 2$) included trainees who had a medical doctoral degree (i.e., MD). In addition, 48% of studies ($n = 12$) included trainees with a bachelor's degree or equivalent, and 32% ($n = 8$) included trainees with less than a bachelor's degree (e.g., associate's degree, certificate, high school diploma).¹ In over a third of the studies (40%, $n = 10$), information regarding trainees' educational background was not reported. Trainees' years of clinical experience varied widely both within and across studies; whereas some trainees reported less than 1 year of clinical experience, others reported over 30. The majority of studies ($n = 20$, 80%) provided some information about trainer qualifications and/or years of experience, although the detail of information varied. Sixty percent of studies ($n = 15$) reported that trainers were members of the Motivational Interviewing Network of Trainers (i.e., MINT); in 12% of studies ($n = 3$), trainers were Department of Veterans Affairs (VA)-certified MI trainers.

Setting characteristics

Twenty studies (80%) provided sufficient information to code the type of program in which training participants provided substance abuse interventions. Just under a quarter of studies (24%, $n = 6$) were conducted with providers in outpatient clinics (e.g., community substance abuse clinics), and another fifth (20%, $n = 5$) were conducted with providers in medical settings (e.g., inpatient units). Two studies (8%) were conducted with providers in residential settings (e.g., community settings with a residential component).² The most commonly coded category for program type was *other* ($n = 9$, 36%), indicating that frequently the training was offered to providers in program types that did not meet our a priori decision rules for coding setting. Examples of these types of programs included trauma centers,²⁴ prison services,³² and pharmacies.³⁵

Format and length of training

Most studies (96%, $n = 24$) provided information about the format of MI trainings.³ Consistent with recommendations,^{3,5,7} the majority of studies (84%, $n = 21$) used a workshop plus feedback/coaching training format as the primary method (i.e., experimental

¹Trainees from a variety of educational backgrounds tended to be included within a single study; as such percentages sum to over one-hundred.

²These two studies conducted trainings in both outpatient and residential settings.

³The one study that did not include information about training³¹ referenced an older paper (i.e., Ager et al., 2005⁵¹) where this information was reported. To be consistent with our a priori decision rules regarding coding, we do not provide information that was not reported in the reviewed paper.³¹

training condition) for at least 1 group of trainees (i.e., in some studies training method was the independent variable). The next most common training method comprised workshops and experiential exercises (e.g., role play, group exercises), used as the primary training method in 8% of studies ($n = 2$). In 1 study (4%), a didactic workshop was used as the primary training method. Although other methods (i.e., readings [$n = 4$, 16%], computer trainings [$n = 1$, 4%]) were used, these served as either the comparison group or steps in a trainee skill-based stepped process²⁷ rather than the primary training format. Just over half of the reviewed studies ($n = 13$, 52%) compared trainings using workshops plus feedback/coaching with another training method. Additionally, in 2 studies (8%), workshops utilizing a general approach to feedback/coaching sessions were compared with trainings in which workshops and enhanced forms of feedback/coaching (i.e., MIA: STEP; MI-LEAF [Motivational Interviewing Language Enhanced Attention and Focus]) were utilized.^{16,40}

The number of training sessions and length of each session varied across studies. Similar to Madson et al.,³ we found that just over half of the studies provided a single training session ($n = 14$, 56%), although the use of 2 training sessions was also common ($n = 5$, 20%). Four training sessions were used in 2 studies (8%). In the remaining 2 studies that reported number of training sessions, trainees underwent either 2 or 5 ($n = 1$, 4%) training sessions (i.e., depending on group assignment) or 8 ($n = 1$, 4%) training sessions. The reporting metric for session length (i.e., number of days vs. hours) varied across studies, limiting our ability to provide summary statistics about the length of sessions. However, as presented in Table 2, studies included in this review tended to provide more total hours of training than those reviewed previously;³ for example, 100% of the studies included in the current review provided more than 8 total hours of training, as compared with 71% of those in the previous review.

Coaching, feedback, and assessment of skills

Perhaps the most promising aspect of the studies included in the current review was a shift toward incorporating coaching and/or feedback into MI training. Despite previous authors stressing the importance of ongoing feedback for the development and maintenance of MI skills,⁷ in a prior review only 6 studies (22%) described the use of supervision and only 1 used ongoing coaching.³ In contrast, 84% of studies ($n = 21$) included in this review reported utilizing at least 1 coaching session, with the majority ($n = 15$, 60%) reporting the use of 4 or more sessions.

As an aspect of coaching, it was common for studies to utilize validated objective measures of performance. Such measures included the Motivational Interviewing Treatment Integrity (MITI; $n = 10$, 40%), the Independent Tape Rater Scale (ITRS; $n = 5$, 20%), the Motivational Interviewing Skill Code (MISC; $n = 2$, 8%), the MI Consultation and Feedback Form (MICAFF), and the Behavior Change Counseling Index (BECCI; $n = 1$ each, 4% each).⁴ In the majority of studies, such measures were used to assess trainees' MI proficiency ($n = 19$, 76%; trainees achieved proficiency in 10 studies) and competency ($n = 19$, 76%; trainees achieved competency in 9 studies).

⁴Some studies utilized multiple measures; as such, percentages sum to over 100.

Training evaluation

With one exception, the validated measures of objective performance that were utilized to provide feedback-based coaching were also utilized to evaluate training outcomes. The only exception was Stein et al. who used the MITI for coaching⁴¹ but the Helpful Responses Questionnaire for outcomes.⁴² In addition to the studies that used validated coding measures based on work samples for both coaching and evaluation, an additional 5 studies (20%) used such measures for evaluation only. The most commonly assessed outcome was MI skills ($n = 23$, 92%), followed by knowledge ($n = 7$, 28%) and self-confidence ($n = 4$, 16%). Although also suggestive of a positive departure from earlier training methods in which few studies did so, just under half of the reviewed studies ($n = 12$, 48%) assessed whether trainees maintained skills after training concluded. In 1 study (4%), trainee satisfaction was assessed. Outcomes classified as other (e.g., trainee readiness to change; utilization of change talk) were assessed in 20% of studies ($n = 5$). Consistent with Madson et al.,³ it was uncommon for studies to assess client outcomes. In the 2 studies that did so,^{16,35} both treatment retention and substance use/abstinence use were assessed, and neither study found these outcomes to be associated with MI training.

Discussion

The goal of this paper was to systematically review the MI training literature for SU treatment, with a focus on the methods and reporting in recent MI training projects. Our review highlights that the literature has grown and improved significantly in quality since prior reviews. For example, the studies reviewed included enhanced descriptions of training methodology, populations, and settings. Moreover, many of the reviewed studies conducted some form of randomized trial that allowed comparison of training conditions. Studies such as these will uncover the optimal conditions to help clinicians acquire MI skills and provide evidence-based implementation methods to incorporate MI into standard practice. Further, a remarkable change over previous reviews was the number of studies that incorporated coaching and the use of valid observational tools. Nonetheless, we also uncovered areas for further development to strengthen the MI training literature.

In line with recommendations for a thorough description of trainee and setting characteristics involved in MI training programs,^{5,7,10} most of the reviewed studies provided adequate details of trainee and setting characteristics. However, the descriptions of trainer characteristics varied. In order to move the science forward, it will be imperative for future studies to include trainer qualifications; to ensure that MI skills are being transferred adequately, we must know that they are being transferred from those who have sufficient knowledge of the technology. Therefore, it is recommended that researchers continue to provide a basic description about trainees and the training setting as well as describe the trainer characteristics and qualifications. Further, information on who is being trained, their training, professional and theoretical (e.g., 12-step, cognitive behavioral) background, and prior experience with MI is important. This information will help clarify whether failures of technology transfer are related to the implementation process (e.g., methodology) or influential determinants of implementation outcome (e.g., trainer, trainee, or organizational issues).⁴³ Training programs and clinical sites will also benefit from this information in

determining the necessary qualifications of trainers in order to expect adequate outcomes from learners.

A promising change in studies of MI training has been the transition from primarily workshop/didactic-based trainings to trainings that incorporate coaching and feedback. In fact, many of the reviewed studies utilized workshop-only conditions as a comparison/control group, to better identify specific advantages of adding feedback/coaching training methods. Although an undeniable step toward progress, an important consideration for future studies is the need for additional description about the exact methods used (i.e., such that a training method could be directly replicated) for control/comparison and study conditions. Additionally, given that feedback is known to enhance training, future progress can be made by examining both the process and content of feedback. Of the reviewed studies, 2 provide excellent examples of how these aspects can be described (e.g., the systematic approach of MIA: STEP; the language focus of MI-LEAF) and compared with “feedback as usual.”^{16,40}

There are various resources describing different MI training exercises and strategies,⁴⁴ and it would benefit MI training research and practice to identify effective experiential MI training strategies. In this regard, researchers may consider modeling their training descriptions after Darnell et al.,²⁴ who provided specifics about the training methods, including the content (e.g., 0–10 scaling questions to identify importance of and confidence in change) covered in training sessions. These detailed descriptions are vital for comparing effects across studies, as well as standardizing training to optimize efficiency and effectiveness of MI training programs and components. Moving forward, quality implementation of MI will entail identifying training components that are necessary for successful outcomes, as well as those components that can be modified based on organization, clinician, or client needs.⁴⁵ Certainly, it is challenging to describe all the specifics of a MI training project given journal word limits. Some of the projects reviewed here^{24,40} or protocol papers²⁶ provide examples for disseminating future MI training projects.

A major finding of the current review was that most studies included some form of coaching and feedback. Research by Forsberg et al. demonstrates one effort to meet these standards.³¹ Clinicians were provided with an initial MI training (workshop with experiential exercises) of 12 hours, followed by 84 hours of additional MI supervision across 11 assessment periods covering 2.5 years. During this time span, a total of 33 coded recordings per clinician were reviewed with a valid integrity measure (i.e., MITI 3.0). Moving forward, additional information about coaching methods can inform an effective, standardized approach to MI training.

We found great variability in the description of coaching methods. More consistent reporting of the processes and procedures would enhance conclusions on how to establish coaching best practices. Specifically, future studies should report the number of coaching sessions intended for each trainee, the number of sessions completed, and the average length of sessions. Some of the studies reviewed used individual coaching, whereas others used group coaching or a combination of approaches. However, we know little about the differential effectiveness of these coaching formats. Similarly, it appears that using valid observational

measures enhance coaching, and it would be important to identify best practices for using these measures. For example, one could investigate the value of focusing on MITI global scores or behavioral counts or both when providing feedback. Similarly, there could be investigations for how this feedback is provided based on using a MI-consistent⁴⁶ or inconsistent coaching style. The coaching and feedback literature may also benefit from incorporating other means of feedback on MI use. Recent developments in assessing client feedback using the Client Evaluation of MI (CEMI)^{47,48} may assist the coaching and feedback process and is worthy of investigation as an additional training element. Empirically identifying these best practices for coaching and feedback will further technology transfer and potentially lower costs associated with implementation, which is particularly important given the resource-intensive training process.

Despite the advancements in MI training, there continues to be a paucity of literature that links training outcomes with client outcomes. We found 2 studies that examined this relationship: Jaffray et al. and Martino et al. linked MI training to client substance use and treatment retention.^{16,35} Moving forward, it will be important to identify not only whether training processes develop clinician competency but ultimately also whether these processes facilitated MI consistent behaviors and positive behavior change in clients. Future MI training research should strive to link training outcomes, including global MI competency and behaviors, to client outcomes, including use behavior and treatment retention, as well as to therapeutic factors such as working alliance and treatment engagement.

This review provides an updated examination of the MI training literature, with a focus on the methods and quality in recent research. Although the findings are encouraging, they need to be considered within study limitations and necessary next steps. First, we limited the review to focus on SU professionals, and it may be valuable to replicate these results with other behaviors and nonprofessionals such as peer specialists/advocates.⁴⁹ Our a priori criteria may have limited some of our findings, and future studies may want to consider broader evaluation criteria to reduce the large number of studies coded as “other.” We also decided to exclude SBIRT training studies from this review. Given the rapid developments in SBIRT,¹⁹ it may be valuable to systematically review those training studies to assess similarities and differences with these findings. Further, we chose to include 1 methods³⁸ and 2 secondary analysis papers^{21,25} that have the potential to inflate MI training components or minimize outcomes. However, these papers were included in this review, as we believe they provide value not better captured elsewhere. Finally, this review addresses just one aspect of MI dissemination and implementation: training. Future investigations of MI training should include diverse methodologies such as qualitative designs and action research as well as mixed methods to better evaluate process and outcomes. Key questions about other factors (e.g., feasibility) that influence the integration of MI into practice settings must be addressed in future research.^{45,50}

Funding

Villarosa-Hurlocker are supported by a training grant (T32-AA018108) from the National Institute on Alcohol Abuse and Alcoholism (NIAAA) in the United States.

References

- [1]. Miller W, Rollnick S. The effectiveness and ineffectiveness of complex behavioral interventions: impact of treatment fidelity. *Contemp Clin Trials*. 2014;37(2):234–241. [PubMed: 24469237]
- [2]. Miller W, Mount K. A small study of training in motivational interviewing: does one workshop change clinician and client behavior? *Behav Cognit Psychother*. 2001;29(4):457–471.
- [3]. Madson M, Loignon A, Lane C. Training in motivational interviewing: a systematic review. *J Subst Abuse Treat*. 2009;36(1): 101–109. [PubMed: 18657936]
- [4]. Barwick M, Bennett L, Johnson S, McGowan J, Moore J. Training health and mental health professionals in motivational interviewing: a systematic review. *Child Youth Serv Rev*. 2012;34(9):1786–1795.
- [5]. Söderlund L, Madson M, Rubak S, Nilsen P. A systematic review of motivational interviewing training for general health care practitioners. *Patient Educ Couns*. 2011;84(1):16–26. [PubMed: 20667432]
- [6]. De Roten Y, Zimmermann G, Ortega D, Despland J. Meta-analysis of the effects of MI training on clinicians' behavior. *J Subst Abuse Treat*. 2013;45(2):155–162. [PubMed: 23537923]
- [7]. Schwalbe C, Oh H, Zweben A. Sustaining motivational interviewing: a meta-analysis of training studies. *Addiction*. 2014;109(8):1287–1294. [PubMed: 24661345]
- [8]. Carroll K Training in vain? *Addiction*. 2016;111(7):1153–1154. [PubMed: 26929091]
- [9]. Hall K, Staiger PK, Simpson A, Best D, Lubman DI. After 30 years of dissemination, have we achieved sustained practice change in motivational interviewing? *Addiction* 2016;111(7):1144–1150. [PubMed: 26216706]
- [10]. Madson MB, Schumacher JA, Baer JS, Martino S. Motivational interviewing for substance use: Mapping out the next generation of research. *J Subst Abuse Treat*. 2016;65:1–5. [PubMed: 26971078]
- [11]. Miller W, Moyers T. Asking better questions about clinical skills training. *Addiction*. 2016;111(7):1151–1152. [PubMed: 26929059]
- [12]. Miller W, Moyers T. Motivational interviewing and the clinical science of Carl Rogers. *J Consult Clin Psychol*. 2017;85(8): 757–766. [PubMed: 28726479]
- [13]. van Tulder MW, Furlan A, Bombardier C, Bouter L, Editorial B. Of The Cochrane Collaboration Back Review Group. Updated method guidelines for systematic reviews in the cochrane collaboration Back Review Group. *Spine* 2003;28(12): 1290–1299. [PubMed: 12811274]
- [14]. Hallgren K, Dembe A, Pace B, et al. Variability in motivational interviewing adherence across sessions, providers, sites, and research contexts. *J Subst Abuse Treat*. 2018;84:30–41. [PubMed: 29195591]
- [15]. Schumacher J, Madson M, Norquist G. Using telehealth technology to enhance motivational interviewing training for rural substance abuse treatment providers: a services improvement project. *Behav Ther* 2011;34(4):64–70.
- [16]. Martino S, Paris M, Añez L, et al. The effectiveness and cost of clinical supervision for Motivational Interviewing: A randomized controlled trial. *J Subst Abuse Treat*. 2016;68:11–23. [PubMed: 27431042]
- [17]. Bellg AJ, Borrelli B, Resnick B, et al. Enhancing treatment fidelity in health behavior change studies: best practices and recommendations from the NIH behavior change consortium. *Health Psychol*. 2004;23(5):443–451. [PubMed: 15367063]
- [18]. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009;151(4):264–269. [PubMed: 19622511]
- [19]. Ager R, Roahen-Harrison S, Rice J, et al. Predictors of adopting motivational enhancement therapy. *Res Soc Work Pract*. 2011;21(1):65–76.
- [20]. Baer J, Wells E, Rosengren D, et al. Agency context and tailored training in technology transfer: a pilot evaluation of motivational interviewing training for community counselors. *J Subst Abuse Treat*. 2009;37(2):191–202. [PubMed: 19339139]

- [21]. Carpenter K, Cheng W, Nunes E, et al. "Old dogs" and new skills: how clinician characteristics relate to motivational interviewing skills before, during, and after training. *J Consult Clin Psychol*. 2012;80(4):560–573. [PubMed: 22563640]
- [22]. Smith J, Carpenter K, Nunes E, et al. Training substance abuse clinicians in motivational interviewing using live supervision via teleconferencing. *J Consult Clin Psychol*. 2012;80(3):450–464. [PubMed: 22506795]
- [23]. Clancy R, Taylor A. Engaging clinicians in motivational interviewing: Comparing online with face-to-face post-training consolidation. *Int J Ment Health Nurs* 2016;25:51–61. [PubMed: 26767548]
- [24]. Darnell D, Dunn C, Atkins D, Ingraham L, Zatzick D. A randomized evaluation of motivational interviewing training for mandated implementation of alcohol screening and brief intervention in trauma centers. *J Subst Abuse Treat*. 2016;60:36–44. [PubMed: 26117081]
- [25]. Decker S, Martino S. Unintended effects of training on clinicians' interest, confidence, and commitment in using motivational interviewing. *Drug Alcohol Depend*. 2013;132(3): 681–687. [PubMed: 23684633]
- [26]. Martino S, Ball S, Nich C, et al. Teaching community program clinicians motivational interviewing using expert and train-the-trainer strategies. *Addiction*. 2011;106(2):428–441. [PubMed: 20925684]
- [27]. Doran N, Hohman M, Koutsenok I. Linking basic and advanced motivational interviewing training outcomes for juvenile correctional staff in California. *J Psychoactive Drugs*. 2011;43(Sup1):19–26.
- [28]. Drapkin M, Wilbourne P, Manuel J, et al. National dissemination of motivation enhancement therapy in the Veterans Health Administration: training program design and initial outcomes. *J Subst Abuse Treat*. 2016;65:83–87. [PubMed: 26951921]
- [29]. Dunn C, Darnell D, Carmel A, et al. Comparing the motivational interviewing integrity in two prevalent models of brief intervention service delivery for primary care settings. *J Subst Abuse Treat*. 2015;51:47–52. [PubMed: 25515624]
- [30]. Forrester D, McCambridge J, Waissbein C, Emlyn-Jones R, Rollnick S. Child risk and parental resistance: can motivational interviewing improve the practice of child and family social workers in working with parental alcohol misuse? *Br J Soc Work*. 2008;38(7):1302–1319.
- [31]. Forsberg L, Lindqvist H, Helgason A. Clinician acquisition and retention of Motivational Interviewing skills: a two-and-a-half-year exploratory study. *Subst Abuse Treat Prev Policy*. 2010;5(1):8. [PubMed: 20465805]
- [32]. Forsberg L, Ernst D, Farbring C. Learning motivational interviewing in a real-life setting: a randomised controlled trial in the Swedish prison service. *Crim Behav Ment Health*. 2011;21(3):177–188. [PubMed: 21706526]
- [33]. Fu S, Roth C, Zillich A, et al. Training primary care clinicians in motivational interviewing: a comparison of two models. *Patient Educ Couns*. 2015;98(1):61–68. [PubMed: 25455795]
- [34]. Isenhardt C, Dieperink E, Lenox R, et al. Training and maintaining motivational interviewing skills in a clinical trial. *J Subst Use*. 2014;19(1–2):164–170.
- [35]. Jaffray M, Matheson C, Bond CM, et al. Does training in motivational interviewing for community pharmacists improve outcomes for methadone patients? A cluster randomised controlled trial. *Int J Pharm Pract*. 2014;22(1):4–12. [PubMed: 23822820]
- [36]. Malan Z, Mash B, Everett-Murphy K. Evaluation of a training programme for primary care providers to offer brief behaviour change counselling on risk factors for non-communicable diseases in South Africa. *Patient Educ Couns*. 2016;99(1):125–131. [PubMed: 26324109]
- [37]. Martino S, Canning-Ball M, Carroll K, Rounsaville B. A criterion-based stepwise approach for training counselors in motivational interviewing. *J Subst Abuse Treat*. 2011;40(4):357–365. [PubMed: 21316183]
- [38]. Martino S, Zimbrea P, Forray A, et al. See one, do one, order one: a study protocol for cluster randomized controlled trial testing three strategies for implementing motivational interviewing on medial inpatient units. *Implementation Sci*. 2015;10(1): 138–149.

- [39]. Mitcheson L, Bhavsar K, McCambridge J. Randomized trial of training and supervision in motivational interviewing with adolescent drug treatment practitioners. *J Subst Abuse Treat.* 2009;37(1):73–78. [PubMed: 19150203]
- [40]. Moyers T, Houck J, Glynn L, Hallgren K, Manuel J. A randomized controlled trial to influence client language in substance use disorder treatment. *Drug Alcohol Depend.* 2017;172:43–50. [PubMed: 28122270]
- [41]. Stein L, Clair M, Soenksen-Bassett S, Martin R, Clarke J. Studying process and proximal outcomes of supervision for motivational interviewing. *Train Educ Prof Psychol.* 2015;9(2):175–182. [PubMed: 26417395]
- [42]. Miller W, Hedrick K, Orlofsky D. The Helpful Responses Questionnaire: A procedure for measuring therapeutic empathy. *J Clin Psychol.* 1991;47(3):444–448. [PubMed: 2066417]
- [43]. Schumacher J, Madson M, Nilsen P. Barriers to learning motivational interviewing: a survey of motivational interviewing trainers' perceptions. *J Addict Offender Couns.* 2014;35(2):81–96.
- [44]. Rosengren DB. *Building Motivational Interviewing Skills: A Practitioner Workbook.* 2nd Edition). New York: Guilford Press; 2017.
- [45]. Damschroder L, Hagedorn H. A guiding framework and approach for implementation research in substance use disorders treatment. *Psychol Addict Behav.* 2011;25(2):194–205. [PubMed: 21443291]
- [46]. Madson MB, Bullock EE, Speed AC, Hodges SA. Supervising substance abuse treatment: specific issues and a motivational interviewing model In Hess AK, Hess KD, Hess TH. eds. *Psychotherapy Supervision: Theory Research and Practice.* 2nd ed. Hoboken, NJ: John Wiley and Sons; 2008:340–358.
- [47]. Madson M, Mohn R, Stein B, et al. Measuring client perceptions of motivational interviewing: factor analysis of the Client Evaluation of Motivational Interviewing Scale. *J Subst Abuse Treat.* 2013;44(3):330–335. [PubMed: 22999814]
- [48]. Madson M, Villarosa M, Schumacher J, Mohn R. Evaluating the validity of the Client Evaluation of Motivational Interviewing Scale in a brief motivational intervention for college student drinkers. *J Subst Abuse Treat.* 2016;65:51–57. [PubMed: 26935329]
- [49]. Tsai J, Klee A, Martino S, et al. Training peer specialists with mental illness in motivational interviewing: A pilot study. *Psychiatr Rehabil J.* 2017;40(4):354–360. [PubMed: 27786521]
- [50]. Meyers D, Durlak J, Wandersman A. The quality implementation framework: A synthesis of critical steps in the implementation process. *Am J Community Psychol.* 2012;50(3–4):462–480. [PubMed: 22644083]
- [51]. Ager RD, Roahen-Harrison S, Toriello PJ, et al. A study on the effectiveness of a brief motivational enhancement therapy training. *J Addict Dis.* 2005;24(Suppl 1):15–31.




Figure 1.
Flow chart of study inclusion.

Table 1.

Study methodology.

Reference	Study design	Program type	N	Education	Professional background	Years of clinical experience
Ager ¹⁹	SA	—	136	M	—	—
Baer ²⁰	SA, RCT	OP	—	—	—	—
Carpenter ²¹	SA, RCT	OP, RES	58	M, BD, NF	—	7.2
<i>Secondary analysis of Smith²²</i>						
Clancy ²³	SA	OTH	64	DD, M, GD, DG, CT, DI	NRS, PSYCH, SW, OTH	2.8/7.1
Dameil ²⁴	SA, RCT, LTS	OTH	30	DD, M, GD, DG, CT, DI	NRS, SW, OTH	>1–31+
Decker ²⁵	SA, RCT, LTS	OP	91	DG	—	9.1–11.5
<i>Secondary analysis of Martino²⁶</i>						
Doran ²⁷	SA, LTS	—	576	DG	GM, NRS, PSYCH, OTH	11.1
Drapkin ²⁸	SA	—	264	—	ADC, CN, NRS, PSY, PSYCH, SW, OTH	—
Dunn ²⁹	SA, RCT	Med	15	M, BD	SW	4.5–15.4
Forrester ³⁰	SA	OTH	42	—	SW	—
Forsberg ³¹	SA, RCT, LTS	OTH	18	—	OTH	8–10
Forsberg ³²	SA, RCT, LTS	OTH	45	—	CJ	—
Fu ³³	SA, RTC	Med	34	—	GM, NRS	11.1–14.1
Isenhardt ³⁴	SA, RTC	Med	2	DD, MD	PSY, PSYCH	—
Jaffray ³⁵	SA, RCT, LTS	OTH	84	—	PHAR	—
Malan ³⁶	OTH	Med	41	MD, GD	GM, NRS	—
Martino ³⁷	SA	OTH	26	DD, M, BD, AD, NF	ADC, NRS, PSYCH, SW, OTH	12.5
Martino ²⁶	SA, RTC	OP	90	DD, M, BD, AD, NF	ADC, GM, NRS, PSYCH, SW, OTH	8.9–10.7
Martino ¹⁶	SA, RTC, LTS	Med	30	—	GM, NRS	—
Martino ³⁸	SA, RTC	OP	66	DD, M, BD, AD, NF, CT	ADC, OTH	8.9
<i>Methods paper</i>						
Mitcheson ³⁹	SA, RTC	OTH	30	—	ADC	—
Moyens ⁴⁰	SA, RTC	OTH	190	—	ADC, CN, NRS, PSYCH, SW, GM	12.4
Schumacher ¹⁵	SA	—	16	BD or higher, less than BD	ADC, OTH	5.7
Smith ²²	SA, RTC	OP, RES	97	M, BD, NF	—	7.8–8.5

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Reference	Study design	Program type	N	Education	Professional background	Years of clinical experience
Stein ⁴¹	SA, RTC, LTS	—	25	M, BD	—	—

Note. SA: stand-alone; RCT: randomized controlled trial; LTS: larger treatment study; OP: outpatient; Res: residential; Med: medical; OTH: other setting; DD: doctoral degree; MD: medical degree; M: master's degree; GD: graduate degree; DG: degree; BD: bachelor's degree; AD: associated degree; CT: certificate; DI: diploma; NF: no formal training; ADC: alcohol and drug counselor; CJ: criminal justice; GM: general medicine; CN: counseling; NRS: nursing; PSY: psychiatry; PSYCH: psychology; SW: social work; PHAR: pharmacy.

Table 2.

Training methods.

Reference	CPU	RE	WO	WEL	WFC	Comparison groups	No. Of sessions	Session length	Number of coaching sessions	Measure used	% completion
Ager ¹⁹	—	—	—	—	—	—	—	—	0	No	100
Baer ²⁰		X	X		X	WO vs. WFC	2/5	15 hours	Not specified	No	85
Carpenter ²¹		X	X	X	X	WO vs. WEL vs. WFC	2	8 hours	5	MITI	51–75
Clancy ²³		X	X	X	X	WO vs. WFC	2	2 days	3	No	76–100
Darnell ²⁴				X	X	WFC vs. waitlist	1	1 day	2–6	MITI	51–75
Decker	X			X	X	RE vs. WFC vs. TNT WFC	1	15 hours	3	ITRS	?
Doran ²⁷			X		X	None	2	3 days/2 days	0	No	No
Drapkin ²⁸					X	None	1	3.5 days	4	MITI	76–100
Dunn ²⁹				X	X	RCP (high dose) vs. NRCP (low dose)	1	5.9 (low dose) to 9.6 hours (high dose)	5	No	76–100
Forrester ³⁰			X	X	X	WEL vs. WFC	2	1 day	5	No	76–100
Forsberg ³¹				X	X	WFC vs. PAU	1	12 hours	28 (estimate)	MITI	26–50
Forsberg ³²			X	X	X	WEL vs. WFC vs. PAU	1	5 days	5	MITI	76–100
Fu ³³			X	X	X	WEL vs. WFC	1	½ day	6	No	76–100
Isenhardt ³⁴				X	X	None	1	1 day	10	MICAFF	76–100
Jaffray ³⁵			X		X	WEL vs PAU	4	—	0	BECCI	76
Malan ³⁶			X		X	None	4	2 hours	0	MITI, OTH	No
Martino ³⁷	X	X			X	None	8	1–2 hours	1–2	ITRS	51–75
Martino ²⁶		X		X	X	RE vs. WFC vs. TNT WFC	1	15 hours	3	ITRS	76–100
Martino ¹⁶			X	X	X	WO vs. WFC	1	1 day	40 (estimate)	ITRS	76–100
Martino ³⁸				X	X	WFC vs. WFC*	1	8 hours	up to 7	ITRS	51–75
Mitcheson ³⁹		X		X	X	WFC vs. RE	1	2 days	4	No	51–75
Moyers ⁴⁰				X	X	WFC vs. WFC*	2	2 days	4 (2 individual, 2 group)	MITI, MISC	76–100
Schumacher ¹⁵				X	X	None	1	2 day	10	MITI	76–100
Smith ²²			X	X	X	WEL vs. WFC vs. WFC-T	1	13 hours	5	MITI	76–100
Stein ⁴¹				X	X	None	-	2–5 days	Not specified	MITI	No

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Note. CPU: computer training; RE: reading; WO: workshop only; WEL: workshop plus experiential; WFC: workshop and feedback/coaching; PAU: practice as usual; CTT: context-tailored training; TNT: train the trainer; RCP: routine care providers; NRCP: nonroutine care providers; WFCT: workshop, feedback/coaching via teleconference; MITI: MI Treatment Integrity Code; MISC: MI Skill Code; BECCI: Behavior Change Counseling Index; ITRS: Independent Tape Rater Scale; MICAFF: VA MI Consultation and Feedback Form.

* In these trials, enhanced forms of WFC were compared with general forms of WFC.

Table 3.

Training evaluation.

Reference	Outcomes measured	Proficiency met	Competency met	Valid coding tool used	Sustained skill assessed	Client outcome measured	Type of client outcomes	Training connected to outcome
Ager ¹⁹	KN, SK	No	No	No	No	No	—	No
Baer ²⁰	SK	Yes	Yes	MITI	Yes	No	—	No
Carpenter ²¹	SK	Yes	Yes	MITI	Yes	No	—	No
Clancy ²³	KN, SC, OTH	—	—	—	No	No	—	No
Dameil ²⁴	SK	Yes	Yes	MITI	Yes	No	—	No
Decker ²⁵	SC, SK	—	—	IRTS	Yes	No	—	No
Doran ²⁷	SK, OTH	Yes	No	VASE-R	No	No	—	No
Drapkin ²⁸	KN, SK	—	Yes	Not reported	Yes	No	—	No
Dunn ²⁹	SK	Yes	—	MITI	No	No	—	No
Forrester ³⁰	KN, SK, OTH	No	Yes	Not reported	No	No	—	No
Forsberg ³¹	SK	Yes	No	MITI	Yes	No	—	No
Forsberg ³²	SK	No	—	MITI	No	No	—	No
Fu ³³	KN, SC, SK	No	No	MITI	No	No	—	No
Isenhardt ³⁴	SK	Yes	No	MICAFF	No	No	—	No
Jaffray ³⁵	SK	No	No	BECCI	No	Yes	ABS, RET	No
Malan ³⁶	SK	No	No	MITI, OTH	No	No	—	No
Martino ³⁷	SK	Yes	Yes	ITRS	No	No	—	No
Martino ³⁶	SK	Yes	Yes	ITRS	Yes	No	—	No
Martino ¹⁶	SAT, KN, SC, OTH	—	—	—	No	No	—	No
Martino ³⁸	SK	—	Yes	ITRS	Yes	Yes	ABS, RET	No
Mitcheson ³⁹	SK	No	No	MITI	Yes	No	—	No
Moyers ⁴⁰	KN, SK, OTH	—	—	MITI, MISC	Yes	No	—	No
Schumacher ¹⁵	SK	No	No	MITI	Yes	No	—	No
Smith ²²	SK	Yes	Yes	MITI	Yes	No	—	No
Stein ⁴¹	SK	No	No	OTH	No	No	—	No

Note. SAT: satisfaction; KN: knowledge; SC: self-confidence; SK: skills; OTH: other; MITI: MI Treatment Integrity Code; BECCI: Behavior Change Counseling Index; ITRS: Independent Tape Rater Scale; VASE-R: video assessment of simulated encounters; ABS: abstinence; RET: treatment retention.