

A REVIEW

Spread and prevention of some common viral infections in community facilities and domestic homes

J. Barker, D. Stevens¹ and S.F. Bloomfield¹

Pharmaceutical Sciences Institute, School of Life and Health Sciences, Aston University, Birmingham and

¹Unilever Research Port Sunlight Laboratory, Bebington, Wirral, UK

569/10/00: received 4 October 2000, revised 21 February 2001 and accepted 7 March 2001

- | | |
|--|---|
| 1. Summary, 7 | 4.2 Respiratory viruses, 12 |
| 2. Introduction, 7 | 4.3 Herpes simplex virus, 13 |
| 3. The burden of viral infection, 8 | 4.4 Varicella virus, 13 |
| 3.1 Gastrointestinal infections, 8 | 4.5 The infectious dose, 13 |
| 3.2 Respiratory infections, 9 | 5. Evidence that hygiene can play a part in preventing transmission of viral infections, 13 |
| 4. Dispersal, persistence and spread of viruses in domestic homes and community settings, 10 | 6. Development of effective hygiene policies for prevention of transmission of viral infections, 16 |
| 4.1 Gastrointestinal viruses, 11 | 7. Conclusions, 17 |
| 4.1.1 Rotavirus, 11 | 8. References, 18 |
| 4.1.2 Norwalk-like viruses, 11 | |
| 4.1.3 Hepatitis A virus, 12 | |

1. SUMMARY

Viruses are probably the most common cause of infectious disease acquired within indoor environments and have considerable impact on human health, ranging from severe life-threatening illnesses to relatively mild and self-limiting or asymptomatic diseases. In particular, viruses causing gastrointestinal and respiratory diseases spread rapidly in the community and cause considerable morbidity. Increasing numbers of people who have impaired immunity, for whom the consequences of infection can be much more serious, are now cared for in 'out of hospital' settings. This review examines the dispersal, persistence and control of some common viruses in the domestic home and in community facilities. There is growing evidence that person-to-person transmission via the hands and contaminated fomites plays a key role in the spread of viral infections and there is a need for wider understanding of the potential for contaminated surfaces to act as unidentified vectors of pathogens in the

transmission cycle. Intervention studies have shown that improved standards of education, personal hygiene (particularly handwashing) and targeted environmental hygiene have considerable impact in the control and prevention of infectious organisms.

2. INTRODUCTION

Nearly one thousand different types of viruses are known to infect humans and it is estimated that they account for approximately 60% of all human infections (Horsfall 1965).

Viruses are spread easily through closed environments such as the home, schools, workplaces, transport systems, etc. Although many of the respiratory and gastrointestinal infections caused by viruses can be asymptomatic or relatively mild and self-limiting (coughs and colds, etc.), they still represent a significant economic burden. Increasing numbers of people who have reduced immunity to infection, for whom the consequences of infection can be much more serious, are now cared for at home. At risk groups include not only the immunocompromised but also the elderly, neonates, pregnant women, hospital patients discharged into the community, individuals using immu-

Correspondence to: J. Barker, Pharmaceutical Sciences Institute, School of Life and Health Sciences, Aston University, Aston Triangle, Birmingham B4 7ET, UK (e-mail: J.E.Barker@aston.ac.uk).

nosuppressive drugs and also those using invasive systems (indwelling catheters) or inhalation systems or devices. Otherwise healthy family members with asthma or allergies also have increased susceptibility to infection. In the UK it is estimated that one in six people in the community belong to an 'at risk' group (Bloomfield 2001). World Health Organisation estimates suggest that, by 2025, there will be more than 800 million people over 65 years old in the world, two-thirds of them in developing countries (Anon. 1998).

Viruses are probably the most common cause of infectious disease acquired within indoor environments. Close personal contact within the home and community settings, such as daycare centres and schools, makes them ideal places for the spread of viral infections. Infected individuals can shed up to 10^{12} virus particles per ml of faeces with the possibility of transfer of the virus by contaminated hands to surfaces in the bathroom or toilet. Viruses that cause tonsillitis, colds, croup, bronchiolitis, influenza, pneumonia and other respiratory tract infections can be spread in aerosolized droplets. Aerosols produced by coughing, sneezing and talking can be inhaled directly by a susceptible host or may settle onto surfaces. Touching hands or fomites, such as eating utensils, towels or doorknobs, inadvertently contaminated with fresh secretions or vomit, etc. from an infected person and then transferring the virus from the hands to the eyes, nose or mouth, are further routes of spread. Infants are especially vulnerable to such infections because they frequently place objects, such as toys, into their mouths. Transfer of viruses to food during handling and preparation via hands and food contact surfaces is an important route of spread of viral gastroenteritis.

Amongst health care professionals there is growing awareness that improved standards of hand, surface and air hygiene in community settings could do much to prevent the spread of viral infections within these environments. The purpose of this paper is to review the evidence base for this assumption. Since viral infections are not easily treated, prevention of infection is still the main route of control. Assessment of the impact of hygiene is made difficult by the general lack of quantitative epidemiological data and, even where evidence for cross-contamination as a causative factor in an outbreak exists, it is always circumstantial. A further problem in assessing whether contamination found on hands or other surfaces might represent a hazard is that the infectious dose can vary significantly according to the pathogenicity of the organism and the immune status of the host. Thus the case for practising good hygiene in these settings rests largely on evidence showing that cross-contamination can occur in these environments coupled with laboratory data demonstrating the efficacy of hygiene procedures in minimizing microbiological contamination.

3. THE BURDEN OF VIRAL INFECTION

3.1 Gastrointestinal infections

In developed countries it is estimated that 30–40% of infectious gastroenteritis cases are attributable to viruses (Thompson 1994). Surveillance data from the UK show that reported outbreaks of viral intestinal infection have increased rapidly over the last 10 years; epidemiological data for 1995 and 1996 (Evans *et al.* 1998) show that rotavirus, astrovirus, Norwalk-like viruses (NLVs; also known as small round structured viruses) and other caliciviruses were responsible for 48% of all reported outbreaks of infectious intestinal disease (IID). Other data indicate that NLVs and rotavirus are the commonest pathogens causing outbreaks of gastroenteritis in homes for the elderly (Djuretic *et al.* 1996; Ryan *et al.* 1997; Dedman *et al.* 1998).

Over the period 1993–96 a UK study involving some 460 000 participants was carried out to evaluate rates of IID in the community and presenting to general practice which has given valuable insights into the epidemiology of viral infections in the community (Wheeler *et al.* 1999). The study indicated that as many as one in five people in the general UK population develop IID each year with an estimated 9.4 million cases occurring annually. It has long been recognized that, since cases and outbreaks related to viral agents are often unreported, the impact of viral intestinal infections may be much greater than national surveillance suggests. The findings of the community study confirmed the validity of this assumption. Wheeler and co-workers estimated that for every one case of rotavirus and NLV reported to national surveillance a further 35 cases of rotavirus and 1562 cases of NLV occur in the community.

UK surveillance covering 1995–96 showed NLV as a significant cause of epidemic gastroenteritis in community residential and nursing homes, accounting for 43% of all reported general gastroenteritis outbreaks (Evans *et al.* 1998). The rate of reported NLV infection reaches a peak in children under 5 years and again in the elderly. Foodborne outbreaks can arise from contaminated raw food such as shellfish and also through secondary contamination from food handlers carrying the virus. Foods implicated in outbreaks are mainly those eaten raw, or those not cooked after handling, e.g. salads, cold meats and fruit.

Worldwide, rotavirus is probably the most important viral pathogen causing diarrhoeal disease in infants, infecting virtually all children aged 3–5 years (Parashar *et al.* 1998a). However, a recent study by Pang *et al.* (2000) of children between 2 months and 2 years of age with acute gastroenteritis, has shown that human caliciviruses are found as commonly as rotaviruses. In developing countries rotavirus accounts for approximately 6% of all diarrhoeal episodes and 20% of all diarrhoea-associated deaths of children under

5 years of age, resulting in an estimated 800 000 childhood deaths each year. It is second only to upper respiratory infections in infants under 2 years old as a major cause of death in the developing world (Glass *et al.* 1997). Each year in the US, 2.7 million children under 5 years old are affected by rotavirus diarrhoea, resulting in 500 000 visits to the doctor and 50 000 hospitalizations (Parashar *et al.* 1998b). A large proportion of hospital admissions due to gastroenteritis in children under 5 years old were caused by rotavirus in both the UK and Hong Kong (Ryan *et al.* 1996; Chan *et al.* 1998). In older children 51% of hospital admissions for acute diarrhoea were associated with rotavirus (Lewis *et al.* 1979). A study by Isaacs *et al.* (1986) showed that 20% of children under 14 years old who visited their doctor with diarrhoea had a rotavirus infection. Indications are that hospital admissions only represent a small percentage of rotavirus infections; the majority will be treated by general practitioners.

Rotavirus infections are highly seasonal, peaking in the winter months (Brandt *et al.* 1982; Ryan *et al.* 1996; Dedman *et al.* 1998). It has been suggested that low humidity and people spending more time indoors contribute to the spread of rotavirus infections (Anon. 1995). Such conditions may make it possible for the virus to be spread by the airborne route through environmental contamination (Brandt *et al.* 1982). A study in the US revealed that rotaviruses infected one or more members in 51% of families, including 28% of children and 13% of adults (Rodriguez *et al.* 1987). Within infected families rotavirus infection was found in 57% of children and 25% of adults. Some adults acquired rotavirus infections a few days after their children's illnesses, suggesting that the children rather than the parents brought infection into the home. Rodriguez *et al.* (1979) also found rotavirus infection in 55% of adult family contacts of children hospitalized with gastroenteritis. In a community study in New Zealand, in families with an index case of rotavirus infection, children were more frequently infected than adults. Once a family member became infected there was a high probability of cross-infection (Grimwood *et al.* 1983).

Among children with diarrhoea attending daycare centres, Lew *et al.* (1991) detected astroviruses and adenoviruses. Astrovirus was significantly more common in children with diarrhoea than those without diarrhoea. Enteric adenoviruses were detected in an equal percentage in children with and without diarrhoea. Children can excrete astrovirus before the onset of diarrhoea and up to 20 d after the diarrhoea has stopped (Mitchell *et al.* 1993). Although astrovirus primarily infects the young, the elderly can also be affected, with reported outbreaks in care homes and hospital wards for the elderly (Gray *et al.* 1987; Lewis *et al.* 1989). Enteric adenoviruses (generally serotypes 40 and 41) are also associated with outbreaks of gastroenteritis in

schools, paediatric hospital wards and nursing homes (LeBaron *et al.* 1990). They may be second to rotavirus as a cause of gastroenteritis in young children (Blacklow and Greenberg 1991).

Globally, hepatitis A virus (HAV) is the most common cause of hepatitis in man (Melnick 1995). Contaminated water or food, particularly filter-feeding shellfish, frequently transmit hepatitis A but other foods are occasionally implicated (raw milk, dairy products and cold meats). The virus is excreted in high numbers in faeces and is spread from person to person primarily by the faecal-oral route. When personal hygiene is not observed, food handlers may unintentionally transfer the virus to food during the incubation period of the disease (Sundkvist *et al.* 2000). Outbreaks of viral hepatitis occur in institutions such as daycare centres, hospitals, nurseries and schools (Bern *et al.* 1992; Dickinson 1992). These outbreaks may lead to secondary cases in the general community (Hadler *et al.* 1980).

3.2 Respiratory infections

Infections caused by influenza viruses, rhinoviruses, coronaviruses and respiratory syncytial viruses (RSVs) are a major health burden. Estimates suggest that adults suffer two to five colds per year and infants and preschool children have about four to eight colds per year (Sperber 1994). Although such infections are often regarded as trivial, taking into account lost days from work and school, hospital admissions and mortality rates in infants and the elderly, the health and economic costs are considerable. Although the common cold can be caused by a number of viruses, rhinoviruses and coronaviruses predominate. Rhinoviruses are responsible for outbreaks of the common cold in the general community such as schools, daycare centres and hospitals (Denny *et al.* 1986; Krilov *et al.* 1986; Kellner *et al.* 1988). Rhinoviruses and coronaviruses have been found to cause a greater disease burden in elderly people living at home, compared with influenza virus or RSV (Nicholson *et al.* 1997). Influenza affects all age groups, but it is the elderly and persons with underlying health problems who are at particular risk from complications of influenza and are more likely to require hospitalization.

Respiratory syncytial virus infections occur all over the world and outbreaks are common in the cold season in temperate climates and in the rainy season in tropical climates. Respiratory syncytial virus is a major cause of respiratory illness in young children, affecting about 90% of children by the age of 2 years (Crowcroft *et al.* 1999; Simoes 1999). School-aged children often carry RSV to their homes and spread infection to younger siblings. Attack rates within families are high, with about 40% of family members, including adults, becoming infected. In most family

outbreaks although more than 95% of infections are symptomatic they are not usually severe (Berglund 1967; Hall *et al.* 1976a). Infants admitted to hospital with RSV bronchiolitis or pneumonia tend to shed the virus abundantly and for prolonged periods allowing ample opportunity for spread (Hall *et al.* 1976b). In adults RSV infection generally results in a 'common cold' type illness although it can sometimes produce a 'flu-like' syndrome indistinguishable from influenza. Antibodies resulting from an early childhood RSV infection do not prevent further RSV infections later in life. Respiratory syncytial virus is known to cause a high incidence of pneumonia and death in the elderly. In England and Wales it is estimated that RSV causes 60–80% more deaths than influenza, causing about 23 000 deaths each winter (Nicholson 1996).

Parainfluenza viruses (PIVs) are a further major group of respiratory pathogens. They cause severe colds, croup, bronchitis and pneumonia in children and adults and in infants the virus can cause life-threatening disease (Hall 1987). Infection is probably spread by aerosols in addition to direct contact with contaminated surfaces (Hall *et al.* 1980). Brady *et al.* (1990) noted that the persistence of PIV on hospital surfaces contaminated with patients' secretions was a potential source of transmission.

4. DISPERSAL, PERSISTENCE AND SPREAD OF VIRUSES IN DOMESTIC HOMES AND COMMUNITY SETTINGS

Cross infection from an infected person to a new host depends on a number of factors, including the number of virus particles shed by the infected person, their stability in the environment, in aerosols or on surfaces and the potential for spread within a closed environment (Valenti 1998). Viruses that increase fluid secretions or irritate the respiratory epithelium induce coughing and sneezing, which in turn increases the shedding and transmission of the virus. Although diarrhoea eliminates organisms from the gut, it increases the potential for contamination of the environment and spread of the virus infection. The more particles shed the greater their survival and the greater the chance of reaching a new host. Equally important, the likelihood of cross infection depends on the number of particles that reach the new host, the immune status of that host and the route by which they become infected.

Virus particles can be shed in large numbers in various body fluids from an infected person or a carrier, including blood, faeces, saliva, urine and nasal secretions. The non-enveloped viruses have greater resistance to drying and thus spread more easily than enveloped viruses, which are less stable in the environment. Laboratory studies show that rotavirus, adenovirus, poliovirus, herpes simplex virus and HAV can survive for significant periods on dry surfaces

(Nerurkar *et al.* 1983; Abad *et al.* 1994). Although few studies have been carried out in domestic homes, studies in children's daycare centres (Lew *et al.* 1991; Keswick *et al.* 1983b; Butz *et al.* 1993) and in hospitals (Samadi *et al.* 1983; Akhter *et al.* 1995) show that viruses can survive on surfaces and that virus transfer and survival on hands play a part in the transmission of infections.

Bellamy *et al.* (1998) investigated the domestic environment for the presence of viruses and body fluids that may contain viruses. Haemoglobin was found on 2% of surfaces (taps, washbasins, toilet bowls and seats), indicating the presence of blood and possible contamination with blood-borne viruses. Amylase (an indicator of saliva, sweat and urine) was found on 29% of surfaces, which were frequently handled or in contact with urine. This highlights that surfaces may remain soiled for some time and may not be thoroughly cleaned. Bellamy *et al.* (1998) also detected enteroviral RNA in three of 448 environmental samples (tap handle, telephone handpiece and toilet bowl). A previous study has shown that enteric viruses can survive on environmental surfaces for up to 60 d (Abad *et al.* 1994).

Virus transmission in childcare facilities was studied using modified cauliflower virus DNA as an environmental marker (Jiang *et al.* 1998). The viral DNA, introduced through treated toy balls, spread within a few hours of handling. Although the marker treated objects were removed after 1 d, the viral DNA continued circulating in the facilities for up to 2 weeks. Hand contact with contaminated surfaces played an important part in transmission. The markers were also detected in the children's homes, on the hands of family members and environmental surfaces, including toys.

Transmission of viruses in a household setting has been recently studied, using bacteriophage ϕ X174 as a model virus with resistance properties similar to polio- or parvoviruses (Rheinbaben *et al.* 2000). Contaminated door handles and skin surfaces were found to be efficient vectors of contamination. At least 14 persons could be contaminated one after another by touching a contaminated door handle. Successive transmission from one person to another could be followed up to the sixth contact person. Transfer from contaminated door handles to other surfaces was also confirmed under everyday life conditions in a flat shared by four students.

Studies focusing on home and community settings are providing a better understanding of how infectious disease is spread in these environments. Such studies suggest that the airborne route is by no means the sole route of transmission of respiratory infections caused by rhinovirus and RSV and that IID, particularly that of viral origin, can arise from various sources of which food is only one. Evans *et al.* (1998) reported that, whereas 174 of 233 outbreaks of infection attributed to *Salmonella* were 'mainly foodborne' and 15

regarded as 'mainly person to person', for 680 reported outbreaks of NLV infection 607 were attributed to person-to-person transfer and only 21 were reported as foodborne.

Although epidemiological and surveillance studies provide vital information on modes of infection transmission during outbreaks, they give only a limited picture of how sporadic person-to-person transmission actually occurs within community and home environments. Although such data are limited, it is generally acknowledged that person-to-person transmission is associated not only with poor hand hygiene but also airborne or surface-to-surface transmission. What cannot be deduced from current data is the relative importance of these different modes of person-to-person transmission and how it may differ for different viral agents and different communities (home, daycare, etc.). In particular, it is difficult to assess the importance of environmental contamination as a source of secondary cases, but recent data show that this can be a significant factor, particularly in the transmission of NLVs (Anon. 1993). Although direct evidence is lacking, outbreaks in hotels and cruise ships in which recurrent waves of infection occurred in successive cohorts of guests strongly suggests transmission of NLV via environmental sites and surfaces (Ho *et al.* 1989; Gellert *et al.* 1994; McEvoy *et al.* 1996; Cheesbrough *et al.* 2000). These and other studies which can be used to assess the routes of transmission of viral infections in community and home settings are discussed in the following sections.

4.1 Gastrointestinal viruses

4.1.1 Rotavirus. Rotavirus is shed in large numbers from an infected person, with faeces often containing $> 10^{12}$ particles per gram. Children and adults can be asymptomatic excretors of rotavirus (Ansari *et al.* 1991a) and rotavirus excretion can persist for up to 34 d after diarrhoea has stopped in symptomatic patients (Pickering *et al.* 1988). More recently, a hospital study showed that 30% of the immunocompetent children excreted rotavirus particles for more than 21 d and as long as 57 d after the onset of diarrhoea (Richardson *et al.* 1998).

Rotavirus can survive on human hands and transfer of infectious virus to animate and non-porous inanimate surfaces has been demonstrated (Sattar *et al.* 1986; Ansari *et al.* 1988). Ward *et al.* (1991) examined the transfer of rotavirus from contaminated surfaces to the mouth and from surfaces to hands to the mouth. All of the volunteers who licked rotavirus-contaminated plates became infected whereas, of those individuals touching the virus-contaminated plates with their fingers and then their mouths, only about half became infected.

A number of studies in child daycare centres have shown that rotavirus can be widely disseminated when outbreaks occur. In one such centre faecal contamination of hands and

the environment was demonstrated during an outbreak of rotavirus diarrhoea (Keswick *et al.* 1983a). Other studies in daycare centres have shown that 16–30% of surfaces sampled can be contaminated with rotavirus. In particular, hand contact surfaces (e.g. refrigerator handle, toilet handles, telephone receivers and toys) and moist surfaces such as sinks, water fountains and water-play tables were contaminated with the virus (Keswick *et al.* 1983a; Wilde *et al.* 1992; Butz *et al.* 1993).

Soule *et al.* (1999) found that there was an increase in the number of environmental surfaces contaminated with rotavirus in a hospital paediatric unit when there was an increase in the number of children suffering from rotavirus gastroenteritis. Of the surfaces in direct contact with children (thermometers, play mats and toys) rotavirus was detected in 63% of samples compared with 36% for surfaces without direct contact (telephones, door handles and washbasins). These findings were similar to those of Akhter *et al.* (1995) who showed that widespread rotavirus contamination of a paediatric ward and playroom correlated with the presence of patients infected with rotavirus.

In a treatment centre in Bangladesh, handwashings from 78% of the attendants of patients with diarrhoea (children under 5 years) were positive for rotavirus antigens (Samadi *et al.* 1983). Rotavirus was also found in handwashings of 19% of attendants of patients with non-rotavirus diarrhoea, indicating that they may have come into contact with other attendants and patients in adjacent beds. This highlights the potential for contaminated hands to spread the infection.

4.1.2 Norwalk-like viruses. Projectile vomiting associated with NLVs is probably a major source of cross-infection because it is estimated that 3×10^7 particles are distributed as an aerosol into the environment during a vomiting attack (Caul 1994). A recent report (Anon. 1993) showed how aerosols produced by vomiting can be inhaled or can contaminate hands or work surfaces, with the potential for subsequent transfer to foods or direct hand-to-mouth transfer. The importance of airborne transmission was demonstrated in a recent outbreak in a restaurant where no food source was detected but an analysis of the attack rate showed an inverse correlation with the distance from a person who had vomited (Marks *et al.* 2000). The potential for secondary transmission via environmental surfaces in semiclosed communities was demonstrated following a wedding reception, where an outbreak of NLV gastroenteritis affected 50% of guests (Patterson *et al.* 1997). The previous day, a kitchen assistant had vomited in a sink that was subsequently used for preparing vegetables eaten by the wedding guests. Further evidence for transmission of NLV via aerosols and environmental surfaces comes from reports of recurrent waves of NLV gastroenteritis occurring in

successive cohorts of guests on a cruise ship over a 2-year period (Ho *et al.* 1989). It was found that the risk of gastroenteritis amongst passengers who shared toilet facilities was twice that of those who had a private bathroom. Nosocomial spread is also a major concern. Norwalk-like virus gastroenteritis in an elderly care unit in a hospital spread rapidly within and between wards, affecting both patients and staff. Analysis of risk exposure showed areas where patients had vomited to be the most significant factor for the spread of NLVs to staff (Chadwick and McCann 1994). A study by Cheesbrough *et al.* (1997) showed that carpets can also harbour NLVs and serve as reservoirs of infection. Two carpet fitters became ill after removing a carpet from a hospital ward 13 d after the last case in a NLV outbreak. Routine vacuuming every day since the outbreak had not removed the virus. During an outbreak of vomiting and diarrhoea due to NLV in a long-stay ward for the mentally ill, 36 environmental samples were collected on the affected ward of which 11 (30%) were positive by reverse transcriptase-polymerase chain reaction. Positive swabs were from lockers, curtains and commodes and were confined to the immediate environment of the affected patients (Green *et al.* 1998). Most recently the potential for environmental spread of NLVs was demonstrated in a prolonged hotel outbreak in successive cohorts of guests (Cheesbrough *et al.* 2000). Environmental sampling demonstrated widespread dissemination of the virus on hand contact and other surfaces. From the patterns of infection it was concluded that, although infectious aerosols were probably the main route of dissemination of infection within a particular cohort of guests, contact with contaminated fomites was the most likely factor responsible for maintaining the outbreak by forming the link between successive cohorts.

4.1.3 Hepatitis A virus. As with other enteric viruses HAV is shed from an infected person in large numbers and is able to survive on environmental surfaces (Mbithi *et al.* 1991) and be readily transferred to hands (Mbithi *et al.* 1992). Fomites are potential risk factors in the spread of the virus, especially in hospital wards, daycare centres or restaurants (Cliver 1983). A recent outbreak of HAV was associated with a public house whose barman had chronic diarrhoea and had served drinks while incubating hepatitis A himself (Sundkvist *et al.* 2000). Fomite transmission by contamination of glasses was the likely route of spread. Assessments of community outbreaks of HAV have shown that persons involved in nappy-changing in daycare centres often handle food and that this is a potential risk for transmission (Hadler and McFarland 1986). Hepatitis A virus may be acquired from children who are excreting HAV, the majority of whom are asymptomatic (Fox *et al.* 1974). A significant percentage (23–52%) of susceptible household contacts of index cases with acute HAV infection are at risk of acquiring

acute HAV infection from the index case (Minuk *et al.* 1994). A higher rate of HAV infection amongst children (71–80%) compared with the parents (29%) suggests that play activity among children is a significant factor for HAV transmission in households.

4.2 Respiratory viruses

It is generally accepted that respiratory viruses, such as those which cause the common cold and flu, are spread from person to person by aerosol transmission due to sneezing and coughing. Nevertheless, there is growing evidence that a significant proportion of flu and particularly cold viruses are spread via hands and surfaces such as handkerchiefs and tissues, tap and door handles, telephones or other surfaces touched by an infected person (Eccles 2000; Goldman 2000). Cross infection can occur either by handshaking or by touching the contaminated surface. Rubbing either the nasal mucosa or the eyes with virus-contaminated hands can cause infection.

Sattar *et al.* (1993) have shown that rhinoviruses can survive on environmental surfaces for several hours. Infectious viruses have been recovered from naturally contaminated objects in the surroundings of persons with rhinovirus colds (Reed 1975). Clean hands can readily pick up the virus by touching or handling such objects (Ansari *et al.* 1991b). As much as 70% of infectious rhinovirus on contaminated hands has been shown to transfer to a recipient's fingers after contact of only 10 s (Gwaltney *et al.* 1978). After handling contaminated coffee cup handles and other objects, more than 50% of subjects developed an infection (Gwaltney and Hendley 1982). Hendley *et al.* (1973) and Reed (1975) have demonstrated that rhinoviruses can survive for several hours on the hands and self-inoculation by rubbing of the nasal mucosa or conjunctivae via virus-contaminated fingers can lead to infection.

Influenza virus can be shed before the onset of symptoms and for up to 7 d after onset and individuals with influenza can be infectious before they develop symptoms and for up to a week afterwards. Both influenza A and B virus have been shown to survive on hard surfaces such as stainless steel and plastic for 24–48 h and on absorbent surfaces such as cloth, paper and tissues for up to 12 h (Bean *et al.* 1982). It was shown that influenza A virus could be transferred from contaminated surfaces to hands for up to 24 h after the surface was inoculated. Epidemiological evidence supports the laboratory data because an influenza outbreak in a nursing home suggested that the virus was spread by staff, through hands contaminated directly with body fluids, or by touching contaminated fomites (Morens and Rash 1995).

There is similar evidence for the environmental survival and spread of PIV and RSV. Ansari *et al.* (1991b) demonstrated the transfer of PIV from stainless steel

surfaces to clean fingers, which suggests that fomites have a role as a reservoir for the spread of the virus. Further, PIV could be recovered from non-absorbent surfaces for as long as 10 h when the surface remained moist. However, Brady *et al.* (1990) found that, when material containing PIV was spread and allowed to dry, virus was only recoverable for up to 2 h. These workers also showed that PIV persisted on the skin for at least 1 h after contamination, which reinforces the need to perform vigorous handwashing before and after contact with patients and their environment. Likewise, Hall *et al.* (1980) showed that RSV was recovered from hands touching surfaces contaminated with fresh secretions from RSV-infected infants. Evidence showing that direct and indirect contact is a key factor in transmission of RSV infection is further reviewed by Goldmann (2000).

4.3 Herpes simplex virus

Humans are the only known reservoir of herpes simplex virus 1 (HSV). The virus is most commonly spread by oral secretions and can be shed by persons with or without symptoms. Herpes simplex virus can be recovered from the skin for up to 2 h after inoculation of the hands with the virus (Bardell 1989). The virus was more readily transmitted from moist drops than from drops which had been allowed to dry, although touching dried virus-containing droplets on the skin with a moistened finger resulted in transmission of the virus. Infectious HSV has also been recovered from environmental surfaces such as doorknobs and toilet seats, although it is not clear what role fomites play in the spread of herpes viruses (Larson and Bryson 1982; Bardell 1990; Bardell 1993).

4.4 Varicella virus

A recent study demonstrated the rapid and broad contamination of the environment with varicella-zoster virus (VZV) when a family member acquired the disease (Asano *et al.* 1999). Eight days after onset of the index case VZV DNA was detected in both samples from the patient and on the surfaces of an air-conditioning filter, a table, television channel push-buttons and a door handle. The virus was also detected on the hands of the parents and children. Two siblings developed the disease 18 d after onset of the index case.

4.5 The infectious dose

The problem in assessing whether contamination in the environment might be a hazard is that the infectious dose can vary significantly according to the immune status of the individual. It is clear that increasingly the variability in immune status of individuals is becoming a significant factor

in community and domestic settings as well as in the hospital environment. Although some viruses survive relatively poorly in the environment, the low infectious dose of many viral pathogens, even for individuals regarded as 'healthy', suggests that, where body fluids naturally contaminate objects with a high viral load, the virus can persist in sufficient numbers to act as sources of infection for several hours, weeks or even months (Sattar and Springthorpe 1999). The infective dose for NLVs may be as low as 10–100 particles, indicating that both aerosol and surface contamination could be a route of transfer of infection (Caul 1994). Likewise, the infective dose for rotavirus may be as few as 10 particles and person-to-person transmission probably perpetuates endemic disease (Ward *et al.* 1986). A minimal infective dose of less than 10 plaque-forming units has been demonstrated for poliovirus (Minor *et al.* 1981). Respiratory viruses also have low infective doses. For rhinoviruses, the infective dose via the nasal route may be less than 1 TCID₅₀ (Couch 1990), i.e. the tissue culture infective dose infecting 50% of the cells. Conversely, PIVs have an infective dose via the intranasal route of 80 TCID₅₀ (Smith *et al.* 1966).

5. EVIDENCE THAT HYGIENE CAN PLAY A PART IN PREVENTING TRANSMISSION OF VIRAL INFECTIONS

Although studies about the survival characteristics of viruses represent an important component in understanding the infection potential and the preventive role of hygiene, much of our knowledge comes from reports of infection outbreaks where hygiene procedures have been defective or from case control studies. Fifteen such reports have been examined in which viral contamination was directly implicated or for which viral agents were likely to have been the cause of the infections. The effects of the hygiene intervention procedures are summarized in Table 1 and relate to daycare and other community centres where the concentration of people and activity provides the most cost-effective setting for evaluation of the impact of hygiene procedures. Although opportunities for cross-contamination and cross-infection may occur less frequently in the home it could be argued that, since the ratio of homes to daycare centres is very large, the impact of these environments on the overall infection rates across a community may not be so dissimilar, even though daycare centres bring more people together. None of the investigations cited relate specifically to the home but Fornasini *et al.* (1992) and Osterholm *et al.* (1992) report studies of disease transmission from daycare centres to the home where it is transferred among family members.

In eight of the 15 studies only the impact of handwashing was evaluated. In a 36-week handwashing education programme in child daycare centres, Black *et al.* (1981) showed that the incidence of diarrhoea in children was

Table 1 The effects of hygiene measures in the control of infection in homes and community settings

Infection/illness	Hygiene interventions			Reported effects	Selected references
	Location	Personal	Environmental		
Rhinovirus infections	Homes	Handwashing with iodine-based handwash		Lower rates of infection in those using the iodine-based as opposed to inactive handwash	Carter <i>et al.</i> (1980)
Diarrhoea, some due to rotavirus or adenovirus	Children's daycare centre	Handwashing by staff and children after toilet activities and before eating		50% reduction in illness compared with non-handwashing group	Black <i>et al.</i> (1981)
Aseptic meningitis (echovirus 30)	Childcare centre	More frequent handwashing among teachers than parents of children		Lower rates of infection	Helfand <i>et al.</i> (1994)
Rotavirus (neonatal)	Hospitals	Education/handwashing		Increased risk for neonatal infection associated with mothers not handwashing during care of the neonate	Kilgore <i>et al.</i> (1996)
Respiratory tract infections (common cold)	Children's daycare centre	Education and careful and frequent handwashing by teachers and children		Reduction in incidence of common cold compared with control centre	Niffenegger (1997)
Respiratory tract infections, viral pathogens identified in 44% of illnesses	Senior daycare centre	Education on handwashing to staff. Use of virucidal alcohol foam		50% decrease in respiratory infections after introducing controls	Falsey <i>et al.</i> (1999)
Respiratory infections (colds)*	Childcare centres	Education/handwashing Use of aseptic nose-wiping		Infection control practices reduced the incidence of colds by 17%	Roberts <i>et al.</i> (2000a)
Respiratory/gastro-intestinal infections*	Child daycare homes	Education/reinforcing handwashing, use of vinyl gloves, alcohol-based handrinse	Disposable nappy changing mats at each nappy change	Symptoms of enteric disease significantly reduced compared with control homes	Butz <i>et al.</i> (1990)
Bronchiolitis (RSV)	Hospital, paediatric ward	Education/handwashing. Staff and parents	Segregation of babies	Seven-fold reduction in incidence of hospital-acquired RSV infection	Isaacs <i>et al.</i> (1991)
NLV infection	Hospital, elderly care unit	Staff reminded of the importance of handwashing	Cleaning and disinfection of the environment, e.g. floors, toilet areas, taps, etc. with hypochlorite solution	Reduction in attack rate after using control measures	Chadwick and McCann (1994)
Respiratory/gastro-intestinal infections*	Pre-school daycare centre	Education/reinforcing handwashing to school staff and families	Surface cleaning and disinfection including toys and school buses	Decrease in incidence of infections	Krilov <i>et al.</i> (1996)
Respiratory tract infections*	Children's daycare homes	Education/handwashing	Removing shared hand drying towels, cleaning of sleeping mats	Rarely washed hands associated with higher incidence of infection	St. Sauver <i>et al.</i> (1998)

Respiratory/gastro-intestinal infections*	Child daycare centres	Education/more frequent handwashing	Cleaning of daycare centres and regular washing of toys	Children and staff had significantly fewer infections compared with control centres	Uhari and Mottonen (1999)
Diarrhoeal illness*	Child daycare centres	Education/reinforcing handwashing to staff and children		When children's compliance with handwashing was high, diarrhoea episodes reduced by 66%	Roberts <i>et al.</i> (2000b)
NLV infection	Cruise ship		Cleaning and disinfection of cabins, communal bathrooms with chlorine-based compound, steam cleaning of soft furnishings	Significant reduction in incidence after cleaning	McEvoy <i>et al.</i> (1996)

*Causative agents not specified.

RSV, Respiratory syncytial virus; NLV, Norwalk-like virus.

significantly reduced compared with two control centres. Kilgore *et al.* (1996) studied the prevalence of neonatal rotavirus infection in Bangladesh. They found an increased risk for neonatal rotavirus infection among infants whose mothers reported no handwashing during care of the neonate.

In a study carried out during the cold and flu season at two daycare centres, fewer colds were reported in the test group of 3–5-year-olds using proper and frequent handwashing techniques than in the control group. In the test centre the proportion of colds remained fairly constant at 18.9% whilst in the control group the proportion of colds increased from 12.7% to 27.8% (Niffenegger 1997). Carter *et al.* (1980) demonstrated that families who used an iodine-based hand disinfectant, known to kill rhinoviruses, had lower rates of infection than families using an inactive handwash. To decrease respiratory infections in senior daycare centres, staff were educated on viral transmission and the value of handwashing (Falsey *et al.* 1999). In the intervention year, the infection rate among those attending the centres was significantly lower than in the previous 3 years, with an almost 50% decrease in the infection rate. Roberts *et al.* (2000a) carried out a randomized controlled trial of the effect of infection control measures on the frequency of upper respiratory infection in childcare. The intervention measures were training of childcare staff about transmission of infection, handwashing and aseptic nose-wiping technique. When compliance with infection control practice was high, the incidence of colds was reduced by 17%. A similar study by these workers also examined the effects of infection control measures on the frequency of diarrhoeal episodes in childcare using a randomized controlled trial (Roberts *et al.* 2000b). They found that, for those centres in which children's compliance with handwashing was high, diarrhoeal episodes were reduced by 66%. In the US, an outbreak of aseptic meningitis due to echovirus 30 was reported amongst parents with children attending a childcare centre. It was found that more frequent handwashing among the teachers compared with the parents of young children was associated with significantly lower rates of infection (Helfand *et al.* 1994).

In six of the 15 studies handwashing combined with environmental decontamination or other control measures was considered, whilst one study which related to NLV infection highlighted only the importance of environmental disinfection. In a preschool daycare centre, respiratory and gastrointestinal infections decreased following implementation of measures which included reinforcing existing handwashing procedures and education of staff and families on issues of infection control including environmental surface cleaning and disinfection and disinfection of toys (Krilov *et al.* 1996). Uhari and Mottonen (1999) evaluated an infection prevention programme for reducing the

transmission of infections in child daycare centres. It was evident that most of the infections that did occur were viral. The programme included increased handwashing, cleaning of the daycare centres and regular washing of toys. Both the children and staff had significantly fewer infections than those in control centres.

St. Sauver *et al.* (1998) studied hygienic practices and the prevalence of respiratory illness in children attending daycare homes. Never or rarely washing hands by both children and carers was associated with a higher frequency of respiratory illness in both family and group daycare homes. Using shared cloth towels rather than individual paper towels and washing of sleeping mats less than once a week were also associated with a higher frequency of upper respiratory infection.

Isaacs *et al.* (1991) reported a sevenfold reduction in the incidence of RSV in a hospital when patients and staff were educated about the importance of handwashing and infected babies were segregated. Before intervention 4.2% of children under 2 years old developed nosocomial RSV, whilst after intervention only 0.6% developed infection. Following implementation of a hygiene intervention programme that included handwashing education, use of gloves, disposable nappy pads and an alcohol-based hand rinse the incidence of enteric illness was lowered in intervention child daycare homes as compared with control homes (Butz *et al.* 1990).

As stated previously, contamination of the environment may be considerable during an outbreak of NLV. Following outbreaks of viral gastroenteritis on consecutive cruises, a ship was cleared and disinfected at the end of the fourth cruise in order to interrupt transmission of NLV (McEvoy *et al.* 1996). Fewer than 10 cases presented in each of the fifth and sixth cruises compared with 195 cases during the fourth cruise. Control measures included cleaning and disinfection of cabins, crew and staff quarters and communal bathrooms and steam cleaning of soft furnishings. Hygiene measures were also introduced into the kitchen. The contamination of soft furnishing in areas where individuals have vomited presents a difficult cleaning problem and steam cleaning has been recommended by McEvoy *et al.* (1996).

During a hospital gastroenteritis outbreak caused by NLV, the attack rate among patients decreased in several wards following the implementation of environmental hygiene procedures (Chadwick and McCann 1994). Infection control measures implemented included cleaning and chemical disinfection (ward floors, toilet areas, toilet seats, taps and spillages of vomit and faeces) to reduce environmental contamination. Hypochlorite solution (1%) was used for disinfection of places contaminated with vomit or faeces and 0.1% hypochlorite for general disinfection of ward floors and toilet areas.

Since the studies highlighted in Table 1 involved the implementation of infection control programmes involving several steps, it is difficult to attribute the effectiveness of the hygiene procedures to one specific aspect because they were inherently multifaceted. Indeed, the results do not prove that the various interventions had a direct effect in decreasing infection rates. It could be argued that the reduced rates were due to variations in individual susceptibility to infection or to a lower incidence of the pathogen amongst the case control groups. Nevertheless, overall, there is convincing circumstantial evidence to suggest that improved standards of hygiene can have a significant impact in reducing the rates of respiratory, intestinal and other viral infections in childcare facilities, domestic homes, hospitals and adult care centres and the circulation of infections between these communities.

6. DEVELOPMENT OF EFFECTIVE HYGIENE POLICIES FOR PREVENTION OF VIRAL INFECTIONS

The data reviewed show how improved standards of education and integrated hygiene measures, including hand and environmental hygiene, could have a significant impact in reducing infectious diseases within community and home environments. In recent years the concept of risk assessment or Hazard Analysis Critical Control Point (HACCP) has successfully controlled microbial risks in food and other manufacturing environments. Traditionally, the public has tended to regard good hygiene as creating an environment free of germs. To devise a hygiene policy that has real public health benefits, it is now accepted that a risk-based approach should also be adopted (Bloomfield and Scott 1997; Jones 1998; Scott 1999).

A risk assessment approach to hygiene starts from the premise that homes and other settings always contain potentially harmful microbes (people, pets, food, etc.) and that good hygiene is not about eradication but about targeting measures in the places and at the times that matter, in order to limit risks of exposure. For both the hands and for environmental surfaces hygiene can be achieved by physical removal of organisms from the surface. Alternatively, organisms can be inactivated *in situ* by a disinfection process or a combination of both physical removal and disinfection. In many situations such as the hands, and cooking and eating utensils, appropriate risk reduction can be achieved using detergent and hot water. However, since, in this situation, hygiene is achieved by removal of the microbes from the surface, if it is to be effective it must be applied in conjunction with a thorough rinsing process with clean water and must take account of the strength of attachment of the microbes to the surface (Eginton *et al.* 1995). Stevens and Holah (1993) showed that

wiping of contaminated abraded surfaces using a sponge followed by a 10-s rinse produced a 3-log reduction in bacterial contamination of stainless steel surfaces but only a 1–1.5-log reduction on enamelled steel, mineral resin and polycarbonate surfaces. Scanning electron microscope studies showed that bacteria were typically retained in surface imperfections, such that surfaces which sustained the most extensive damage retained higher numbers of bacteria.

Studies by Schurmann and Eggers (1985) showed that enteric viruses may be more strongly bound to the skin surface and that the inclusion of an abrasive substance, such as sand or aluminium hydroxide, in the handwash preparation is advisable to achieve effective virus removal. Recent studies of the transmission of viruses in a household setting using bacteriophage ϕ X174 as a model showed that virus spread was not prevented by the usual standards of hand hygiene as practised in the household (Rheinbaben *et al.* 2000). The virus was reisolated after 24 h from the hands of all persons tested even after normal use and cleaning of the hands.

Biocides that have activity against both enveloped and non-enveloped viruses include chlorine- and iodine-releasing agents, peracids and ozone (Rotter 1997; Sattar and Springthorpe 1999). Biocide effectiveness depends on the nature of the virus, the surface carrier, the presence of interfering substances such as organic soil and hard water salts and the contact time. Although these compounds can be used for disinfection of environmental surfaces they are generally too toxic and irritant for use on the skin. In achieving decontamination of hands, although handrub and handwash products currently available may have good activity against bacterial pathogens, activity against viral contamination is variable and depends on the type of virus. Rotter (1997) suggested that, although alcoholic handrubs are effective against enveloped viruses such as influenza, PIV, herpes and RSV, activity against non-enveloped viruses such as rotaviruses, rhinovirus, poliovirus, adenovirus, NLV and hepatitis virus is limited unless extended contact times (up to 10 min) are used. Similarly, agents such as triclosan and chorhexidine have some activity against enveloped virus but are not considered effective against non-enveloped viruses.

7. CONCLUSIONS

It is well established that viruses are shed in large numbers and can survive for long periods on surfaces or fomites commonly found in many environments and this emphasizes the possible role of surfaces in the transmission of viruses. Faeces can contain up to 10^{12} virus particles per gram and vomit up to 10^7 per millilitre so the potential for hand and environmental contamination is considerable. Viral shedding may begin before the onset of symptoms and may continue for several days or even weeks after the symptoms have ceased. Virus

transfer from surfaces to hands, fingers and food has been demonstrated. Other studies have shown a high rate of spread once a viral infection is introduced into a family home or institution. Improved handwashing and surface hygiene procedures have been shown to interrupt the transmission of viral infections via hands, surfaces or fomites.

Although the importance of hygiene and most particularly handwashing cannot be over-emphasized as a means of reducing infections it can be difficult to enforce even in healthcare facilities where staff should be aware of the infection risks. Studies have shown that handwashing compliance amongst healthcare workers is variable (Daniels and Rees 1999; McGuckin *et al.* 1999; Nishimura *et al.* 1999; Pittet *et al.* 1999). In a department of surgery, clinicians washed their hands between examinations in only 41% of cases (Daniels and Rees 1999). In continuous video-camera surveillance of an intensive care unit personnel complied with handwashing in 71% of entries, whereas visitors of patients complied in 94% of entries (Nishimura *et al.* 1999). A recent survey of beliefs and attitudes towards hygiene in domestic homes showed that over 74% recognized handwashing as a key preventative measure in ensuring food safety and 66% believed that surface cleaning was also important. However, the respondents admitted that they would not carry out these procedures as frequently as they thought they should (Mathias 1999).

The importance of hands in the transmission of virus infections is well recognized and many of the studies cited in this review relate specifically to handwashing interventions. Increasingly however, there is evidence that cross contamination via surfaces is a significant contributory factor. Most particularly hand contact with contaminated surfaces is likely to be the cause of such cross contamination. Sattar and Springthorpe (1999) and Lieberman (1994) emphasized that, if hygiene programmes are to be effective, hand hygiene education must be integrated with education about the importance of surface and air hygiene in prevention of infection transmission.

To motivate changes in attitude to hygiene it will be necessary to gain acceptance that homes and other community settings will always contain potentially harmful microbes and that good hygiene is not about eradication but about targeting the correct measures at the times that matter, in order to reduce infection risks. There is a need for wider understanding of the potential for contaminated surfaces to act as unidentified vectors of pathogens, in the recontamination of hands, during the infection transmission cycle. The epidemiological evidence to date shows that raising awareness about the importance of key procedures, such as through handwashing and surface hygiene (particularly hand and food contact surfaces), will have a considerable impact in the control and prevention of infectious organisms.

8. REFERENCES

- Abad, F.X., Pinto, R.M. and Bosch, A. (1994) Survival of enteric viruses on environmental fomites. *Applied and Environmental Microbiology* **60**, 3704–3710.
- Akhter, J., Al-Hajjar, S., Myint, S. and Hussain Qadri, S.M. (1995) Viral contamination of environmental surfaces on a general paediatric ward and playroom in a major referral centre in Riyadh. *European Journal of Epidemiology* **11**, 587–590.
- Anon. (1993) Outbreaks of viral gastroenteritis associated with SRSVs. *PHLS Microbiology Digest* **10**, 2–8.
- Anon. (1995) A working party of the Public Health Laboratory service Salmonella Committee. The prevention of human transmission of gastro-intestinal infections, infestations and bacterial intoxications. *Communicable Disease Report Review* **5**, R158–R172.
- Anon. (1998) *The World Health Report 1998 Life in the 21st Century, a Vision for All*. Geneva: World Health Organisation.
- Ansari, S.A., Sattar, S.A., Springthorpe, G.A., Wells, G.A. and Tostowaryk, W. (1988) Rotavirus survival on human hands and transfer of infectious virus to animate and non-porous inanimate surfaces. *Journal of Clinical Microbiology* **26**, 1513–1518.
- Ansari, S.A., Springthorpe, V.S. and Sattar, S.A. (1991a) Survival and vehicular spread of human rotaviruses — possible relation to seasonality of outbreaks. *Reviews of Infectious Diseases* **13**, 448–461.
- Ansari, S.A., Springthorpe, V.S., Sattar, S.A., Rivard, S. and Rahman, M. (1991b) Potential role of hands in the spread of respiratory viral infections — studies with human parainfluenza virus 3 and rhinovirus 14. *Journal of Clinical Microbiology* **29**, 2115–2119.
- Asano, Y., Yoshikawa, T., Ihira, M., Furukawa, M., Suzuki, K. and Suga, S. (1999) Spread of varicella-zoster virus DNA to family members and environments from siblings with varicella in a household. *Paediatrics* **103**, E611–E613.
- Bardell, D. (1989) Hand-to-hand transmission of herpes simplex virus type 1. *Microbios* **59**, 93–100.
- Bardell, D. (1990) Survival of herpes simplex virus type 1 on some frequently touched objects in the home and public buildings. *Microbios* **63**, 145–150.
- Bardell, D. (1993) Survival of herpes simplex virus type 1 in saliva and tap water contaminating some common objects. *Microbios* **74**, 81–87.
- Bean, B., Moore, B.M., Peterson, L.R., Gerding, D.N. and Balfour, H.H. (1982) Survival of influenza viruses on environmental surfaces. *Journal of Infectious Diseases* **146**, 47–51.
- Bellamy, K., Laban, K.L., Barrett, K.E. and Talbot, D.C.S. (1998) Detection of viruses and body fluids which may contain viruses in the domestic environment. *Epidemiology and Infection* **121**, 673–680.
- Berglund, B. (1967) Respiratory syncytial virus infections in families. *Acta Paediatrica Scandinavica* **56**, 395–404.
- Bern, C., Martinez, J., de Zoysa, I. and Glass, R.I. (1992) The magnitude of the global problem of diarrhoeal disease: a ten year update. *WHO Bulletin* **70**, 705–714.
- Black, R.E., Dykes, A.C., Anderson, K.E., Wells, J.G., Sinclair, S.P., Gary, G.W. Jr, Hatch, M.H. and Gangarosa, E.J. (1981) Handwashing to prevent diarrhoea in day-care centres. *American Journal of Epidemiology* **113**, 445–451.
- Blacklow, N.R. and Greenberg, H.B. (1991) Viral gastroenteritis. *New England Journal of Medicine* **325**, 252–264.
- Bloomfield, S.F. (2001) How hygienic should our homes be? *Professional Care of Mother and Child* **11**, 3–4.
- Bloomfield, S.F. and Scott, E. (1997) Cross-contamination and infection in the domestic environment and the role of chemical disinfectants. *Journal of Applied Microbiology* **83**, 1–9.
- Brady, M.T., Evans, J. and Cuartas, J. (1990) Survival and disinfection of parainfluenza viruses on environmental surfaces. *American Journal of Infection Control* **18**, 18–23.
- Brandt, C.D., Kim, H.W., Rodriguez, W.J., Arrobio, J.O., Jeffries, B.C. and Parrott, R.H. (1982) Rotavirus gastroenteritis and weather. *Journal of Clinical Microbiology* **16**, 478–482.
- Butz, A.M., Fosarelli, P., Dick, J., Cusack, T. and Yolken, R. (1993) Prevalence of rotavirus on high-risk fomites in day care facilities. *Paediatrics* **92**, 202–205.
- Butz, A.M., Larson, E., Fosarelli, P. and Yolken, R. (1990) Occurrence of infectious symptoms in children in day care homes. *American Journal of Infection Control* **18**, 347–353.
- Carter, C.H., Hendley, J.O., Mika, L.A. and Gwaltney, J.M. (1980) Rhinovirus inactivation by aqueous iodine *in vitro* and on skin. *Proceedings of the Society for Experimental Biology and Medicine* **165**, 380–383.
- Caul, E.O. (1994) Small round structured viruses — airborne transmission and hospital control. *Lancet* **343**, 1240–1242.
- Chadwick, P.R. and McCann, R. (1994) Transmission of a small round structured virus by vomiting during a hospital outbreak of gastroenteritis. *Journal of Hospital Infection* **26**, 251–259.
- Chan, P.K.S., Tam, J.S., Nelson, E.A.S., Fung, K.S., Adeyemi-Doro, F.A.B., Fok, T.F. and Cheng, A.F. (1998) Rotavirus infection in Hong Kong: epidemiology and estimates of disease burden. *Epidemiology and Infection* **120**, 21–325.
- Cheesbrough, J.S., Barkess-Jones, L. and Brown, D.W. (1997) Possible prolonged environmental survival of small round structured viruses. *Journal of Hospital Infection* **35**, 325–326.
- Cheesbrough, J.S., Green, J., Gallimore, C.I., Wright, P.A. and Brown, D.W.G. (2000) Widespread environmental contamination with Norwalk-like viruses (NLV) detected in a prolonged hotel outbreak of gastroenteritis. *Epidemiology and Infection* **125**, 93–98.
- Clover, D.O. (1983) *Manual on Food Virology*. Publication No. VPH/83.46. Geneva: World Health Organisation.
- Couch, R.B. (1990) Rhinoviruses. In *Fields Virology* eds. Fields, B.N., Knipe, D.M., Chanock, R.M., Hirsch, M.S., Melnick, J.L., Monath, T.P. and Roizman, B. pp. 607–629. New York: Raven Press.
- Crowcroft, N.S., Cutts, F. and Zambon, M.C. (1999) Respiratory syncytial virus: an underestimated cause of respiratory infection, with prospects for a vaccine. *Communicable Disease and Public Health* **2**, 234–241.
- Daniels, I.R. and Rees, B.I. (1999) Handwashing: simple but effective. *Annals of the Royal College of Surgeons of England* **81**, 117–118.
- Dedman, D., Laurichesse, H., Caul, E.O. and Wall, P.G. (1998) Surveillance of small round structured virus infection in England and Wales, 1990–5. *Epidemiology and Infection* **121**, 139–149.
- Denny, F.W., Collier, A.M. and Henderson, F.W. (1986) Acute respiratory infections in day care. *Review of Infectious Diseases* **8**, 527–532.
- Dickinson, C.J. (1992) Hepatitis A — new information on an old virus. *Hepatology* **16**, 1099–1101.

- Djuretic, T., Ryan, M.J., Fleming, D.M. and Wall, P.G. (1996) Infectious intestinal disease in elderly people. *Communicable Disease Report Review* **6**, R107–R112.
- Eccles, R. (2000) International Scientific Forum on Home Hygiene. Spread of common colds and influenza. World Wide Web. [Http://www.ifh-homehygiene.org/newspage/new05.htm](http://www.ifh-homehygiene.org/newspage/new05.htm).
- Eginton, P.J., Gibson, H., Holah, J., Handley, P.S. and Gilbert, P. (1995) Quantification of the ease of removal of bacteria from surfaces. *Journal of Industrial Microbiology* **15**, 305–310.
- Evans, H.S., Madden, P., Douglas, C., Adak, G.K., O'Brien, S.J., Djuretic, T., Wall, P.G. and Stanwell-Smith, R. (1998) General outbreaks of infectious intestinal disease in England and Wales: 1995 and 1996. *Communicable Disease and Public Health* **1**, 165–171.
- Falsey, A.R., Criddle, M.M., Kolassa, J.E., McCann, R.M., Brower, C.A. and Hall, W.J. (1999) Evaluation of a handwashing intervention to reduce respiratory illness rates in senior day-care centers. *Infection Control and Hospital Epidemiology* **20**, 200–202.
- Fornasini, M., Reves, R.R., Murray, B.E. and Pickering, L.K. (1992) Trimethoprim-resistant *Escherichia coli* in households of children attending day care centres. *Journal of Infectious Diseases* **166**, 326–330.
- Fox, M.K., Langner, S.B. and Wells, R.W. (1974) How good is handwashing practice? *American Journal of Nursing* **74**, 1676–1678.
- Gellert, G.A., Waterman, S.H. and Ewert, D. (1994) An outbreak of gastro-enteritis caused by small round structured virus in a geriatric convalescent facility. *Infection Control and Hospital Epidemiology* **11**, 459–464.
- Glass, R.I., Bresee, J.S., Parashar, U.D., Miller, M.A. and Gentsch, J.R. (1997) Rotavirus vaccines at the threshold. *Nature Medicine* **3**, 10–11.
- Goldmann, D.A. (2000) Transmission of viral respiratory infections in the home. *Paediatric Infectious Disease Journal* **19**, S97–S102.
- Gray, J.J., Wreghitt, T.G., Cubitt, W.D. and Elliott, P.R. (1987) An outbreak of gastroenteritis in a home for the elderly associated with astrovirus type 1 and human calicivirus. *Journal of Medical Virology* **23**, 377–381.
- Green, J., Wright, P.A., Gallimore, C.I., Mitchell, O., Morgan-Capner, P. and Brown, D.W.G. (1998) The role of environmental contamination with small round structured viruses in a hospital outbreak investigated by reverse-transcriptase polymerase chain reaction assay. *Journal of Hospital Infection* **39**, 39–45.
- Grimwood, K., Abbott, G.D., Fergusson, D.M., Jennings, L.C. and Allan, J.M. (1983) Spread of rotavirus with families: a community-based study. *British Medical Journal* **287**, 57–59.
- Gwaltney, J.M. and Hendley, J.O. (1982) Transmission of experimental rhinovirus infection by contaminated surfaces. *American Journal of Epidemiology* **116**, 828–833.
- Gwaltney, J.M. Jr, Moskalski, P.B. and Hendley, J.O. (1978) Hand-to-hand transmission of rhinovirus colds. *Annals of Internal Medicine* **88**, 463–467.
- Hadler, S.C. and McFarland, L. (1986) Hepatitis in day care centers: epidemiology and prevention. *Reviews of Infectious Diseases* **8**, 548–557.
- Hadler, S.C., Webster, H.M., Erben, J.J., Swanson, J.E. and Maynard, J.E. (1980) Hepatitis A in daycare centers — a community-wide assessment. *New England Journal of Medicine* **302**, 1222–1227.
- Hall, C.B. (1987) Parainfluenza viruses. In *Textbook of Paediatric Infectious Diseases* eds Feigin, R.D. and Cherry, J.D. pp. 1632–1646. Philadelphia: W.B. Saunders.
- Hall, C.B., Douglas, R.G. Jr and Geiman, J.M. (1976b) Respiratory syncytial virus infections in infants: quantitation and duration of shedding. *Journal of Paediatrics* **89**, 11–15.
- Hall, C.B., Douglas, R. Jr and Geiman, J.M. (1980) Possible transmission by fomites of respiratory syncytial virus. *Journal of Infectious Diseases* **141**, 98–102.
- Hall, C.B., Geiman, J.M., Biggar, R., Kotok, D.I., Kogan, P.M. and Douglas, R.G. (1976a) Respiratory syncytial virus infections within families. *New England Journal of Medicine* **294**, 414–419.
- Helfand, R.F., Khan, A.S., Pallansch, M.A., Alexander, J.P., Meyers, H.B., DeSantis, R.A., Schonberger, L.B. and Anderson, L.J. (1994) Echovirus 30 infection and aseptic meningitis in parents of children attending a child-care centre. *Journal of Infectious Diseases* **169**, 1133–1137.
- Hendley, J.O., Wenzel, R.P. and Gwaltney, J.M. (1973) Transmission of rhinovirus colds by self inoculation. *New England Journal of Medicine* **288**, 1361–1364.
- Ho, M.-S., Glass, R.I., Monroe, S.S., Madore, H.P., Stine, S., Pinsky, P.F., Cubitt, D., Ashley, C. and Caul, E.O. (1989) Viral gastroenteritis aboard a cruise ship. *Lancet* **2**, 961–965.
- Horsfall, F.L. Jr (1965) General principles and historical aspects. In *Viral and Rickettsial Infections of Man* ed. Horsfall, F.L. Jr & Tamm, I. pp. 1–10. New York: Lippincott.
- Isaacs, D., Day, D. and Crook, S. (1986) Childhood gastroenteritis: a population study. *British Medical Journal* **293**, 545–546.
- Isaacs, D., Dickson, H., O'Callaghan, C., Sheaves, R., Winter, A. and Moxon, E.R. (1991) Handwashing and cohorting in prevention of hospital acquired infections with respiratory syncytial virus. *Archives of Disease in Childhood* **66**, 227–231.
- Jiang, X., Dai, X., Goldblatt, S., Buescher, C., Cusack, T.M., Matson, D.O. and Pickering, L.K. (1998) Pathogen transmission in child care settings studied by using a cauliflower virus DNA as a surrogate marker. *Journal of Infectious Diseases* **177**, 881–888.
- Jones, M.V. (1998) Application of HACCP to identifying hygiene risks in the home. *International Biodeterioration and Biodegradation* **41**, 191–199.
- Kellner, G., Popow-Kraupp, T., Kundi, M., Binder, C., Wallner, H. and Kunz, C. (1988) Contribution of rhinoviruses to respiratory viral infections in childhood: a prospective study on a mentally hospitalized infant population. *Journal of Medical Virology* **25**, 455–469.
- Keswick, B.H., Pickering, L.K., DuPont, H.L. and Woodward, W.E. (1983a) Prevalence of rotavirus in children in day-care centres. *Journal of Paediatrics* **103**, 85–86.
- Keswick, B.H., Pickering, L.K., DuPont, H.L. and Woodward, W.E. (1983b) Survival and detection of rotaviruses on environmental surfaces in day care centres. *Applied and Environmental Microbiology* **46**, 813–816.
- Kilgore, P.E., Unicomb, L.E., Gentsch, J.R., Albert, M.J., McElroy, C.A. and Glass, R.I. (1996) Neonatal rotavirus infection in Bangladesh: strain characterization and risk factors for nosocomial infection. *Paediatric Infectious Diseases* **15**, 672–677.
- Krilov, L.R., Barone, S.R., Mandel, F.S., Cusack, T.M., Gaber, D.J. and Rubino, J.R. (1996) Impact of an infection control program in a specialized preschool. *American Journal of Infection Control* **24**, 167–173.
- Krilov, L., Pierik, L., Keller, E., Mahan, K., Watson, D., Hirsch, M., Hamparian, V. and McIntosh, K. (1986) The association of

- rhinoviruses with lower respiratory tract diseases in hospitalized patients. *Journal of Medical Virology* **19**, 345–352.
- Larson, T. and Bryson, Y. (1982) Fomites and herpes simplex virus: the toilet seat revisited. *Paediatric Research* **16**, 244, abstract no. 995.
- LeBaron, C.W., Furutan, N.P., Lew, J.F., Allen, J.R., Gouvae, V., Moe, C. and Monroe, S.S. (1990) Viral agents of gastroenteritis: public health importance and outbreak management. *Morbidity and Mortality Weekly Report* **39**, 1–24.
- Lew, J.F., Moe, C.L., Monroe, S.S., Allen, J.R., Harrison, B.M., Forrester, B.D., Stine, S.E., Woods, P.A., Hierholzer, J.C., Herrmann, J.E., Blacklow, N.R., Bartlett, A.V. and Glass, R.I. (1991) Astrovirus and adenovirus associated with diarrhoea in children in day care settings. *Journal of Infectious Diseases* **164**, 673–678.
- Lewis, D.C., Lightfoot, N.F., Cubitt, W.D. and Wilson, S.A. (1989) Outbreaks of astrovirus type 1 and rotavirus gastroenteritis in a geriatric inpatient population. *Journal of Hospital Infection* **14**, 9–14.
- Lewis, H.M., Parry, J.V. and Davies, H.A. (1979) A year's experience of the rotavirus syndrome and its association with respiratory illness. *Archives of Disease in Childhood* **54**, 125–128.
- Lieberman, J.M. (1994) Rotavirus and other viral causes of gastroenteritis. *Paediatric Annals* **23**, 529–535.
- Marks, P.J., Vipond, I.B., Carlisle, D., Deakin, D., Fey, R.E. and Caul, E.O. (2000) Evidence for airborne transmission of Norwalk-like virus (NLV) in a hotel restaurant. *Epidemiology and Infection* **124**, 481–487.
- Mathias, K.A. (1999) The use of consumer knowledge, beliefs and attitudes in the development of a local authority strategy for domestic food hygiene education. MPhil Thesis. Open University.
- Mbithi, J.N., Springthorpe, V.S., Boulet, J.R. and Sattar, S.A. (1992) Survival of hepatitis A virus on human hands and its transfer on contact with animate and inanimate surfaces. *Journal of Clinical Microbiology* **30**, 757–763.
- Mbithi, J.N., Springthorpe, V.S. and Sattar, S.A. (1991) Effect of relative humidity and air temperature on survival of Hepatitis A virus on environmental surfaces. *Applied and Environmental Microbiology* **57**, 1394–1399.
- McEvoy, M., Blake, W., Brown, D., Green, J. and Cartwright, R. (1996) An outbreak of viral gastroenteritis on a cruise ship. *Communicable Disease Report Review* **6**, R188–R192.
- McGuckin, M., Waterman, R., Porten, L., Bello, S., Caruso, Juzaitis, B., Mazer, S. and Ostrawski, S. (1999) Patient education model for increasing handwashing compliance. *American Journal of Infection Control* **27**, 309–314.
- Melnick, J.L. (1995) History and epidemiology of Hepatitis A virus. *Journal of Infectious Diseases* **171** (Suppl. 1), S2–S8.
- Minor, T.E., Allen, C.I., Tsiatis, A.A., Nelson, D.B. and Alessio, D.J. (1981) Human infective dose determinations for oral poliovirus type 1 vaccine in infants. *Journal of Clinical Microbiology* **13**, 388–389.
- Minuk, G.Y., Ding, L.X., Hannon, C. and Sekla, L. (1994) The risks of transmission of acute hepatitis A and B virus infection in an urban centre. *Journal of Hepatology* **21**, 118–121.
- Mitchell, D.K., Van R., Morrow, A.L., Monroe, S.S., Glass, R.I. and Pickering, L.K. (1993) Outbreaks of astrovirus gastroenteritis in day care centres. *Journal of Paediatrics* **123**, 725–732.
- Morens, D.M. and Rash, V.M. (1995) Lessons from a nursing home outbreak of influenza A. *Infection Control and Hospital Epidemiology* **16**, 275–280.
- Nerurkar, L.S., West, F., May, M., Madden, D.L. and Sever, J.L. (1983) Survival of herpes simplex in water specimens collected from hot tubs in spa facilities and on plastic surfaces. *Journal of the American Medical Association* **22**, 3081–3083.
- Nicholson, K.G. (1996) Impact of influenza and respiratory syncytial virus on mortality in England and Wales from January, 1975 to December, 1990. *Epidemiology and Infection* **116**, 51–63.
- Nicholson, K.G., Kent, J., Hammersley, V. and Cancio, E. (1997) Acute viral infections of upper respiratory tract in elderly people living in the community: comparative, prospective, populations based study of disease burden. *British Medical Journal* **315**, 1060–1064.
- Niffenegger, J.P. (1997) Proper handwashing promotes wellness in child care. *Journal of Paediatric Health Care* **11**, 26–31.
- Nishimura, S., Kagehira, M., Kono, F., Nishimura, M. and Taenaka, N. (1999) Handwashing before entering the intensive care unit: what we learned from continuous video-camera surveillance. *American Journal of Infection Control* **27**, 367–369.
- Osterholm, M.T., Reves, R.R., Murph, J.R. and Pickering, L.K. (1992) Infectious diseases and child day care. *Paediatric Infectious Diseases Journal* **11**, S31–S41.
- Pang, X.L., Honma, S., Nakata, S. and Verikari, T. (2000) Human caliciviruses in acute gastro-enteritis of young children in the community. *Journal of Infectious Diseases* **181**, S288–S294.
- Parashar, U.D., Bresee, J.S., Gentsch, J.R. and Glass, R.I. (1998a) Rotavirus. *Emerging Infectious Diseases* **4**, 561–570.
- Parashar, U.D., Holman, R.C., Clarke, M.J., Bresee, J.S. and Glass, R.I. (1998b) Hospitalizations associated with rotavirus diarrhoea in the US, 1993 through 1995: surveillance based on the new ICD-9-CM rotavirus specific diagnostic code. *Journal of Infectious Diseases* **177**, 7–13.
- Patterson, W., Haswell, P., Fryers, P.T. and Green, J. (1997) Outbreak of small round structured virus gastroenteritis arose after kitchen assistant vomited. *Communicable Disease Report Review* **7**, R101–R103.
- Pickering, L.K., Bartlett, A.V., Reeves, R.R. and Morrow, A. (1988) Asymptomatic excretion of rotavirus before and after rotavirus diarrhoea in children in day care centres. *Journal of Paediatrics* **112**, 361–365.
- Pittet, D., Mourouga, P. and Pereger, T.V. (1999) Compliance with handwashing in a teaching hospital. *Annals of Internal Medicine* **130**, 126–130.
- Reed, S.E. (1975) An investigation on the possible transmission of rhinovirus colds through indirect contact. *Journal of Hygiene (Cambridge)* **75**, 249–258.
- Rheinbaben, F.V., Schünemann, S., Broß, T. and Wolff, M.H. (2000) Transmission of viruses via contact in a household setting: experiments using bacteriophage ϕ X174 as a model virus. *Journal of Hospital Infection* **46**, 61–66.
- Richardson, S., Grimwood, K., Gorrell, R., Palombo, E., Barnes, G. and Bishop, R. (1998) Extended excretion of rotavirus after severe diarrhoea in young children. *Lancet* **351**, 1844–1848.
- Roberts, L., Jorm, L., Patel, M., Smith, W., Douglas, R.M. and McGilchrist, C. (2000a) Effect of infection control measures on the frequency of upper respiratory infection in child care: a randomised, controlled trial. *Paediatrics* **105**, 738–742.
- Roberts, L., Jorm, L., Patel, M., Smith, W., Douglas, R.M. and McGilchrist, C. (2000b) Effect of infection control measures on the

- frequency of diarrhoeal episodes in child are: a randomised, controlled trial. *Paediatrics* **105**, 743–746.
- Rodriguez, W.J., Kim, H.W., Brandt, C.D., Schwartz, R.H., Gardner, M.K., Jeffries, B., Parrott, R.H., Kaslow, R.A., Smith, J.L. and Kapikain, A.Z. (1987) Longitudinal study of rotavirus infection and gastroenteritis in families served by a paediatric medical practice: clinical and epidemiological observations. *Paediatric Infectious Diseases* **6**, 170–176.
- Rodriguez, W.J., Kim, H.W., Brandt, C.D., Yolken, R., Richard, M., Arrobio, J.O., Schwartz, R.H., Kapikian, A.Z., Chanock, R.M. and Parrott, R.H. (1979) Common exposure outbreak of gastroenteritis due to type 2 rotavirus with high secondary attack rates within families. *Journal of Infectious Diseases* **140**, 353–357.
- Rotter, M.L. (1997) Handwashing and hand disinfection. In *Prevention and Control of Nosocomial Infections* ed. Wenzel, R.P. pp. 1052–1068. Philadelphia: Lippincott Williams & Wilkins.
- Ryan, M.J., Ramsey, M.E., Brown, D., Gay, N.J., Farrington, C.P. and Wall, P.G. (1996) Hospital admissions attributable to rotavirus infection in England and Wales. *Journal of Infectious Diseases* **174**, S12–S18.
- Ryan, M.J., Wall, P.G., Adak, G.K., Evans, H.S. and Cowden, J.M. (1997) Outbreaks of infectious intestinal disease in residential institutions in England and Wales 1992–4. *Journal of Infection* **34**, 49–54.
- Samadi, A.R., Huq, M.I. and Ahmed, Q.S. (1983) Detection of rotavirus in handwashings of attendants of children with diarrhoea. *British Medical Journal* **286**, 188.
- Sattar, S.A., Jacobsen, H., Springthorpe, S., Cusack, T. and Rubino, J. (1993) Chemical disinfection to interrupt the transfer of Rhinovirus type 14 from environmental surfaces to hands. *Applied and Environmental Microbiology* **59**, 1579–1585.
- Sattar, S.A., Lloyd-Evans, N. and Springthorpe, V.S. (1986) Institutional outbreaks of rotavirus diarrhoea: potential role of fomites and environmental surfaces as vehicles for virus transmission. *Journal of Hygiene* **96**, 277–289.
- Sattar, S.A. and Springthorpe, S. (1999) Viricidal activity of biocides. Activity against human viruses. In *Principles and Practice of Disinfection, Preservation and Sterilization*, 3rd edn. eds Russell, A.D., Hugo, W.B. and Ayliffe, G.A.J. pp. 168–186. Oxford: Blackwell Science.
- Schurmann, W. and Eggers, H.J. (1985) An experimental study on the epidemiology of enteroviruses — water and soap washing of poliovirus 1 contaminated hands, its effectiveness and kinetics. *Medical Microbiology and Immunology* **174**, 221–236.
- Scott, E. (1999) Hygiene issues in the home. *American Journal of Infection Control* **27**, S22–S25.
- Simoes, E.A.F. (1999) Respiratory syncytial virus infection. *Lancet* **354**, 847–852.
- Smith, C.B., Purcell, R.H., Bellanti, J.A. and Chanock, R.M. (1966) Protective effect of antibody to parainfluenza type 1 virus. *New England Journal of Medicine* **275**, 1145–1149.
- Soule, H., Genoulaz, O., Gratacap-Cavallier, B., Mallaret, M.R., Morand, P., Francois, P., Bin, D.L.D., Charvier, A., Bost-Bru, C. and Seigneurin, J.-M. (1999) Monitoring rotavirus environmental contamination in a paediatric unit using polymerase chain reaction. *Infection Control and Hospital Epidemiology* **20**, 432–434.
- Sperber, S.J. (1994) The common cold. *Medicine* **11**, 235–242.
- St. Sauver, J., Khurana, M., Kao, A. and Foxman, B. (1998) Hygienic practices and acute respiratory illness in family and group day care homes. *Public Health Reports* **113**, 544–551.
- Stevens, R.A. and Holah, J.T. (1993) The effect of wiping and spray-wash temperature on bacterial retention on abraded domestic sink surfaces. *Journal of Applied Bacteriology* **75**, 91–94.
- Sundkvist, T., Hamilton, G.R., Hourihan, B.M. and Hart, I.J. (2000) Outbreak of hepatitis A spread by contaminated drinking glasses in a public house. *Communicable Disease and Public Health* **3**, 60–62.
- Thompson, S.C. (1994) Infectious diarrhoea in children: controlling transmission in the child care setting. *Journal of Paediatrics and Child Health* **30**, 210–219.
- Uhari, M. and Mottonen, M. (1999) An open randomized controlled trial of infection prevention in child day-care centres. *Journal of Paediatrics and Infectious Diseases* **18**, 672–677.
- Valenti, W.M. (1998) Selected viruses of nosocomial importance. In *Hospital Infections*, 4th edn. eds Bennett, J.V. and Brachman, P.S. pp. 637–664. Philadelphia: Lippincott-Raven.
- Ward, R., Bernstein, D., Knowlton, D., Sherwood, J., Yung, E., Cusack, T., Rubino, J. and Schiff, G. (1991) Prevention of surface-to-human transmission of rotavirus by treatment with disinfectant spray. *Journal of Clinical Microbiology* **29**, 1991–1996.
- Ward, R.L., Bernstein, D.I., Young, E.C., Sherwood, J.R., Knowlton, D.R. and Schiff, G.M. (1986) Human rotavirus studies in volunteers: determination of infectious dose and serological response to infection. *Journal of Infectious Disease* **154**, 871–880.
- Wheeler, J.G., Sethi, D., Cowden, J.M., Wall, P.G., Rodrigues, L.C., Tompkins, D.S., Hudson, M.J. and Roderick, P.J. (1999) Study of infectious intestinal disease in England: rates in the community, presenting to general practice and reported to national surveillance. *British Medical Journal* **318**, 1046–1055.
- Wilde, J.R., Pickering, L., Eiden, J. and Yolken, R. (1992) Detection of rotaviruses in the day care environment by reverse transcriptase polymerase chain reaction. *Journal of Infectious Disease* **166**, 507–511.