



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

VIRUS 00954

Heterogeneity of the L2 gene of field isolates of bluetongue virus serotype 17 from the San Joaquin Valley of California

Carlos A. de Mattos, Cecilia C.P. de Mattos, Bennie I. Osburn
and N. James MacLachlan *

*Department of Veterinary Pathology, School of Veterinary Medicine, University of California,
Davis, CA 95616, USA*

(Received 27 May 1993; revision received 16 August 1993; accepted 30 August 1993)

Summary

Genome segment 2 (L2) from six field isolates of bluetongue virus (BTV) serotype 17 was sequenced by cycling sequencing after the amplification of the viral cDNA by the polymerase chain reaction. The viruses were isolated from sheep, cattle and a goat in the San Joaquin Valley of California during the years 1981 and 1990. These viruses exhibit divergent patterns of neutralization with BTV 17-specific monoclonal antibodies. The six L2 genes of the BTV 17 field isolates all encode a protein of 955 amino acids. Similarity of the nucleotide sequences of the L2 genes with respect to the prototype strain ranges between 93.8% and 95.1%, whereas the similarity between the field isolates ranges from 96.8% to 99.1%. Although very closely related, the L2 gene of each virus is distinct. Furthermore, mutations in the L2 gene of field isolates of BTV do not consistently follow a linear pattern of accumulation over time. Some amino acid changes in the VP2 protein of field strains were conserved over time, whereas others were not correlated with the year of isolation and some substitutions were unique to individual viruses. The predicted VP2s constitute a group of non-identical, but closely related proteins. Phylogenetic analyses suggest that the viral variants which co-circulate in the San Joaquin Valley could evolve by different evolutionary pathways.

Bluetongue virus; Sequencing; Evolution

* Corresponding author.

Introduction

A variety of RNA viruses such as influenza virus, foot and mouth disease virus, rhabdoviruses and some alphaviruses are characterized by extensive genetic polymorphism (Domingo et al., 1980; Nichol, 1988; Rico-Hesse et al., 1988; Webster et al., 1992). This genetic diversity is responsible for differences in antigenic properties, tissue tropism and virulence exhibited by individual members of the same virus group or serotype (Wege et al., 1981; Faragher et al., 1988). Even viruses isolated at the same time and from a common geographical region may be genetically heterogenous, as may RNA viruses isolated from a single outbreak of disease or even from a single infected host (Sobrino et al., 1986; Domingo, 1989; Kinnunen et al., 1990).

Bluetongue virus (BTV), the prototype member of the genus Orbivirus in the Reoviridae family, is also genetically heterogenous (Roy, 1989; Heidner et al., 1991). Infection of wild and domestic ruminants occurs throughout much of the tropical and temperate areas of the world, likely wherever competent vector gnats occur. Of the 24 known serotypes of BTV only five, serotypes 2, 10, 11, 13 and 17, have been isolated in the US since 1953. BTV has 10 viral dsRNA genome segments which encode at least 10 viral polypeptides, seven of which are structural and three nonstructural. The structural proteins form a two-layered particle which consists of a core constituted by VP1, VP3, VP4, VP6 and VP7, and an outer capsid formed by VP2 and VP5 which are encoded by genome segments 2 (L2) and 5 (M2) respectively (Roy, 1989). VP2 is responsible for neutralization and serotype specificity of BTV (Roy, 1989), although VP5 might affect neutralization by its conformational influence on VP2 (Cowley and Gorman, 1989; Mertens et al., 1989). The genetic diversity of the BTV serogroup is clearly established (Roy, 1989; Heidner et al., 1991; de Mattos et al., 1992) with gene segments L2 and M2 being the most divergent amongst the prototype viruses of different BTV serotypes (Roy, 1989; Heidner et al., 1991). The genetic diversity that characterizes this serogroup is the result of both genetic drift from individual point mutations, and reassortment of individual viral gene segments (Roy, 1989; de Mattos et al., 1991). The genetic variability of the BTV serogroup is reflected in its antigenic diversity. The antigenic diversity of US BTV prototype and field isolates has previously been characterized with neutralizing monoclonal antibodies (Appleton and Letchworth, 1983; Greider and Schultz, 1990; MacLachlan et al., 1992; Rossitto and MacLachlan, 1992). Phenotypic variation is specially pronounced amongst isolates of serotype 17 (Appleton and Letchworth, 1983; MacLachlan et al., 1992; Rossitto and MacLachlan, 1992).

Phylogenetic studies have helped to clarify the evolution and epidemiology of viral infections such as influenza and Venezuelan equine encephalitis (Yamashita et al., 1988; Weaver et al., 1992). There is less information available regarding the evolution of the BTV serogroup. Evolutionary relationships among prototype and some field strains of different BTV serotypes have been characterized using the complete or partial nucleotide sequences of conserved or variable genes (Gould and Pritchard, 1990, 1991; Kowalik and Li, 1991; Hwang et al., 1992a,b; Yang et

al., 1992). Field isolates of the same BTV serotype obtained from different host species in the same or different years and from the same geographical region have yet to be adequately compared by sequence analysis. The purpose of this study was to further characterize the L2 gene of phenotypically distinct field isolates of BTV serotype 17 isolated in the San Joaquin Valley of California in 1981 and 1990, and to evaluate the evolutionary relationship of these field isolates to each other and with the five US prototype strains of BTV serotypes 2, 10, 11, 13 and 17.

Materials and Methods

Viruses and RNA purification

The L2 genes of six independently isolated strains of BTV 17 obtained from ruminants in the San Joaquin Valley of California during the years 1981 and 1990 were characterized. The origin of each isolate is listed in Table 1. The isolates were processed as described (Heidner et al., 1991; MacLachlan et al., 1992). The viruses were propagated in BHK 21 cells (Heidner et al., 1991) and the viral dsRNA was phenol extracted, ethanol precipitated and lithium chloride purified (de Mattos et al., 1989).

PCR amplification and cycling sequencing

Fifteen synthetic 22-mer and one 19-mer oligonucleotide primers were constructed from the published sequence of the L2 gene of BTV 17 (Ghiashi et al., 1987). These correspond to nucleotide positions (Fig. 1): 1–19, 230–251, 338–359, 661–682, 921–942, 1000–1021, 1262–1283, 1519–1540, 1631–1652, 1922–1943, 1949–1970, 2148–2169, 2316–2337, 2494–2515, 2731–2752, 2902–2923. Different combinations of these primers were used to reverse transcribe the positive and negative strand of viral nucleic acid using mouse Moloney leukemia virus reverse transcriptase (M-MuLV) (Akita et al., 1992). cDNAs were amplified using the polymerase chain reaction as described by Akita et al. (1992). The annealing temperature for the PCR amplification was adapted according to the characteristics of each primer pair used, ranging from 55°C to 59°C. The PCR products were

TABLE 1
Bluetongue virus serotype 17 field isolates studied

Field isolate	Species of origin	Year of isolation	County of isolation	State
17B81	Bovine	1981	Fresno	CA
17C81	Caprine	1981	Kern ^a	CA
17O81	Ovine	1981	Kern ^a	CA
17B90Z	Bovine	1990	Fresno	CA
17O90X	Ovine	1990	Kern	CA
17O90Y	Ovine	1990	San Joaquin	CA

^a Field isolates obtained from the same farm.

purified using a commercial kit (Gene-Clean, BIO 101). Both strands of the cDNAs were sequenced by cycle sequencing using a commercial kit (f-mol Kit, Promega). The sequencing reaction was done with 500 ng of cDNA, 20 pM of primer, an initial 2 min incubation at 95°C, followed by 30 cycles of 95°C for 1 min, 57°C for 1 min and 70°C for 1 min. A final extension step, using terminal deoxynucleotidyl transferase, was done at 37°C for 60 min. Each sequencing

		70
BTV17	<u>GTTAAAAGACTGATCCACCATGGAGGAGTTCGTCA</u> TTCCAGTCTATTCA <u>GAGATGAGATTCCATATGCG</u>	
17B81	C..
17C81
17081	C..
17B90Z	G..C..
17090X	G..C..
17090Y	G..C..
	71	140
BTV17	TTACTAACGAGATA <u>CCCTCTGGCGATA</u> CAGACAA <u>ATGTTAAAATAGAGGATGTTGAAGGAAACATAATG</u>	
17B81G.....T.....	G.....
17C81A.....	G.....
17081G.....T.....	G.....
17B90Z	..G..G.....T.....	C.....G.....
17090X	..G..G.....T.....	C.....G.....
17090Y	..G..G.....T.....	C.....G.....
	141	210
BTV17	TTGTAA <u>AGATTCC</u> TGAATCCGATATGATAGATATACCAAA <u>ATTAA</u> CGATTGTAGAGGCCATGAATTATAA	
17B81G.....	G.....GA.....
17C81G.....	G.....A.....
17081G.....	G.....A.....
17B90Z	G.....A.....
17090X	G.....A.....
17090Y	G.....A.....
	211	280
BTV17	ACCAGCGAGAA <u>ACGATGGGATCGTGTACCTAGATTACTTGATATAACATTAC</u> CGCCTTATGATGATPAGA	
17B81A.....	T.....
17C81A.....	T.....
17081	C.....A.....	T.....
17B90ZA.....	C.....T.....
17090XA.....	C.....T.....
17090Y	C.....A.....	C.....T.....
	281	350
BTV17	AAATCGAC <u>AAAAGCGCTCGAGGGATAGAGTTCATGACGAACGCTAGATGGATGAAATGGGCCATAGATG</u>	
17B81T.....
17C81	...A...A...T.....
17081	...A...A...T.....
17B90Z	...A.....	T.....
17090X	...A.....	T..C.....
17090Y	...A.....	T.....
	351	420
BTV17	ATAGA <u>ATGGATATCCAACCTCTAAAGTGACTT</u> AGATCATTATTGTTCCGTCAACC <u>ATCAACTCTTAA</u>	
17B81G.....	C.....G.....
17C81G.....	C.....G.....
17081G.....	C.....G.....
17B90ZG.....	C.....T.....G.....
17090XG.....	C.....T.....G.....
17090YG.....	C.....T.....G.....

Fig. 1. Nucleotide sequences of the plus-strand of the L2 gene of the US prototype and field isolates of BTV 17. The start and stop codons are underlined and bases identical to the prototype are represented by dots. The arrows indicate the examples described in the text.

reaction was repeated with PCR products from several different amplifications to control for artifacts.

Computer analyses

The nucleotide sequence of the L2 gene as well as the predicted amino acid sequences of VP2 of each of the six field isolates and the five US BTV prototype strains were analyzed using the Wisconsin Package (Genetic Computer Group Inc., 1991). Phylogenetic analyses were done with the PHYLIP package version 3.41

	421	490
BTV17	CTCGTTGTTAAAGCGAATGCTGCCAACGCTGATACGATCTATTGATTATTCCCGCTTGAGGACTAT	
17B81C..C.....C.....	A.....A..C..
17C81C..C.....C.....	A.....A..C..
17081C..C.....C.....	A.....A..C..
17B90ZC..C.....	A.....A..C..
17090XC..C.....	A.....A..C..
17090YC..C.....	A.....A..C..
	491	560
BTV17	AAAAAGAGATGTAACCATAACAAATCTTGTATTATTGAGAAAGTTAACGAATATGGAGTTGTTCCACGCGT	
17B81C.....	G..T..
17C81GA.....C.....	G..T..
17081GA.....C.....	G..T..
17B90ZC.....	G..T..
17090XC.....	G..T..
17090YC.....	G..T..
	561	630
BTV17	TGCAAGGTGCTGCGTACAGTATCAAATCAAGCTACGAATTAGTGGCATACTCGAAAGAGGAAGGCCTGGA	
17B81A.....	A.....A..T..
17C81A.....	A.....T..
17081A.....	A.....T..
17B90ZA.....T.....	A.....A..TT..
17090XA.....T.....	A.....A..TT..
17090YA.....T.....	A.....A..TT..
	631	700
BTV17	GGGACTTATGTGGTAGGACAGCCAAATGGATAACACTGACTAGAGGAACGCGAATAGGCAATAGTGGAA	
17B81C..G.....G.....	
17C81C..A.....G.....	
17081C..A.....G.....	
17B90ZCA.....	
17090XCA.....	
17090YCA.....	
	701	770
BTV17	TTATCCTATGAACGGTCATTCGAGCATGGTCCAGGTAAAGCGTAAATGGAAAGATTCCAGACGAGATAG	
17B81	...A.....	G.T.....AG..T..
17C81	C..C.T.....T..C..A.....	G.T.....AG..
17081	C..C.T.....T..C..A.....	G.T.....AG..
17B90Z	...C.T.....T..C..A.....	G..TT.....AG..
17090X	...C.T.....T..C..A.....	G..TT.....AG..
17090Y	...C.T.....T..C..A.....	G..TT.....AG..
	771	840
BTV17	CGAACGAGATCGCGCAACTGAATAGAATAAGGGCAGAGTGGATAACAGCCACATACGATAGAGGCAGGAT	
17B81G.....	G.G..T..
17C81G.....	G.G..T..
17081G.....	G.G..T..
17B90ZG.....	G.G..T..
17090X	...T..A.....	G.G..T..
17090Y	...G.....	G.G..T..

Fig. 1 (continued).

(Felsenstein, 1991). The first 19 nucleotides of the 5' end and the last 22 of the 3' end were not included in the phylogenetic studies because primers were used to amplify the gene. The genetic distance between any two nucleotide sequences was calculated using the DNADIST program based in the Kimura two parameter model (Kimura, 1980). The phylogenetic analyses were carried out utilizing a distance matrix method, based on the least-square method (Fitch and Margoliash, 1967), and implemented in the FITCH program of the PHYLIP package. This procedure makes no assumptions about the relative rates of evolution in all

	841	910
BTV17	TAGAGCGCTAGAGCTATGCAGTATCCTTTCCACGATTGGCGTAAGATGCTAAATACGCATGAGGAGCCG	
17B81A.....	A.G.C.....A.....
17C81AG.....	A.G.....
17O81AG.....	A.G.C.....A.....
17B90ZAG.....	G.C.....A.A.....
17O90XAG.....	G.C.....A.A.....
17O90YAG.....	G.C.....A.A.....
	911	980
BTV17	AAAGATGAAATGGACCTATCGACGAGATTCAAGTTAACCTTGATGAGAAATTAAACAGAGCGGATTCA	
17B81T.....A.....C.....C.A.....	A.C.....
17C81T.....A.A.....C.....A.A.....	A.C.....
17O81T.....A.A.....C.....C.A.....	A.C.....
17B90ZT.....A.A.....C.....C.A.....	A.C.....
17O90XT.....A.A.....C.....C.A.....	A.C.....
17O90YT.....A.A.....C.....C.A.....	A.C.....
	981	1050
BTV17	AAACATGTTAATATTTGGTGTAAAGAGGTCCAGCACAGATGAAGGAAGATTCTACGCTCTGATCGCAAT	
17B81	T.T.....T.....
17C81	T.....T.....
17O81CC.....G.....	T.....T.....
17B90ZCC.....G.....	T.....T.....
17O90XCC.....G.....	T.....T.....
17O90Y	T.....T.....
	1051	1120
BTV17	CGCAGCGACAGATACACAAAAGGGCAGAGTGTGGAGAACGAATCCGTATCCATGCTTGCAGGGTGCTTTA	
17B81G.....T.....A.....	A.....
17C81G.....T.....A.....	A.....
17O81T.....A.....	A.....
17B90ZG.....T.....A.....	A.....
17O90XG.....T.....A.....	A.....
17O90YG.....T.....A.....	A.....
	1121	1190
BTV17	GTTGCAGCTGACTGTGAATTAGGTGATGTTACAGTACACTTCGACGTGTGTACATGGAGTCTAAGGC	
17B81C.....	G.....
17C81C.....G.C.....C.G.....	C.G.....
17O81C.....G.C.....C.....	C.....
17B90ZC.....C.....	G.....
17O90XC.....C.....A.G.....T.....	A.G.....T.....
17O90YC.....C.....	G.....
	1191	1260
BTV17	CAGAATATGGACAGCAGCAGCAGTTAGAGAACATAAAACGTCTTAATCGTATAAATTATTGAA	
17B81	.G..G.....A.....	T.....C.....
17C81	.G..G.....G.....	
17O81	.G..G.....G.....	
17B90Z	.G..G.....	T.....
17O90X	.G..G.....	
17O90Y	.G..G.....	T.....

Fig. 1 (continued).

branches. The L2 gene from epizootic hemorrhagic disease virus (EHDV) serotype 1 was used as an outgroup for the phylogenetic analyses because it is distantly related to the homologous genome segment from the BTV serogroup and it is clearly recognized as a distinct orbivirus serogroup (Gould and Pritchard, 1991; Iwata et al., 1992). Previous studies on orbivirus phylogeny have placed EHDV-1 outside the BTV serogroup (Gould and Pritchard, 1991; Iwata et al., 1992; Roy,

	1261	1330
BTW17	TTCAAACCTAGCAGTTGGTGACCAGATTATTCAATTGGCGTTATGAGGTTAAAGCATCGCGGAGACGACC	
17B81G..C..C.....A.....T
17C81G..C..C.....AT
17081G..C..C.....A.....AT
17B90ZG..C..C.....A.....T
17090XG..C..C.....A.....T
17090YG..C..C.....A.....T
	1331	1400
BTW17	TATGACAGCGGATATAATGTGTCGGCATGAAGCTGAGGAGGATGAACATTATGTAAAATTAAATGAGGACA	
17B81C.....C.....
17C81A.....C.....A.....G.....G.....C.....C.....
17081A.....C.....A.....G.....G.....C.....C.....
17B90ZT.....C.....C.....C.....
17090XT.....C.....G.....C.....
17090YT.....C.....G.....C.....
	1401	1470
BTW17	AATATAAAGAAATGCTGGACAGAACATGATTCAAGGGTGGGTGGGATCAGGAAAGATTAAACTTCATAACAT	
17B81C.....	
17C81C.....	
17081C.....	
17B90ZG.....G.....	
17090XC.....	
17090YC.....	
	1471	1540
BTW17	ACTGACGGACCCAAACCTTATTGACGATTGATTTGAAAAAGACGCGTATCTGAATTCA CGGTCCGAGTTA	
17B81T.....T.....
17C81T.....T.....
17081T.....T.....
17B90ZT.....T.....T.....
17090XT.....T.....T.....
17090YT.....T.....C.....T.....T.....
	1541	1610
BTW17	GTTTGCCGGATTATTCGACAAATGGATCAGTTCACCAATGTTAACCGCGCTTAAGAATTACTAAAG	
17B81T.....	
17C81	
17081	
17B90ZA.....
17090XA.....
17090Y	
	1611	1680
BTW17	GGGAGATCGGAACATCAAAAAGGATGACCCATGGAACAACCGCGCGTACGTGGATACTTAAGCCCT	
17B81G.....T.....A.....T.....
17C81G.....T.....A.....C.....T.....
17081G.....T.....A.....C.....T.....C
17B90ZGT.....G.....T.....A.....C.G.T.T.C
17090XGT.....G.....T.....A.....C.....T.....C
17090YGT.....G.....T.....A.....C.....T.....C

Fig. 1 (continued).

1992). Although EHDV was first isolated in 1955, historical data indicate that the virus has been present in the US at least since 1890 (Shope et al., 1960). A bootstrap analysis with 100 replicates was done to assess confidence limits of the branch pattern. The results are presented as a majority rule consensus tree which shows the most frequent branching order (Felsenstein, 1985). A value > 95% is considered significant.

	1681	1750
BTV17	TGCGGAATCGTTAGATTTGTTCTGGGCCACTACGATCTGGCTTCTATTTTGACGAGACGTTG	
17B81G.....G.....T.....A.....G.....	
17C81G.....G.....T.....A.....G.....G.....	
17O81G.....G.....T.....A.....G.....	
17B90ZG.....G.....T.....A.....G.....G.....	
17O90XG.....G.....T.....A.....G.....G.....	
17O90YG.....G.....T.....A.....G.....G.....	
	1751	1820
BTV17	AGCTAAACAGGAACAGTCGCCGTTTTCAATATTGAGTCAGCTCGATTTCCCGCCTACGC	
17B81A.....	
17C81A.....C.....	
17O81C.....	
17B90ZA.....	
17O90XA.....	
17O90YA.....	
	1821	1890
BTV17	AGCTAACAGGAGATGCCGTATGCCACATTCAAGCGGAGCGTTATACGTTAGGAAAGTCGCGCTATT	
17B81C.....	
17C81C.....C.....	
17O81C.....C.....	
17B90ZC.....C.....	
17O90XC.....C.....	
17O90YC.....C.....	
	1891	1960
BTV17	TTTGATCGGAATTATGAAAAGTTAACGATCTACATGAAGGTATGGAACACCAGAGATATGTGCAT	
17B81	...A.....	T..AGC.....
17C81	...A.....A.....	T..A.C.....
17O81	...A.....A.....	T..A.C.....
17B90Z	...A.....	T..C.....
17O90X	...A.....	T..C.....
17O90Y	T..C.....
	1961	2030
BTV17	CCGTCGACTGGCGGACGTATCAGAACCGTGCTAGAGATGAAGGATTCCCTGCCAGCTTACGTGTTTG	
17B81T.....A..T.....	CC..T..T..A..C..T..C..
17C81T.....T.....	CC..T..T..A.....C..C..
17O81T.....T.....	CC..T..T..A.....C..C..
17B90ZT.....T.....	CC..T..T..A.....T..C..
17O90XT.....T.....	CC..T..T..A.....T..C..
17O90YC..T.....T.....	CC..T..T..A.....T..C..
	2031	2100
BTV17	TGATTGATTACATCTTGAAAAACGTAAACAGCTACGTGATACCAAAGAGGCGAGGTACATTGTGTATCT	
17B81	C.....
17C81	C.....
17O81	C.....
17B90ZG..G.....	A.....G.C.....T..
17O90XG..G.....	A.....C.....T..
17O90YC..G.....	A.....G.C.....T..

Fig. 1 (continued).

Results

L2 nucleotide sequence comparison of BTV 17 field isolates

The six L2 genes of the BTV 17 field isolates all are 2923 nucleotides in length, and each encodes a protein of 955 amino acids. The initiation codons are located at position 20–22 and the termination codons at position 2885–2887 (Fig. 1). Nucleotide sequences of the BTV 17 genomes were compared using the sequence of the prototype strain of BTV 17 as reference (Ghiasi et al., 1987). There are no

	2101	2170
BTW17	AATTCAAAGTCTTACTGGGACACAACGGCTGAGTGTAAAGTCGACGTTCCGAATTCTTCCAACGA	
17B81C.....GA.....C.G.AA.....C.....T.....	
17C81C.....GA.....C.G.AA.....C.....T.....	
17081C.....GA.....C.G.AA.....	
17B90ZC.....GA.....C.G.AA.....	
17090XC.....T.GA.....C.G.AA.....C.....T.....	
17090YC.....GA.....C.G.AA.....C.....T.....G..	
	2171	2240
BTW17	TTGTPGATGCTAAAAGAGATCAAATTGTGCCGATCTGAATGTGATTAACCTCCTCCCTGTATGTTCC	
17B81	..A..A.....G.....T.....C.T.....	
17C81	..A..A.....G.....T.....T.....C.....	
17081	..A..A.....G.....T.....T.....C.....	
17B90Z	..A..A.C.T.....G.....T.....C.....	
17090X	..A..A.C.T.....G.....T.....C.....	
17090Y	..A..A.C.T.....G.....T.....C.....	
	2241	2310
BTW17	TTGTCATGATAACATCTCGTATTGGCATAGACAGTGGTCGATTCCAATGGTACTGTTGACGATACGGAT	
17B81C.....A.....G.....	
17C81	...C.....A.....	
17081C.....A.....	
17B90ZT.....C.....A.....	
17090XT.....C.....A.....	
17090YT.....C.....A.....	
	2311	2380
BTW17	CAAGTTAACCTGTAGAGGTTGGTCATATGCGAATAGATTGGTTCAAAAGTTTATGAACCTTACA	
17B81C.....	
17C81	
17081C.....C.G.....A.....	
17B90Z	..G.....C.....C.G.CAA.....A.....T.....	
17090X	..G.....C.....C.G.A.....A.....T.....	
17090Y	..G.....C.....C.G.A.....A.....T.....	
	2381	2450
BTW17	CGGTTTACCCCTGGTAGTTGAGAAAAACAGATTGCCGAAGATATAACATAAAGAGTTGGTGTGGTCG	
17B81CA.....G.....G.....A.....	
17C81CA.....G.....G.....G.....G.....	
17081CA.....G.....G.....G.....G.....	
17B90ZCA.....G.....G.....G.....A.....	
17090XCA.....G.....G.....G.....A.....	
17090YCA.....G.....G.....G.....C.....	
	2451	2520
BTW17	CTTCGAATATTACACCAATACAAAAATTCCAGGGAAATGTGCATACACCAAGTAATGACTACGAAAAT	
17B81G.....C.....	
17C81G.....C.....	
17081G.....C.....	
17B90ZT.....G.....C.....G.....	
17090XT.....G.....C.....G.....	
17090YT.....G.....C.....G.....	

Fig. 1 (continued).

deletions or insertions in any of the field isolates with respect to the prototype strain. Many of the mutations of the L2 gene of BTV 17 field isolates are silent. Sequence differences amongst the L2 genes most often are due to transitional mutations. Similarity of the nucleotide sequences of the L2 gene of BTV 17 viruses ranges between 93.8% and 95.1% (Table 2). The L2 genes of the 1981 viruses differ from one another in 25–71 nucleotides and thus are more closely related to each other than they are to the prototype from which they differ in 140–164

	2521	2590
BTW17	GGATGTATTGAGAGTACATTTATCTTCCTTATGTGCAGGTCTGGCGGACTCAGTCGTATATAACATTACCA	
17B81A.A.....G.....G.....T..TA.....	
17C81AGA.....G.....T..TA.....	
17O81AGA.....G.....T..TA.....	
17B90ZA.A.....G.....T..TA.....	
17O90XA.A.....G.....T..TA.....	
17O90YA.A.....G.....T..TA.....	
	2591	2660
BTW17	GTTGCGCATCCTAAAAATGCATCGTTCTAAATAATTGTGGGAGACGACAAATTGGAACCGCATACCGCTT	
17B81G.....T.....	
17C81G.....T.....	
17O81G.....T.....	
17B90ZG.....T.....	
17O90XG.....T.....	
17O90YT.....G..G.....T.....	
	2661	2730
BTW17	CAGAACAGATAGTTAGTCGGTATAATTACTCACGTAAGCACATTGTGGAATTGTATCCGTCACTATCGG	
17B81A.....T.C.....CG.....	
17C81A.....CG.....	
17O81A.....CG.....	
17B90ZA.....CG.....	
17O90XA.....A..CG.....	
17O90YA.....CG.....	
	2731	2800
BTW17	GCAGAACAGTCAGTTGAGAGTTCATACATCTGGAATTGTAAAACACCGTGTATGCGACAAGTTCATTCTA	
17B81T.....C.....T.....	
17C81T.....C.....T.....	
17O81T.....C.....T.....	
17B90ZT.....C.....T.....	
17O90XT.....C.....T.....	
17O90YT.....T.....	
	2801	2870
BTW17	AAACACAAAGTCAAGGTGATATTAGTGAGGATGCCGGGTACGTTTCGAAATGATGAATTAAATGACGA	
17B81	
17C81T.....	
17O81A.....	
17B90Z	
17O90X	
17O90Y	
	2871	2923
BTW17	AACTATTGAATGT <u>CTAGGT</u> CCTATGACACGGACCGATGCCCTCTTACACTTAC	
17B81	
17C81	
17O81	
17B90Z	
17O90X	
17O90Y	

Fig. 1 (continued).

TABLE 2

Bluetongue virus 17 field isolates: nucleotide and amino acid sequence comparison

	BTV 17 ^a	17B81 ^b	17C81 ^b	17O81 ^b	17B90Z ^b	17O90X ^b	17O90Y ^b
BTV 17	—	140 ^c (95.1) ^d	159 (94.5)	164 (94.3)	178 (93.8)	181 (93.8)	179 (93.9)
17B81	33 ^e (96.44) ^f	—	69 (97.6)	71 (97.6)	91 (96.9)	90 (96.8)	89 (96.9)
17C81	45 (95.3)	25 (97.4)	—	25 (99.1)	88 (97)	89 (96.9)	85 (97.1)
17O81	42 (95.6)	25 (97.4)	9 (99.6)	—	74 (97.5)	78 (97.3)	76 (97.3)
17B90Z	43 (95.5)	26 (97.3)	28 (97.1)	25 (97.4)	—	28 (99)	26 (99.1)
17O90X	42 (95.6)	24 (97.5)	27 (97.1)	24 (97.5)	8 (99.2)	—	34 (98.8)
17O90Y	47 (95.1)	28 (97.1)	30 (96.9)	27 (97.2)	12 (98.7)	14 (98.5)	—

^a BTV 17 prototype virus.^b BTV 17 field isolates.Upper section: ^c number of nucleotide changes; ^d percentage of nucleotide sequence homology.Lower section: ^e number of amino acid changes; ^f percentage of amino acid sequence identity.

nucleotides. Similarly, the L2 genes of 1990 viruses differ from one another in only 26–34 nucleotides whereas they differ from the prototype in 178–181 nucleotides (Table 2). The viruses isolated in 1981 and 1990 also are more closely related to each other than they are to the prototype, however they differ from one another in 74–91 nucleotides despite the fact they were isolated from the same geographical area at an interval of only 9 years (Table 2). Some mutations were unique to a single virus (e.g., nucleotide 91 in 17C81 and 325 in 17O90X), whereas others were common to all field isolates (e.g., positions 180, 958 and 2142). Mutations sometimes were common to all 1981 (e.g., positions 132 and 658) or all 1990 isolates (e.g., positions 403 and 1486), but were unique to the isolates of that year. Some mutations were present in two of the 1981 (e.g., position 167 in 17C81 and 17O81) or 1990 (e.g., position 1599 in 17B90Z and 17O90X) viruses, while the other field strains had the same nucleotide as the prototype. Mutations which are common to all the field isolates suggest an ancestor-descendant relationship but other mutations clearly are not consistent with this hypothesis. Specifically, mutations in the 1981 viruses were not consistently present in the 1990 isolates and some mutations were unique to individual viruses regardless of year of isolation, which indicates that these mutations are not cumulative. Comparison of the nucleotide sequences of the BTV 17 prototype and field isolates indicates that although very closely related, the L2 gene of each virus is distinct. Furthermore, mutations in the L2 gene of field isolates of BTV do not necessarily follow a linear pattern of accumulation over time.

Predicted VP2 amino acid sequence comparison

The deduced amino acid sequence of VP2 of the six field isolates and the BTV 17 prototype strain are shown in Fig. 2. The number of amino acid substitutions in the VP2 of the six field isolates ranges from 33 in 17B81 to 47 for 17O90Y (Table 2). Mutations were randomly distributed throughout VP2, although extensive regions of homology occur at amino acid 1–37, 90–156, 397–436, 604–643 and positions 903–955 where only VP2 from 17C81 has a tryptophan at position 940

instead of the glycine present in all other proteins (Fig. 2). These regions might be conserved because they are critical to maintain the function of VP2.

Some amino acids are unique to the prototype strain (e.g., position 54), or to a particular field isolate (e.g., 17B81: position 895; 17C81: 742; 17B90Z: 552; 17O90X: 351 and 17O90Y: 677). In contrast, specific amino acids were shared by the prototype and all the 1981 field isolates (e.g., positions 535 and 765), or by the

	1	70
BTW17	MEEFVIPVYSEDEIPIVALLSRYPLAIQTNKIEDVEGKHNVVKIPESDMIDIPKLTVIEMNYKPARNDG	
17B81R...E.....R.R..	
17C81R.....V..R.....	
17O81R.....V..R.....N..	
17B90ZR.....	
17O90XR.....	
17O90YR.....N.....	
	71	140
BTW17	IVVPRLLDITLRAYDDRKSFKSARGIEFMTNARWMKWAIDDRMDIQPLKVTLHDYC SVNHQLFNCVVKAN	
17B81	
17C81G.T.....	
17O81T.....	
17B90Z	
17O90X	
17O90Y	
	141	210
BTW17	AANADTIYYDYFPLEDYKKRCNHTNL DLLRSLTNMELFH ALQGAAYS IKSSYELVAYSERGSLEETYVVG	
17B81H.....N..E.....A..	
17C81H.R.....N.....AI..	
17O81H.R.....N.....AI..	
17B90ZH.....N..E.....A..	
17O90XH.....N..E.....A..	
17O90YH.....N..E.....A..	
	211	280
BTW17	QPKWIHLTRGTRIGNSGLSYERFISSMVQVS VNGKIPDEIANEIAQINRIRAEWITATYDRGRIRALELC	
17B81T.....I.....SV.....V.....	
17C81P.....I.....S.....V.....A.....	
17O81P.....I.....S.....V.....A.....	
17B90ZP.....I.....S.....V.....A.....	
17O90XP.....I.....S.....A.....	
17O90YP.....I.....S.....V.....A.....	
	281	350
BTW17	SILSTIGRKMLNTHEEPKDEM DLSTRFQFKLDEKFNRADSEHVNIFGVRGPATDEGRFYALIAATDTQ	
17B81	N.....I.....P.....	
17C81	K.....I.....E.....P.....	
17O81	K.....I.....P.....A.....	
17B90Z	K.....P.....A.....	
17O90X	K.....P.....A.....	
17O90Y	K.....P.....	
	351	420
BTW17	KGRVWRTNPYPCLRGALVAECELGDVYSTLRRVYTWSLRPEYGQHERQLENNKYVFNRINLFDSNLAVG	
17B81R.....	
17C81R.....R.....	
17O81R.....	
17B90ZR.....	
17O90X	M.....A.....	
17O90YR.....	

Fig. 2. Alignment of the deduced amino acid sequences of the VP2 proteins of the US prototype and field isolates of BTW 17. Amino acids identical to those of the prototype are denoted by a dot.

Examples described in the text are indicated by arrows.

prototype and all the 1990 viruses (e.g., positions 38 and 290). Furthermore amino acid substitutions occur which were common to only two of the 1981 isolates (e.g., position 159 in 17C81 and 17O81) or the 1990 isolates (e.g., position 527 in 17B90Z and 17O90X). The same amino acid change was shared by some viruses isolated in different years, while the other isolates had the same amino acid as the prototype (e.g., position 64). The data indicate that some amino acid changes in VP2 of field strains were conserved through time, whereas others were not correlated with the year of isolation and some substitutions were unique to individual viruses. The

	421		490
BTV17	DQIIHWRYEVKASAETTYDSGYMCRHEAEEDELLCKINEDKYKEMLDRMIQGGWDQERFKLHNILTDPNL		
17B81	D.
17C81N..N.....	G.....	D.....
17O81N..N.....	G.....	D.....
17B90Z
17O90X	D.....
17O90Y	D.....
	491	▼	560
BTV17	LTIDFEKDAYLNSRSSELVLPDYFDKWISSPMFNARLRTIKGEIGTSKKDDPWNNAVRGYIKPLAESLDF		
17B81L.....	S.....
17C81L.....	S.....
17O81L.....	SP.....
17B90ZL.....	K.....V.....	RSP.....
17O90XL.....	K.....V.....	SP.....
17O90YQ.....L.....V.....	SP.....
	561	▼	630
BTV17	VLGPYYDLRLLFDETSLKQEQAQFQYLSQLDDFPALTQLRGDAVCPhSGGALYTFRKVALFLIGNYE		
17B81G.....	
17C81G.A.....T.....	
17O81G.....T.....	
17B90ZG.A.....T.....	
17O90XG.A.....T.....	
17O90YG.A.....T.....	
	631	▼	700
BTV17	KLSPDLHEGMHQRYVHPSTGGTYQKRVLEMKDSCQLTCFVIDYIFEKREQLRDTKEARYIVYLIQSLTG		
17B81A.....C.....P.....M.....	
17C81T.....C.....P.....	
17O81T.....C.....P.....	
17B90ZT.....C.....P.....M.....	G.....V.....	
17O90XT.....C.....P.....M.....	G.....	
17O90YT.....C.....P.....M.....	Q.....G.....V.....	
	701	▼	770
BTV17	TQRQLSRLSTFPNFFQRLLMLKEIKFVRDLNVINFLPLMFLVHDNISYWHRQWSIPMVLFDDTIKLIPVE		
17B81D..K.....	
17C81D..K.....	A.....S.....	
17O81D..K.....S.....	
17B90ZD..K.....T..V.....S.....	R.....
17O90XD..K.....T..V.....S.....	R.....
17O90YD..K.....G..T..V.....S.....	R.....
	771	▼	840
BTV17	VGAYANRGFGKSFMNFTRFHPGELKKKQIAEDIHKFGVVAFEYYNTKISQGNVHTPVMTTKMDVLRVH		
17B81S.....S.....	KI.
17C81S.....M.....S.....	KI.
17O81S.....M.....S.....	KI.
17B90ZQ.....S.....M.....S.....	KI.
17O90XS.....M.....S.....	KI.
17O90YS.....M.....S.....	KI.

Fig. 2 (continued).

predicted VP2s constitute therefore a group of non-identical, but closely related proteins. The similar amino acid constitution of these proteins is reflected in the similarity of their predicted hydrophobic profiles and secondary structure as predicted by the methods of Kyte and Doolittle (1982) and Chou and Fasman (1974) respectively. Some cysteine (position 657), proline (positions 320 and 664) and glycine (position 574) residues were conserved amongst all field isolates, but absent in the prototype strain. Conversely, a proline was present in the prototype (position 553) and absent in all field isolates. These amino acids can significantly influence secondary structure of the proteins, so their presence or absence could alter the local or overall conformation of VP2 and so induce important phenotypic changes. Even minor conformational changes can have significant effects on virus neutralization or virulence, as has been demonstrated for other RNA viruses (Tuffereau et al., 1989; Aldovini and Young, 1990; Wakefield et al., 1992; De-Maula et al., 1993).

Phylogenetic analyses of field and prototype strains of BTV 17

Phylogenetic relationships between the L2 genes of the five US prototype strains of BTV and the six BTV 17 field isolates were determined using a distance matrix method (Fig. 3) and bootstrap analysis (Fig. 4). Two distinct monophyletic groups are evident, one which includes BTV 2 and BTV 13, and the other which includes the prototype strains of BTV 10, BTV 11 and BTV 17 as well as all six field isolates of BTV 17. This topology is supported by the high confidence values (100%) obtained by bootstrap analyses (Fig. 4).

The monophyletic grouping of BTV 10, 11 and 17 is supported by the common deletions present in the nucleotide sequences of their L2 genes. These deletions are evident by alignment with the L2 gene of EHDV 1. The deletions occur in groups each of three nucleotides between nucleotides 301–302, 384–385, 499–500, 1327–1328, 1374–1375 and 1478–1479 of the prototype strains as compared to EHDV 1. The prototype and field isolates of BTV 17 also share a common deletion between position 1827–1828 which is not present in the other four US serotypes of BTV. Additional deletions are present between positions 1919–1920

	841		910
BTW17	LSSLCAGLADSVVYTL	PVAHPKKCIVLII	VGDDKLEPHTRSEQIVSRYNYSRKHICGIVSVTIGQNSQLR
17B81I.....	T.....V.....
17C81I.....V.....
17O81I.....V.....
17B90ZI.....V.....
17O90XI.....	I.V.....
17O90YI.....V..EE..V.....

	911		955
BTW17	VHTSGIVKHRCDFILKHCKVILVRMPGYVFGNDELMTKLLNV		
17B81	
17C81	W.....	
17O81	
17B90Z	
17O90X	
17O90Y	

Fig. 2 (continued).




Fig. 3. Phylogenetic tree of the six BTV 17 isolates studied and the five US BTV prototype strains. EHDV 1 was used as an outgroup.

of the prototype strains of BTV 10 and BTV 11, and positions 1916–1917 of the prototype and field isolates of BTV 17. The common deletions in the L2 genes of BTV 10, 11 and 17 suggest that these deletions occurred in a common ancestor prior to the divergence of each serotype. BTV 10 and BTV 11 do cluster as a single group with a common origin, however this topology is not statistically significant (61%) by bootstrap analysis. This lack of defined relationship within the second




Fig. 4. Tree diagram showing the majority rule consensus tree resulting from 100 bootstrap replicates of the six L2 genes of the BTV 17 field isolates and the five US prototype strains. The number at each node indicates the number of times that the group above and to the right is found.

TABLE 3
Genetic distances of the BTV 17 field isolates and US BTV and EHDV1 prototype strains

	17B81 ^a	17C81 ^a	17O81 ^a	17B90Z ^a	17B90X ^a	17O90Y ^a	BTV 17 ^b	BTV 10 ^b	BTV 11 ^b	BTV 2 ^b	BTV 13 ^b	EHDV 1 ^c
17B81	0.0000	0.0250	0.0250	0.0322	0.0329	0.0315	0.0523	0.4381	0.4315	1.0496	1.0343	1.6177
17C81	0.0000	0.0091	0.0311	0.0318	0.0318	0.0300	0.0575	0.4396	0.4412	1.0547	1.0454	1.6188
17O81	0.0000	0.0261	0.0261	0.0275	0.0279	0.0594	0.4411	0.4385	1.0558	1.0395	1.6138	
17B90Z	0.0000	0.0098	0.0098	0.0091	0.0091	0.0645	0.4357	0.4385	1.0469	1.0418	1.5947	
17O90X	0.0000	0.0119	0.0119	0.0657	0.4299	0.4329	1.0444	1.0440	1.5984			
17O90Y		0.0000	0.0645	0.0645	0.4366	0.4368	1.0573	1.0519	1.6130			
BTV 17			0.0000	0.0000	0.4263	0.4335	1.0375	1.0492	1.6525			
BTV 10				0.0000	0.4547	1.0676	1.0817	1.7577				
BTV 11					0.0000	1.0806	1.0484	1.7952				
BTV 2						0.0000	1.8653	2.1702				
BTV 13							0.0000	2.1557				
EHDV 1								0.0000	0.0000			

^a BTV 17 field isolates.^b US BTV prototype strains.^c EHDV 1 prototype strain.

monophyletic group is further evidenced by comparison of the genetic distances of the BTV 17 field isolates to the prototype strains of BTV 10 and 11. The L2 genes of the prototype BTV 17 and field isolates 17C81 and 17B90Z are slightly more closely related to BTV 10 than they are to BTV 11, as determined by genetic distance matrices (Table 3). In contrast, the genetic distance values of 17O90X, 17B81 and 17O81 are slightly closer to BTV 11 than they are to BTV 10, whereas the field isolate 17O90Y is almost equidistant between the L2 genes of BTV 10 and BTV 11 (Table 3). The evolutionary relationships amongst viruses within the monophyletic group of BTV 10, 11 and 17, therefore, are undefined.

The phylogenetic analyses indicate that the prototype and field isolates of BTV 17 have descended from a common ancestor, as would be expected. Bootstrap analyses placed a confidence value of 100% on this monophyletic group. Isolate 17B81 is placed as a sister in an independent branch of the tree (confidence value of 100%). Although the tree topology indicates that all the other viruses (17O81, 17C81, 17O90X, 17B90Z and 17O90Y) could be descendants from a common immediate ancestor, the low bootstrap value of 93% does not support this concept. This suggests that these viruses could derive through different evolutionary pathways within the lineage. The two field viruses 17O81 and 17C81, although isolated from the same farm and sharing the same ancestor (100% confidence value), have diverged from one another perhaps to successfully replicate in the different host species from which they were isolated. The topology of the tree indicates that the three 1990 field isolates also were derived from a common origin (100% confidence value), however the branching pattern which has two viruses (17O90X and 17B90Z) emerging from the same node is not statistically significant (54% confidence value), indicating that these three viruses too might have arisen via different evolutionary pathways. The phylogenetic analysis of the L2 genes of prototype and field strains of BTV 17 suggests that the unique viral variants of a single BTV serotype, which co-circulate in the same geographical region, could evolve from different evolutionary pathways.

Discussion

The objective of this study was to further characterize the evolution and genetic variation of BTV 17 in a localized region of California by sequencing the L2 gene of field strains of BTV 17. The L2 gene was selected as it encodes outer capsid protein VP2 which expresses the neutralizing determinants of BTV, thus the L2 gene is the only serotype specific genome segment of BTV. The viruses compared in the study were selected from a panel of field isolates with different neutralization profiles (MacLachlan et al., 1992). In spite of their restricted origin, both geographically and spatially, the L2 genes of these viruses were heterogenous which is consistent with the phenotypic variability of these viruses.

The data indicate that the five US BTV serotypes segregate into two monophyletic groups with BTV 2 and 13 forming one group and BTV 10, 11 and 17 the other. This is in agreement with previous phylogenetic analyses based on the

sequence of the L2 and VP2 of prototype strains of BTV (Gould and Pritchard, 1990; Roy, 1992). Whereas it previously was concluded using L2 gene sequences that prototype strains of BTV 10 and BTV 17 are more closely related to one another than they are to BTV 11 (Gould and Pritchard, 1990; Kowalik and Li, 1991), our analyses with field isolates of BTV 17 demonstrate that the relationship between the viruses of this group is less certain. Sequencing of the L2 gene of field strains of BTV 10 and BTV 11 likely will be necessary to more accurately define the evolutionary relationships between BTV 10, 11 and 17. Earlier phylogenetic studies of the BTV serogroup used the nucleotide sequences from conserved BTV genes (Gould and Pritchard, 1991; Hwang et al., 1992a), and these analyses produced trees which suggest different evolutionary relationships between the BTV serotypes than those we obtained. These divergent results likely can be attributed to the evolutionarily conserved nature of the genes characterized (Hwang et al., 1992b), and to the likelihood of reassortment of viral genes amongst viruses which co-circulate. Reassortment of gene segments clearly occurs amongst strains of the four serotypes of BTV present in the western US (Heidner et al., 1991), thus the serotype-specific L2 gene must be used to characterize the evolutionary relationships amongst viruses isolated from a single region.

Analyses of genomes from field strains of other RNA viruses indicate that multiple variant genomes may occur which differ from the average consensus sequence that represents the prototype virus sequence. Different prototype consensus sequences exist for viruses corresponding to particular geographical regions (Domingo and Holland, 1988). Our data indicate that the L2 gene sequences of BTV 17 isolates made in California differed from each other at the nucleotide level by an average of 2.3%, whereas they differed from the prototype BTV 17 virus isolated in Wyoming in 1962 by an average of 5.8%. Our results also identified mutations common to all the California field isolates with respect to the prototype virus. These genetic differences might reflect the comparison of two different prototype consensus sequences of viruses which have evolved in different geographical areas. Hence, it would be interesting to evaluate the concept of geographical prototype by comparing the US BTV 17 prototype virus with field isolates obtained in Wyoming as well as those obtained in California.

Results of this study clearly demonstrate genetic heterogeneity amongst field isolates of the same BTV serotype which were obtained from the same geographical region at an interval of nine years. Heterogeneity was even demonstrated in the L2 genes of two isolates obtained from the same farm in the same year (17O81 and 17C81). The accumulation of mutations in the field isolates is not proportional to the interval between their years of isolation (Table 2), and some nucleotide changes present in the 1981 isolates were absent in those from 1990. The data are not, therefore, consistent in all instances with a direct ancestor-descendant relationship between the field isolates, rather individual viruses might evolve via different evolutionary pathways. This conclusion is further supported by the phylogenetic analyses which indicate that although all field isolates are included in the same monophyletic group, they do not share an immediate common ancestor.

The natural replication cycle of BTV, which includes an insect vector and

various ruminant hosts, offers significant opportunity for the generation of genetically novel viruses. First, reassortment of individual viral genome segments can occur in both the vector and the ruminant host (Roy, 1989; Heidner et al., 1991). Second, it is likely that only viruses best adapted to replication in either the vector or the ruminant host will be amplified. The lack of proof-reading capability of the BTV RNA polymerase also might contribute to genetic diversity during replication in the vector or ruminant host (Kowalik and Li, 1991). Finally, immune selection of variant viruses likely occurs in infected ruminants. All these phenomena might account for the different evolutionary pathways apparently followed by various field strains of BTV 17, and to the generation of the related but not identical viruses which co-circulate in the San Joaquin Valley.

Acknowledgements

We gratefully acknowledge Dr. John Gillespie, Department of Genetics, University of California, Davis, for his advice and helpful discussions. This work was supported by USDA Competitive Grant 91-37204-6407, and funds provided by the Livestock Diseases Research Laboratory and USDA under the Animal Health Act, 1977, Public Law 95-113.

References

- Akita, G.Y., Chinsangaram, J., Osburn, B.I., Ianconescu, M. and Kaufman, R. (1992) Detection of bluetongue virus serogroup by polymerase chain reaction. *J. Vet. Diagn. Invest.* 4, 400-405.
- Aldovini, A. and Young, R.A. (1990) Mutations of RNA and protein sequences involved in human immunodeficiency virus type 1 packaging result in production of noninfectious virus. *J. Virol.* 64, 1920-1926.
- Appleton, J.A. and Letchworth, G.L. (1983) Monoclonal antibody analysis of serotype restricted and unrestricted bluetongue viral antigenic determinants. *Virology* 124, 286-299.
- Chou, P.Y. and Fasman, G.D. (1974) Prediction of protein conformation. *Biochemistry* 13, 222-245.
- Cowley, J.A. and Gorman, B.M. (1989) Cross-neutralization of genetic reassortants of bluetongue virus serotypes 20 and 21. *Vet. Microbiol.* 19, 37-51.
- de Mattos, C.A., de Mattos C.C., Dangler, C.A., Osburn, B.I., Ianconescu, M. and Kaufman, R. (1992) Hybridization relatedness of Israeli and US bluetongue virus (BLU) serotypes using cDNA probes from BLU virus strain 11-UC8. *Arch. Virol.* 122, 23-33.
- de Mattos, C.C., de Mattos, C.A., Osburn, B.I., Dangler, C.A., Doi, R. and Chuang, R.Y. (1989) Recombinant DNA probe for serotype-specific identification of BTV 17. *Am. J. Vet. Res.* 50, 536-541.
- de Mattos, C.C., de Mattos, C.A., Osburn, B.I., Ianconescu, M. and Kaufman, R. (1991) Evidence of genome segment 5 reassortment in bluetongue virus field isolates. *Am. J. Vet. Res.* 52, 1794-1798.
- DeMaula, C.D., Heidner, H.W., Rossitto, P.V., Pierce, C.M. and MacLachlan, N.J. (1993) Neutralization determinants of United States bluetongue virus serotype ten. *Virology* 195, 292-296.
- Domingo, E. (1989) RNA virus evolution and the control of viral disease. *Prog. Drug Res.* 33, 93-133.
- Domingo, E., Davila, M. and Ortín, J. (1980) Nucleotide sequence heterogeneity of RNA from natural population of foot and mouth disease virus. *Gene* 11, 333-346.
- Domingo, E. and Holland, J.J. (1988) High error rates, population equilibrium and evolution of RNA replication systems. In: E. Domingo, J.J. Holland and P. Ahlquist (Eds.), *RNA Genetics*, Vol. III. Variability of RNA Genomes, pp. 3-36. CRC Press, Boca Raton, FL.

- Faragher, S., Meek, A., Rice, C. and Dalgarno, L. (1988) Genome sequences of a mouse-avirulent and mouse-virulent strain of Ross River virus. *Virology* 163, 509–526.
- Felsenstein, J. (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783–791.
- Felsenstein, J. (1991) PHYLIP (phylogeny inference package) 3.4 Manual. University of Washington, Seattle, Department of Genetics SK-50, WA 98195.
- Fitch, W.M. and Margoliash, E. (1967) Construction of phylogenetic trees. *Science* 155, 279–284.
- Genetics Computer Group (1991) Program Manual for the GCG Package, Version 7, 575 Science Drive, Madison, WI 53711.
- Ghiashi, H., Fukuso, A., Eshita, Y. and Roy, P. (1987) Identification of conserved and variable regions in the neutralization VP2 gene of bluetongue virus. *Virology* 160, 100–109.
- Gould, A.R. and Pritchard, L.I. (1990) Relationships amongst viruses revealed by comparisons of capsid and outer capsid coat protein nucleotide sequences. *Virus Res.* 17, 31–52.
- Gould, A.R. and Pritchard, L.I. (1991) Phylogenetic analyses of complete nucleotide sequence of the capsid protein (VP3) of Australian epizootic hemorrhagic disease of deer virus (serotype 2) and cognate genes from other orbiviruses. *Virus Res.* 21, 1–18.
- Greider, F.B. and Schultz, K.T. (1990) Antigenic topography of bluetongue virus 17. *Microbiol. Path.* 8, 299–303.
- Heidner, H.W., Iezzi, L.G., Osburn, B.I. and MacLachlan, N.J. (1991) Genetic variation and evolutionary relationships amongst bluetongue viruses endemic in the United States. *Virus Res.* 21, 91–109.
- Hwang, G.Y., Chiou, J.K., Yang, Y.Y. and Li, J.K. (1992a) Comparative sequence analyses of the cognate structural protein VP6 genes of five US bluetongue viruses. *Virus Res.* 24, 315–323.
- Hwang, G.Y., Yang, Y.Y., Chiou, J.F. and Li, J.K. (1992b) Sequence conservation among the cognate non-structural NS3/NS3A protein genes of six bluetongue viruses. *Virus Res.* 23, 151–161.
- Iwata, H., Yamagawa, M. and Roy, P. (1992) Evolutionary relationships among the gnat transmitted orbiviruses that cause African horse sickness, bluetongue, and epizootic hemorrhagic disease as evidence by their capsid proteins sequences. *Virology* 191, 251–261.
- Kimura, M. (1980) A simple model for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* 16, 111–120.
- Kinnunen, L., Huuvalainen, A., Poyry, T. and Ho, T. (1990) Rapid molecular evolution of wild type 3 poliovirus during infection in individual hosts. *J. Gen. Virol.* 71, 317–324.
- Kowalik, T.F. and Li, J.K. (1991) Bluetongue virus evolution: sequence analyses of the genomic S1 segments and major core protein VP7. *Virology* 181, 749–755.
- Kyte, J. and Doolittle, R.F. (1982) A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* 157, 105–132.
- MacLachlan, N.J., Rossitto, P.V., Heidner, H.W., Iezzi, L.G., Yilmaz, T.D., DeMaula, C.D. and Osburn, B.I. (1992) Variation amongst the neutralizing epitopes of bluetongue viruses isolated in the United States in 1979–1981. *Vet. Microbiol.* 31, 303–316.
- Mertens, P.P.C., Pedley, S., Cowley, J., Burroughs, J.N., Corteyn, A.H., Jeggo, M.H., Jennings, D.M. and Gorman, B.M. (1989) Analysis of the roles of bluetongue virus outer capsid proteins VP2 and VP5 in determination of virus serotype. *Virology* 170, 561–565.
- Nichol, S.T. (1988) Genetic diversity of enzootic isolates of vesicular stomatitis virus New Jersey. *J. Virol.* 62, 572–579.
- Rico-Hesse, R., Roehrig, J.T., Trent, D.W. and Dickerman, R.W. (1988) Genetic variation of Venezuelan equine encephalitis virus strains of the ID variety in Colombia. *Am. J. Trop. Hyg.* 38, 195–204.
- Rossitto, P.V. and MacLachlan, N.J. (1992) Neutralizing epitopes of the serotypes of bluetongue virus present in the US. *J. Gen. Virol.* 73, 1947–1952.
- Roy, P. (1989) Bluetongue virus genetics and genome structure. *Virus Res.* 13, 179–206.
- Roy, P. (1992) Bluetongue virus proteins. *J. Gen. Virol.* 73, 3051–3064.
- Shope, R.E., MacNamara, L.G. and Mangold, R. (1960) A virus-induced epizootic hemorrhagic disease of the Virginia white-tailed deer (*Odocoileus virginianus*). *J. Exp. Med.* 3, 155–170.
- Sobrino, F., Palma, E.L., Bech, E., Davila, M., de la Torre, J.C., Negro, P., Villanueva, N., Ortin, J. and Domingo, E. (1986) Fixation of mutations in the viral genome during an outbreak of foot and mouth disease: heterogeneity and rate variations. *Gene* 50, 149–159.

- Tuffereau, C., Leblois, H., Benejean, J., Coulon, P., Lafay, F. and Flamand, A. (1989) Arginine or lysine in position 333 of ERA and CVS glycoprotein is necessary for rabies virulence in adult mice. *Virology* 172, 206–212.
- Wakefield, J.K., Jablonski, S.A. and Morrow, C.D. (1992) In vitro enzymatic activity of human immunodeficiency virus type 1 reverse transcriptase mutants in the highly conserved YMDD amino acid motif correlates with the infectious potential of the proviral genome. *J. Virol.* 66, 6806–6812.
- Weaver, S.C., Bellew, L.A. and Ricco-Hesse, R. (1992) Phylogenetic analysis of alphaviruses in the Venezuelan equine encephalitis complex and identification of the source of epizootic viruses. *Virology* 191, 282–290.
- Webster, B.G., Bean, W.J., Gorman, O.T., Chambers, T.M. and Kawaoka, Y. (1992) Evolution and ecology of influenza A viruses. *Microbiol. Rev.* 56, 152–179.
- Wege, H., Stephenson, J.R., Wege, H. and Meulen, V. (1981) Genetic variation of neurotropic and non-neurotropic murine coronaviruses. *J. Gen. Virol.* 54, 67–74.
- Yamashita, M., Krystal, M., Fitch, W. and Palese, P. (1988) Influenza B virus evolution: Co-circulating lineages and comparison of evolutionary pattern with those of influenza A and C viruses. *Virology* 163, 112–122.
- Yang, Y.Y., Chiou, J.F., Hwang, I.J. and Li, J.K. (1992) Evolutionary analyses of five US bluetongue viruses using the cognate S2 genes. *Virus Res.* 25, 241–249.