




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Hydroxychloroquine as an aerosol might markedly reduce and even prevent severe clinical symptoms after SARS-CoV-2 infection



A. Klimke^{a,b,*}, G. Hefner^c, B. Will^d, U. Voss^{a,e}

^a Vitos Klinikum Hochtaunus, Friedrichsdorf, Germany

^b Heinrich Heine University Duesseldorf, Germany

^c Klinik für Forensische Psychiatrie, Vitos Rheingau Eltville, Germany

^d Rheinische Friedrich-Wilhelms-Universität Bonn, University Hospital (UKB), Germany

^e Goethe-Universität Frankfurt am Main, Abt. Allgemeine Psychologie II, Germany

ARTICLE INFO

Keywords:

COVID-19

Hydroxychloroquine aerosol

Pharmacotherapy

Prevention

SARS-CoV-2

ABSTRACT

Covid-19 is a new coronavirus disease first described in December 2019. This respiratory illness is severe and potentially fatal. Severe cases make up to 15%, lethality ranges between 1.5 and more than 10%. What is urgently needed is an efficient pharmacological treatment for the treatment of severe cases. During the infection of alveolar epithelial cells of the lung, the ACE2 receptor has a central function. The antimalarial drugs chloroquine phosphate (CQ) and hydroxychloroquine (HCQ) impair in vitro the terminal glycosylation of ACE2 without significant change of cell-surface ACE2 and, therefore, might be potent inhibitors of SARS-CoV-2 infections. Starting inhibition at 0.1 μM , CQ completely prevented in vitro infections at 10 μM , suggesting a prophylactic effect and preventing the virus spread 5 h after infection. In a first clinical trial, CQ was effective in inhibiting exacerbation of pneumonia, improving lung imaging findings, promotion of virus-negative conversion, and shortening the disease. In addition, HCQ, which is three times more potent than CQ in SARS-CoV-2 infected cells (EC50 0.72 μM), was significantly associated with viral load reduction/disappearance in COVID-19 patients compared to controls. Theoretically, CQ and HCQ could thus be effectively used in the treatment of SARS-CoV pneumonia. From a pharmacological standpoint, however, the major problems of oral treatment with these drugs are possible severe side effects and toxicity. Concretely, this relates to (a) the inconsistent individual bioavailability of these drugs at the alveolar target cells, depending on intestinal resorption, hepatic first-pass metabolism and accumulation in liver, spleen and lung, and (b) the need for a relatively high concentration of 1–5 μM at the alveolar surface.

Therefore, we propose in a first dose estimation the use of HCQ as an aerosol in a dosage of 2–4 mg per inhalation in order to reach sufficient therapeutic levels at the alveolar epithelial cells. By using a low-dose non-systemic aerosol, adverse drug reactions will markedly be reduced compared with oral application. This increase in tolerability enables a broader use for prevention and after contact with an infected person, which would be an advantage especially for the high-risk, often multi-morbid and elderly patients.

Empirical data on self-medication with a one-week aerosol application by two of the authors is presented. Inhalation was well tolerated without relevant side effects.

Introduction/background

The 2019-nCoV pandemic is rapidly spreading with currently more than a million reported cases and 57.000 deaths worldwide [1]. While the majority of infections with Covid-19 are asymptomatic or show a mild course [2,3], up to 15% of patients develop a severe and potentially fatal respiratory illness with an estimated lethality ranging between 1.5 and more than 10% [4]. Currently, as no effective antiviral

pharmacological treatment is available, lethality seems to depend on individual risk factors including age, gender and hypertension and regarding the course of the disease in cases of severe pneumonia, on the availability of large-scale intensive care units and extracorporeal membrane oxygenation (ECMO) [5,6].

Regarding pharmacological treatment, several clinical trials have recently been initiated, e.g. with lopinavir/ritonavir [7], remdesivir peptide (EK1) [8] and interferon alpha [9]. The latter originally being

* Corresponding author. Vitos Klinikum Hochtaunus, Klinik für Psychiatrie und Psychotherapie, Emil-Sioli-Weg 1-3, D- 61381 Friedrichsdorf, Germany.

E-mail address: ansgar.klimke@vitos-hochtaunus.de (A. Klimke).

used for the treatment of hepatitis B (novaferon, Beijing Genova Biotech). However, the clinical efficacy of these new drugs for COVID-19 has not been proven and their safety has not been extensively tested.

Further, CQ and HCQ have been discussed as promising, cost-effective and easily available agents in the treatment of COVID-19 [10–12]. These are relatively old agents, introduced primarily for prophylaxis and treatment of malaria and also for rheumatic diseases [13]. In vitro cell cultures, HCQ seems to be a more potent inhibitor of infection with SARS-CoV-2 than chloroquine [14,15]. Both drugs, taken orally, may cause severe side effects [16] ranging from psychiatric symptoms [17] to ocular toxicity [18] to myocardial dysfunction [19], even at only low overdosage [20]. These side effects may be severe and, therefore, limit widespread application in vivo, at least at the present time. As Drosten [21] pointed out, a major limiting factor of oral hydrochloroquine treatment lies in the fact that the drug may not reach the target, i.e. the surface of the lungs. Instead, it is metabolized internally, requiring treatment dosages which, in turn, may cause toxic and sometimes even lethal side effects.

Here, we propose a different approach to the investigation of the effectiveness of the hydrochloroquine drug that would imply a direct application of the active agent to the respiratory tract.

The hypothesis/theory

We hypothesize that HCQ especially *as an aerosol application* will prevent or at least markedly reduce the replication rate of the SARS-CoV-2 virus *in the early phase of the infection* and subsequently substantially lower the number of severe pneumonias and casualties.

Why this hypothesis is different from current thinking

This hypothesis is new since the major assumption in ongoing clinical studies and actual recommendations is that HCQ and CQ should be used in oral application form in patients with severe covid-19 pneumonia and only when other treatment strategies have failed. However, the typical clinical course of this infection suggests that the virus load in the respiratory tract increases stepwise starting with mild symptoms and ending in up to 15% of patients with severe and potentially life-threatening pneumonia [4]. Therefore, the treatment with a drug which inactivates the cell receptor for the virus should start after exposition with high risk, e.g., when one person was infected very recently with the virus or is in the early phase of the disease. Moreover, our hypothesis differs from the standard recommendation to try HCQ/CQ in a late phase of the disease when other antiviral drugs failed. We believe that a respiratory virus infection should be treated very early because the severe acute respiratory syndrome is caused by ion channel activity of the viroporin 3a which activates the NLRP3 inflammasome [22]. Unfortunately, as of now, there is no evidence yet that HCQ/CQ has any inhibiting effect on this inflammasome activation.

How has this idea evolved?

The idea to propose application of HCQ/CQ as aerosol is generated because one major objection against the clinical efficacy of these drugs is that they have to be administered in relatively high oral dosages. Such high dosages may have several toxic side effects, strongly limiting their utilizability as preventive treatment. An aerosol application of drugs which are primarily intended to act on the respiratory system is well established for several drugs, e.g. in the treatment of asthma with corticosteroids, e.g. budesonide [23,24] and beta mimetica, e.g. fenoterol [25], and in the early treatment of influenza (during the first 48 h) with neuraminidase blockers like zanamivir [26]. Moreover, there are reports of undergoing clinical studies of aerosol interferon alpha (novaferon) for treatment of COVID-19 [27] leading us to advocate a clinical trial to evaluate also HCQ/CQ in this application form.

Evaluation of the hypothesis

Why hydroxychloroquine might be efficacious in COVID-19

It has been demonstrated that the SARS-CoV-2 virus enters ACE2-expressing cells including alveolar epithelial cells of the lung and in other organs [28–30], which has been shown before also for SARS-CoV-1. Therefore, during the infection of alveolar epithelial cells of the lung, the ACE2 receptor has a central function [31].

The antimalarial drugs CQ and HCQ impair the terminal glycosylation of ACE2 without significant change of cell-surface. ACE2 increases the local pH value, which reduces the activity of cathepsin L needed for hydrolysis of the viral S protein and might influence the generation of pro-inflammatory cytokines [32]. Therefore CQ might be a potent inhibitor of SARS-CoV infection [29,33,34].

In vitro, CQ, starting with 0.1 μM inhibited and, moreover, completely prevented SARS-CoV infections in cell cultures at 10 μM , suggesting a prophylactic effect and preventing the virus spread 5 h after infection. HCQ is threefold more potent than CQ in SARS-CoV-2 infected cells, resulting in the clinical recommendation to treat orally with 800 mg at the first day and at 400 mg on the following four days [27].

Two major points of action are postulated for CQ/HCQ in the treatment of COVID-19: (1) they perturb the terminal glycosylation of the ACE2 protein and inhibit cell-binding of the virus, (2) the alkalinizing medication elevates the pH value of the endosomes. Since Cathepsin L requires an acidic environment to crack the viral S protein, virus induced activities are reduced. Other possible explanations for the antiviral effect include a reduction in MAK kinase activation or a disruption of the maturation of the viral M protein. Further, CD8 + cells directed against the virus may be activated and thereby lead to a decrease of pro-inflammatory cytokines [32].

Chinese scientists from Peking University found low inhibitory concentrations of 0.72 μM for the hydroxy derivative in direct comparison to chloroquine (5.47 μM) in cell culture experiments with the pathogen SARS-CoV-2. Both figures are EC50 values, that is, concentrations that cause half-maximal inhibition of virus replication. They calculated that with an initial dose of 400 mg twice daily and a subsequent dose of 200 mg hydroxychloroquine sulfate twice daily there should be significantly higher concentrations in the lung tissue than needed for in vitro inhibition. However, the concentrations have not been measured and clinical studies that examine the possible benefit of such treatment are not yet available [15].

Another group of Chinese virologists published slightly different EC50 values for the in vitro inhibition of viral replication by CQ and HCQ (3.8 μM and 4.1 μM at 0.02 MOI = multiplicities of infection). As the results depend on the type of cell culture, the cells used, the amount of viruses used for infection, the time of evaluation and other factors, this is not surprising [8,14].

Moreover, a first clinical study with more than 100 patients demonstrated superiority of chloroquine phosphate to controls in inhibiting exacerbation of pneumonia, improving lung imaging findings, promotion of virus-negative conversion, and shortening the disease without severe adverse reactions [11]. In a non-randomized open-label trial in 20 patients, hydroxychloroquine treatment was significantly associated with viral load reduction/disappearance in COVID-19 patients compared to controls [12].

Why the oral application form might have disadvantages

From a pharmacological standpoint, one major problem of oral treatment with chloroquine phosphate or hydroxychloroquine is that relatively high concentrations between 1 and 10 μM are needed to inhibit the glycosylation of the soluble ACE2 receptor at the alveolar surface in the lung. Therefore, the bioavailability of these drugs in the pulmonary target regions depends individually on intestinal resorption,

hepatic first-pass metabolism and diffusion from the blood to the alveolar cells. This could in part be compensated by higher oral dosages, which are, however, limited by adverse and potentially toxic side effects.

Another problem can be that CQ and HCQ are accumulated in the liver, spleen, kidney, lungs and other organs. 100 to 500 times of the chloroquine plasma concentration is found in parenchyma cells and 1000 times in pigmented cells. Around 55% of it is bound to plasma proteins (pharmaceutical product information).

From a clinical standpoint, potential fatal arrhythmias and drug-induced sudden cardiac death in rare cases or in case of overdose can be a resulting problem.

How can the efficacy of HCQ/CQ as aerosol be clinically evaluated in terms of prevention and dampening the severity of the course of illness?

Our hypothesis prompts to be tested in controlled randomized clinical studies. Currently, an aerosol preparation of HCQ is not commercially available. Therefore, experimental data is needed to examine whether it is possible to achieve and maintain an antiviral concentration of 1–5 μM in the alveolar cells which has been proven effective in vitro. In a first approximation, one can assume that in the normal adult, the lungs weigh approximately 1.000 g which is 0.014 of the whole body weight of 70 kg. Although it is difficult to predict in the lung the effects of blood perfusion and on the other hand accumulation of HCQ, it seems reasonable to substitute 2–400 mg HCQ b.i.d. in oral form (as recommended) and a bioavailability of 0.79 [35] with 2.2–4.4 mg HCQ b.i.d. per inhalation.

Since HCQ/CQ is an old drug with a known profile of side effects, also for much higher dosages, it might also be appropriate to perform open pilot studies with higher numbers of cases and compare (in a non-randomized study) the clinical course and final result in patients with similar symptoms but without HCQ inhalation.

For objective testing of the preventive potential, persons at risks like therapeutic staff following direct unprotected contact with corona patients as well as relatives of patients who have tested positive for SARS-CoV-2 and who are living in the same household should inhale HCQ in comparison to placebo during the incubation period of 5–7 days without symptoms. We would predict that the number of subsequent infections of the contact persons will be significantly lower in the verum group.

To test whether HCQ as aerosol lowers the severity of the course, a randomized comparison should be performed including patients with beginning mild symptoms who were actually tested SARS-CoV-2 positive in order to evaluate the effect on clinical symptoms including fever, cough, beginning pneumonia and other complications.

Finally, HCQ as aerosol could be tested preferentially as co-medication in more severely ill patients where an indication for oral HCQ treatment is given. The prediction would be that those with the co-medication will have a better treatment outcome than those in oral monotherapy with HCQ.

Empirical (own) data

Since there is currently no commercial aerosol application available, two of the authors (A.K., clinical director and B.W., medical student working at an ICU with acute COVID-19 patients) decided to test tolerability and possible side effects of the inhalation of HCQ, starting with a dosage of 1 mg b.i.d. (dissolved in 2 ml of 0.9% NaCl solution and using a commercial nebulizer generating a droplet size $< 5 \mu\text{m}$ which can reach the alveolar space) which was stepwise increased up to 4 mg per day over a period of one week. Inhalation was well tolerated without relevant side effects. The only observation was after 4 days the feeling of a transient bitter taste in the mouth which lasted 2–3 h after the inhalations.

Consequences of the hypothesis and discussion

Why this treatment strategy is of potential great importance

If our hypothesis is true, HCQ as an aerosol might not only reduce the side effect potential of the oral application form but can also be clinically used as an efficient antiviral agent in the early phase of COVID-19 and eventually lower the rate of severely ill patients and fatalities. This might have great relevance for further prognosis and treatment of this often fatal disease.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.mehy.2020.109783>.

References

- [1] <https://coronavirus.jhu.edu/map.html>, downloaded April 3, 2020.
- [2] Novel, C. P. E. R. E. (2020). The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi*, 41(2), 145.
- [3] Wu, Z., & McGoogan, J. M. (2020). Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *Jama*.
- [4] Baud D, Qi X, Nielsen-Saines K, Musso D, Pomar L, Favre G. Real estimates of mortality following COVID-19 infection. *Lancet Infect Dis* 2020.
- [5] Hamer, O. W., Salzberger, B., Gebauer, J., Stroszczyński, C., & Pfeifer, M. (2020, March). CT morphology of COVID-19: Case report and review of literature. In *RöFo-Fortschritte auf dem Gebiet der Röntgenstrahlen und der bildgebenden Verfahren*. © Georg Thieme Verlag KG.
- [6] Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet* 2020.
- [7] Cao B, Wang Y, Wen D, et al. A trial of lopinavir–ritonavir in adults hospitalized with severe Covid-19. *N Engl J Med* 2020.
- [8] Lai CC, Shih TP, Ko WC, Tang HJ, Hsueh PR. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and corona virus disease-2019 (COVID-19): the epidemic and the challenges. *Int J Antimicrob Agents* 2020;105924.
- [9] Dong L, Hu S, Gao J. Discovering drugs to treat coronavirus disease 2019 (COVID-19). *Drug Discover Therapeut* 2020;14(1):58–60.
- [10] Colson P, Rolain JM, Lagier JC, Brouqui P, Raoult D. Chloroquine and hydroxychloroquine as available weapons to fight COVID-19. *Int J Antimicrob Agents* 2020;105932.
- [11] Gao, J., Tian, Z., & Yang, X. (2020). Breakthrough: Chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Bioscience trends*.
- [12] Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents* 2020;105949.
- [13] Lu H. Drug treatment options for the 2019-new coronavirus (2019-nCoV). *Biosci Trends* 2020;14(1):69–71.
- [14] Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* 2020;30(3):269–71.
- [15] Yao X, Ye F, Zhang M, et al. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020.
- [16] Goel P, Gerriets V. Chloroquine. In *StatPearls [Internet]*. StatPearls Publishing; 2019.
- [17] Good MI, Shader RI. Lethality and behavioral side effects of chloroquine. *J Clin Psychopharmacol* 1982;2(1):40–7.
- [18] Schwartzman S, Samson CM. Are the Current Recommendations for Chloroquine and Hydroxychloroquine Screening Appropriate? *Rheumatic Dis Clinics* 2019;45(3):359–67.
- [19] Blignaut M, Espach Y, van Vuuren M, Dhanabalan K, Huisamen B. Revisiting the Cardiotoxic Effect of Chloroquine. *Cardiovasc Drugs Ther* 2019;33(1):1–11.
- [20] Frisk-Holmberg M, Bergkvist Y, Domeij-Nyberg B, Hellström L, Jansson F. Chloroquine serum concentration and side effects: Evidence for dose-dependent kinetics. *Clin Pharmacol Ther* 1979;25(3):345–50.
- [21] <https://www.pharmazeutische-zeitung.de/virologe-drosten-nimmt-stellung-zu-chloroquin/>; downloaded 04.02.2020.
- [22] Chen, I. Y., Moriyama, M., Chang, M. F., & Ichinohe, T. (2019). Severe acute

- respiratory syndrome coronavirus viroporin 3a activates the NLRP3 inflammasome. *Front Microbiol*, 10.
- [23] Chen Y. Adjunctive Corticosteroid Therapy for Patients with Severe Novel Coronavirus Pneumonia (COVID-19): a Randomized Controlled Trial. <http://www.chictr.org.cn/showproj.aspx?proj=48777> [Accessed 03 April 2020].
- [24] Pauwels RA, Löfdahl CG, Postma DS, Tattersfield AE, O'Byrne P, Barnes PJ, et al. Effect of inhaled formoterol and budesonide on exacerbations of asthma. *N Engl J Med* 1997;337(20):1405–11.
- [25] Salome CM, Schoeffel RE, Woolcock AJ. Effect of aerosol and oral fenoterol on histamine and methacholine challenge in asthmatic subjects. *Thorax* 1981;36(8):580–4.
- [26] Cass LM, Efthymiopoulos C, Bye A. Pharmacokinetics of zanamivir after intravenous, oral, inhaled or intranasal administration to healthy volunteers. *Clin Pharmacokinet* 1999;36(1):1–11.
- [27] Qiu, R., Wei, X., Zhao, M., et al. (2020). Outcome reporting from protocols of clinical trials of Coronavirus Disease 2019 (COVID-19): a review. medRxiv.
- [28] Gentile I, Abenavoli L. COVID-19: perspectives on the potential novel global threat. *Rev Recent Clinic Trials* 2020;15(2):1.
- [29] Jia HP, Look DC, Shi L, et al. ACE2 receptor expression and severe acute respiratory syndrome coronavirus infection depend on differentiation of human airway epithelia. *J Virol* 2005;79(23):14614–21.
- [30] Zhang Z. Therapeutic effect of hydroxychloroquine on novel coronavirus pneumonia (COVID-19). <http://www.chictr.org.cn/showproj.aspx?proj=48880> [Accessed 14 Feb 2020].
- [31] Su, Z., & Wu, Y. (2020). A Multiscale and Comparative Model for Receptor Binding of 2019 Novel Coronavirus and the Implication of its Life Cycle in Host Cells. bioRxiv.
- [32] Devaux, C. A., Rolain, J. M., Colson, P., & Raoult, D. (2020). New insights on the antiviral effects of chloroquine against coronavirus: what to expect for COVID-19?. *Int J Antimicrobial Agents*, 105938.
- [33] Kearney, J. (2020). Chloroquine as a Potential Treatment and Prevention Measure for the 2019 Novel Coronavirus: A Review.
- [34] Keyaerts E, Vijgen L, Maes P, Neyts J, Van Ranst M. In vitro inhibition of severe acute respiratory syndrome coronavirus by chloroquine. *Biochem Biophys Res Commun* 2004;323(1):264–8.
- [35] McLachlan AJ, Tett SE, Cutler DJ, Day RO. Bioavailability of hydroxychloroquine tablets in patients with rheumatoid arthritis. *Rheumatology* 1994;33(3):235–9.