



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

journal homepage: [www.elsevier.com/locate/bjbas](http://www.elsevier.com/locate/bjbas)

## Full Length Article

# Prevalence of *Salmonella* and *E. coli* in neonatal diarrheic calves

F.R. El-Seedy<sup>a</sup>, A.H. Abed<sup>a,\*</sup>, H.A. Yanni<sup>b</sup>, S.A.A. Abd El-Rahman<sup>b</sup>

<sup>a</sup> Bacteriology, Department of Mycology and Immunology, Faculty of Veterinary Medicine, Beni-Suef University, Beni Suef, Egypt

<sup>b</sup> Animal Health Research Institute, Dokki, Egypt

### ARTICLE INFO

#### Article history:

Received 18 October 2015

Received in revised form 24

November 2015

Accepted 28 November 2015

Available online 2 March 2016

#### Keywords:

Neonatal calf diarrhea (NCD)

*Salmonella*

enterotoxigenic *E. coli* (ETEC)

### ABSTRACT

Neonatal calf diarrhea remains one of the most important problems faced by livestock, causing great economic losses. This study investigated the prevalence of *Salmonella* and *Escherichia coli*, especially enterotoxigenic *E. coli* (ETEC), in diarrheic calves. Fecal samples were collected from 127 diarrheic calves up to 3 months of age at 12 farms from different governorates in Egypt. 119 bacterial isolates (93.7%) were recovered and the prevalences of *Salmonella* and *E. coli* in diarrheic calves were 18.1% and 75.6%, respectively. Serotyping of *Salmonella* isolates revealed that *S. Enteritidis* and *S. Typhimurium* were the most prevalent serotypes, representing 60.9% and 30.4%, respectively, while *S. Dublin* was 8.7%. Serogrouping of *E. coli* isolates showed that 10 O-serogroups were obtained where O<sub>26</sub> and O<sub>103</sub> were the most prevalent (17.7% of each). *Salmonella* serotypes showed positive results with PCR test using oligonucleotide primer amplifying 521 bp fragment of *invA* gene of *Salmonella* while 70% of *E. coli* serogroups possessed ETEC virulent gene (K99). The *in-vitro* antibiotic sensitivity test indicated that *Salmonella* serotypes showed high sensitivity against enrofloxacin, spectinomycin and neomycin while *E. coli* isolates showed high sensitivities against marbofloxacin, spectinomycin and neomycin only.

© 2016 Beni-Suef University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Neonatal calf diarrhea (NCD) is one of the most common diseases in young animals, causing huge economic and productivity losses to bovine industry worldwide (Cho and Yoon, 2014). In 2007, The National Animal Health Monitoring System

(NAHMS) for U.S. dairy reported that 57% of unweaned calf mortality was due to diarrhea especially in calves less than one month old (USDA Dairy, 2007). In Egypt, NCD continues to be the 1st cause of calf mortality, which ranges between 27.4% and 55% of the total deaths in young calves (Ahmed, 1980). The economic losses occur not only from mortality but also from other costs including treatment, diagnostics, labor, veterinary

\* Corresponding author. Lecturer of Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef, 62512, Egypt. Tel.: +201148824447; fax: +20822327982.

E-mail addresses: [aboabedelmasry@yahoo.com](mailto:aboabedelmasry@yahoo.com), [ahmed.moawad@vet.bsuef.edu](mailto:ahmed.moawad@vet.bsuef.edu) (A.H. Abed).

<http://dx.doi.org/10.1016/j.bjbas.2015.11.010>

2314-8535/© 2016 Beni-Suef University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

intervention and decreased number of herd replacements as well as subsequent chronic ill thrift and impaired growth performance (Bazeley, 2003).

NCD is a multifactorial syndrome including pathogen (infectious NCD) as well as non-infectious factors related to the animal (immunological and nutritional status), the environment or the management (Izzo et al., 2011). Because of the multifactorial nature of NCD, it is difficult to be controlled effectively (Cho and Yoon, 2014).

Infectious diarrhea is the most significant cause of morbidity and mortality in neonatal dairy calves throughout the world and it can be caused by many pathogens including viruses (coronavirus and rotavirus), protozoa (*Cryptosporidium parvum*) and bacteria (Izzo et al., 2011). Among bacteria, enterotoxigenic *Escherichia coli* (ETEC) and *Salmonellae* are the most economically important pathogens (Achá et al., 2004), although other bacteria have also been identified as causes of enteric disease and NCD, e.g. *Campylobacter* species (Myers et al., 1984) and *Clostridium* species (Cho et al., 2010). Although co-infection is considerably observed in diarrheic calves, sometimes single infection is recorded. Farm geographic location, management practices, as well as herd size affect considerably the prevalence of the pathogen. (Cho and Yoon, 2014). Identification of the possible causative agent in outbreaks of diarrhea is important to allow targeted preventative measures, such as vaccination, and identification of possible risk factors or sources of infection (Izzo et al., 2011).

*Salmonella enterica* colonizes the digestive of both adult cattle and calves, but the infection is often recorded in the first 3 months of age and often causing severe symptoms (Fossler et al., 2005). *S. enterica* serovar *typhimurium* (*S. Typhimurium*) and serovar *dublin* (*S. Dublin*) are the most common causative agents of salmonellosis in cattle causing acute and systemic diarrheal diseases, respectively (Cho and Yoon, 2014). Diarrhea due to *Salmonella* infection is watery and mucoid with the presence of blood and fibrin (Fossler et al., 2005). Calves can shed *Salmonella* for variable periods of time and intermittently depending on the degree of infection (Cho and Yoon, 2014). Furthermore, after the disappearance of the organism from the intestinal tract, up to 5% of recovered animals may become carriers shedding the organism in their feces (Jay, 2000). Infected cattle and carriers can serve as source of infection for other animals or even human through food-borne routes or direct contact and so the determination of *Salmonella* strains in fecal samples is not only important for the diagnosis of salmonellosis, but also essential to identify the carriers (Warnick et al., 2003).

Among *E. coli* pathogroups, the most common cause of NCD is ETEC stains that produce the K99 (F5) adhesion antigen (*E. coli* K99+) and heat-stable (STa or STb) and/or heat-labile (LT1 or LT2) enterotoxins (Kaper et al., 2004). *E. coli* causes a watery diarrhea and weakness in 1–4 day old newborn calves. Death usually occurred within 24 hours due to severe dehydration (Cho et al., 2010). The fimbrial adhesion F5 (K99) promotes the adhesion of bacterial cells to glycoproteins on the epithelial surface of the jejunum and/or ileum, and bacterial enterotoxin also causes damage to the epithelial cells, resulting in fluid secretion and diarrhea (Acres, 1985).

The present study aimed to investigate the prevalence of *Salmonella* and *E. coli*, especially ETEC, in diarrheic calves in Egypt.

## 2. Material and methods

### 2.1. Samples

Fecal samples were collected every 4 weeks for a period of 5 months from the rectum of 127 cross-breed diarrheic calves up to 3 months of age at 12 farms from different governorates in Egypt along the period of April to November 2013. All samples were transferred in an ice box to the laboratory with minimal delay for bacteriological examination.

### 2.2. Bacteriological examination

One gram of each fecal sample was diluted in 3 mL of sterile saline. Samples were cultured and identified according to Quinn et al. (2002).

For isolation of *Salmonella* strains, a loopful from the diluted specimens was inoculated into selenite F broth with overnight incubation at 37 °C. Then, a loopful was streaked out onto MacConkey's agar, xylose lysine deoxycholate and *Salmonella*–*Shigella* agar media and incubated at 37 °C for 18–24 hours. Suspected colonies were subjected to biochemical testing according to Collee et al. (1996).

For isolation of *E. coli* strains, a loopful from the diluted specimens was inoculated into MacConkey's agar and incubated at 37 °C for 18–24 hours. Lactose fermenter (pink) colonies were streaked onto and eosin methylene blue agar and confirmed as *E. coli* using the standard biochemical tests according to Collee et al. (1996).

### 2.3. Serological identification

#### 2.3.1. Serotyping of *Salmonella*

*Salmonella* serovars were identified serologically by slide agglutination test using diagnostic polyvalent and monovalent O and H *Salmonella* antisera according to Kauffman–White scheme (Kauffmann, 1974).

#### 2.3.2. Serogrouping of *E. coli* isolates

*E. coli* serogroups were identified serologically by slide agglutination test using standard polyvalent and monovalent *E. coli* antisera according to Edwards and Ewing (1972).

### 2.4. Polymerase chain reaction

PCR was applied on 3 *Salmonella* isolates representing all the recovered serotypes as well as 10 *E. coli* isolates representing all the recovered serogroups. In case of *Salmonella*, oligonucleotide primers that amplify a 521 bp fragment for the *invA* (invasion A) gene in *Salmonella* serovars was used. Meanwhile, in case of *E. coli*, oligonucleotide primers that amplify a 314 bp fragment for the K99 (F5) gene in *E. coli* species was used (Table 1).

### 2.5. Antibiotic susceptibility testing

All *Salmonellae* (23 isolates) and *E. coli* (96 isolates) recovered from diarrheic calves were tested for their antimicrobial susceptibility to 12 different antimicrobial discs including

**Table 1 – Primers used in PCR.**

Gene	Primer Sequence 5'-3'	Size of product	Reference
invA(F)	TTGTTACGGCTATTTTGACCA	521 bp	Swamy et al. (1996)
invA(R)	CTGACTGCTACCTTGCTGATG		
F5(F)	TATTATCTTAGGTGGTATGG	314 bp	Frank et al. (1998)
F5(R)	GGTATCCTTAGCAGCAGTATTC		

enrofloxacin (10 µg), marbofloxacin (10 µg), gentamycin (10 µg), erythromycin (15 µg), cefotaxime sodium (30 µg), amoxicillin (10 µg), penicillin (10 µg), tetracycline (30 µg), streptomycin (10 µg), trimethoprim-sulphamethoxazol (1.25 + 23.75 µg), spectinomycin (20 µg) and neomycin (20 µg) (Oxoid, Basing Stoke, UK). Antimicrobial susceptibility testing was performed using disc diffusion method on Muller Hinton agar according to [Clinical and Laboratory Standards Institute \(CLSI\) \(2012\)](#). The antibiotic susceptibility was based on the induced inhibition zones according to the guidelines of the [CLSI \(2012\)](#).

### 3. Results

#### 3.1. Bacterial isolation

Out of 127 collected fecal samples from diarrheic calves, 119 (93.7%) bacterial isolates were recovered, including 23 (18.1%) *Salmonella* serovars and 96 (75.6%) *E. coli* strains ([Table 2](#)).

#### 3.2. Serological identification

##### 3.2.1. Serotyping of *Salmonella* isolates

The 23 *Salmonella* isolates were serotyped as 14 *S. Enteritidis* (60.9%), 7 *S. Typhimurium* (30.4%) and 2 *S. Dublin* (8.7%).

##### 3.2.2. Serogrouping of *E. coli* isolates

Out of 96 *E. coli* isolates, 10 O-serogroups were identified. The serogroups O<sub>26</sub> and O<sub>103</sub> were the most prevalent representing 17 isolates (17.7%) for each followed by serogroups O<sub>127</sub>; 14 isolates (14.6%) and O<sub>119</sub>; 13 isolates (13.6%). While the serogroups O<sub>86</sub>, O<sub>111</sub> and O<sub>157</sub> represented 5 isolates (5.2%) for each, meanwhile O<sub>44</sub> and O<sub>158</sub> serogroups represented 4 isolates (4.2%) and finally serogroup O<sub>78</sub>; 3 isolates (3.1%). Moreover, there were 9 (9.4%) isolates were untyped with the available antisera.

**Table 2 – Prevalence of *Salmonella* and *E. coli* in diarrheic calves.**

No. of samples	<i>Salmonella</i>		<i>E. coli</i>		Total		Negative isolation	
	No.	%	No.	%	No.	%	No.	%
127	23	18.1	96	75.6	119	93.7	8	6.3

#: was calculated according to the number of the collected samples.

### 3.3. PCR results

#### 3.3.1. PCR of *Salmonella* serovars

PCR results revealed that all the tested isolates confirmed morphologically, biochemically and serologically as being *Salmonella* showed positive results with PCR assay with oligonucleotide primer that amplified a 521 bp fragment of *invA* gene ([Fig. 1](#)).

#### 3.3.2. PCR of *E. coli* serogroups

PCR results revealed that out of the tested 10 *E. coli* serogroups, 7 serogroups (70%) possessed ETEC virulence gene (K99) including O<sub>26</sub>, O<sub>44</sub>, O<sub>78</sub>, O<sub>86</sub>, O<sub>111</sub>, O<sub>119</sub> and O<sub>127</sub> while O<sub>103</sub>, O<sub>157</sub> and O<sub>158</sub> not possessed K99 gene ([Fig. 2](#)).

### 3.4. Antibiotic susceptibility testing


Results of *in-vitro* sensitivity tests of *Salmonella* isolates against 12 antimicrobial agents revealed that *S. Typhimurium* and *S. Enteritidis* isolates showed high sensitivities against marbofloxacin, enrofloxacin, trimethoprim-sulphamethoxazol, spectinomycin and neomycin. On the other hand, high resistances were observed against amoxicillin, gentamycin, cefotaxime sodium, streptomycin, erythromycin, tetracycline and amoxicillin. Meanwhile, *S. Dublin* isolates were sensitive to enrofloxacin, cefotaxime sodium, penicillin, tetracycline, streptomycin, spectinomycin and neomycin. On the contrary, they were resistant to marbofloxacin, gentamycin, erythromycin, amoxicillin and trimethoprim-sulphamethoxazol. Concerning *E. coli* isolates, the different serogroups as well as the untyped group showed different degrees of sensitivity against the tested antibiotics. Mostly, *E. coli* isolates showed high resistances against the majority of antimicrobials, meanwhile, high sensitivity was observed against marbofloxacin, spectinomycin and neomycin only.

## 4. Discussion

Neonatal calf diarrhea remains as one of the most important problems faced by livestock, causing great economic losses. Calves are at greatest risk of diarrhea within the first month of life and the incidence of diarrhea decreases with age ([Garcia et al., 2000](#)).

The multifactorial nature of NCD makes this disease hard to control effectively. Therefore, prevention and control of such disease must be based on a good understanding of those disease complexities during the calving period before disease outbreaks ([Cho and Yoon, 2014](#)). Identification of the possible causative agent in outbreaks of diarrhea is important to allow targeted preventative measures, such as vaccination, and identification of possible risk factors or sources of infection ([Izzo et al., 2011](#)). *E. coli* and *Salmonella* are the most common identified pathogens in scouring calves less than 2 months of age ([Achá et al., 2004](#)). Their prevalences vary by geographical location of the farms, farm management practices, and herd size ([Cho and Yoon, 2014](#)).

In the present study, the prevalences of *E. coli* and *Salmonella* in neonatal diarrheic calf were 18.1% and 75.6%, respectively ([Table 2](#)). The negative bacterial isolation in some




**Fig. 1 – The PCR amplification of *invA* gene of *Salmonella* serovars showing positive amplicons at 521 bp. DNA size marker (M). Lane 1: positive control (*S. Typhimurium* strain). Lane 2: negative control. Lane 3: *S. Enteritidis*. Lane (4): *S. Typhimurium*. Lane (5): *S. Dublin*.**

fecal samples (6.3%) did not mean absence of microbial infection but may attribute to infection with bacteria requiring specific or enriched culture media and can't grow on the used culture media such as *Clostridium* species (Cho et al., 2010) and *Campylobacter* species (Myers et al., 1984) or viral infection; requiring tissue culture, or even protozoal infection. Fricker (1987) elucidated that bacteria were not detectable due to the number of colony forming units are below the detection limit of the assay (especially in case of *Salmonella* in feces). Additionally, the presence of antibiotic residues may explain falsely negative bacteriological results because the withdrawal time is not regarded in our herds.

The prevalence of *Salmonella* in the present study was similar to those of other studies in Egypt (Riad et al., 1998: 18.2%; Seleim et al., 2004: 17.5%; and Youssef and El-Haig, 2012: 18.66%) while higher than that reported by Haggag and Khaliel (2002) (4%), and Younis et al. (2009) (4.09%). On contrary, this result was much lower than that reported by Moussa et al. (2010) (43.53%). Varying prevalences of salmonellosis in calves were recorded in diarrheic calves in African countries such as Mozambique

(5%; Achá et al., 2004) and Algeria (66.6% in calves at the end of the first month; Akam et al., 2004), and other countries such as India (5%; Joon and Kaura, 1993) and Australia (23.8%; Izzo et al., 2011). Differences of the prevalence rates of *Salmonella* in diarrheic calves in comparison to the previous studies could be explained in the light of species and geographical locations and hygienic measures, and these factors significantly influence the prevalence of salmonellosis in calves (Younis et al., 2009).

The prevalence of *E. coli* in the current study had nearly coincided with the findings of Awad et al. (1979: 80%) and Haggag and Khaliel (2002: 82%), while higher than those of Azzam et al. (2006: 5.4%), El-Shehedi et al. (2013: 35.83%), Osman et al. (2013: 63.6%), and Hassan (2014: 50%), and lower than that obtained by Ibrahim (1995: 100%). In comparison to other countries, the present results were similar to that of Hemashenpagam et al. (2009) in India (75%), while it was higher than those obtained by Joon and Kaura (1993) in India (23%), Viring et al. (1993) in Sweden (11.5%), Khan and Khan (1997) in Pakistan (54%), Steiner et al. (1997) in Germany (42%), Bendali et al. (1999) in France



**Fig. 2 – The PCR amplification of K99 virulence gene of *E. coli* showing positive amplicons at 314 bp. M: DNA size marker (100–1000 bp). Lane 1: Negative control. Lane 2: Positive control (*E. coli* serogroup O<sub>86</sub>). Lane (3–12): The tested serogroups (O<sub>26</sub>, O<sub>44</sub>, O<sub>78</sub>, O<sub>86</sub>, O<sub>103</sub>, O<sub>111</sub>, O<sub>119</sub>, O<sub>127</sub>, O<sub>157</sub> and O<sub>158</sub>, respectively).**

(20.3%), [Valdivia-Andy et al. \(2000\)](#) in Mexico (63.7%), [Oporto et al. \(2008\)](#) in Northern Spain (35.9%), and [Izzo et al. \(2011\)](#) in Australia (17.4%). On contrary, this result was lower than that of [Pourtaghi et al. \(2013\)](#) in Iran (86.7%). The differences of the prevalence rates of *E. coli* in diarrheic calves may be attributed also to the geographical locations and management practice as well as hygienic measures where ETEC infection occurs mainly through ingestion of contaminated food or water ([Cho and Yoon, 2014](#)).

In the current study, serotyping of *Salmonella* isolates revealed that *S. Enteritidis* (SE) and *S. Typhimurium* (ST) were the most prevalent serotypes with 60.9% and 30.4%, respectively, while *S. Dublin* (SD) represented 8.7%. *Salmonella* serovars isolation frequencies vary from one location to another due to different management and hygienic measures as well as geographical, environmental and individual variations ([Veling et al., 2002](#)). The predominance of ST and SE serovars among diarrheic calves detected in the present study was supported by many previous reports in Egypt ([Seleim et al., 2004](#); [Younis et al., 2009](#); [Moussa et al., 2010](#)) and [Youssef and El-Haig, 2012](#)). In addition; this finding substantiated the reports from the other countries ([Murray, 1994](#), in Australia, and [Smith-Palmer et al., 2003](#), in Scotland).

*S. Typhimurium* and *S. Enteritidis* are non-host-adapted serovars and so they are less likely to establish a "carrier state" in the recovered animals. Infection with a non-host-adapted *Salmonella* results in transient shedding of organisms (3–16 weeks). The most common sources of infection are contaminated food and water. Approximately 40% of all animal-origin feed additives (bone or fish meals) are contaminated with *Salmonella*. Human sewage has also been tracked down as a source of infection in some herd outbreaks. Rodents, birds, and other animals spread infection through their feces and their carcasses ([Donaldson et al., 2006](#)). On contrary, *S. Dublin* is host-adapted in cattle, and commonly produces a carrier state in the recovered animals. It can survive intracellularly inside the mononuclear phagocytic cells in the face of high serum neutralizing antibody titers. Intermittent shedding occurs, which is maximized by stress factors. In outbreaks of *S. Dublin* infection, carriers are the most likely source. These asymptomatic carriers can shed organisms in their feces and milk for the remainder of their lives. Calves, which are fed raw, contaminated milk from the bulk tank, can be infected by this route ([Lance et al., 1992](#)).

Serogrouping of *E. coli* isolates showed that 10 O-serogroups were identified; of them O<sub>26</sub> and O<sub>103</sub> were the most prevalent groups (17.7% each) followed by O<sub>127</sub> (14.6%) and O<sub>119</sub> (13.6%). These results run hand to hand with those of [Ibrahim \(1972\)](#) and [Abd-Elrahman \(2013\)](#). The present findings were also nearly similar to that obtained by [El-Shehedi et al. \(2013\)](#) and [Osman et al. \(2013\)](#). Concerning the other countries, [Tamaki et al. \(2005\)](#) in Japan found that the most common *E. coli* serotypes isolated from diarrheic fecal samples of calves were O<sub>119</sub>, O<sub>111</sub>, O<sub>126</sub>, and O<sub>78</sub>. Similarly, [Badouei et al. \(2010\)](#) recovered O<sub>157</sub>:H<sub>7</sub>, O<sub>111</sub>, and O<sub>26</sub> serotypes from diarrheic and non-diarrheic calves and the most common serogroups was O<sub>26</sub> (18.4%).

*Salmonella invA* gene has become one of the most popular PCR target sequences ([Rahn et al., 1992](#)) and its amplification now has been recognized as an international standard for detection of *Salmonella* ([Malorny et al., 2003](#)). The *invA* gene

encodes a protein in the inner membrane of bacteria, which is responsible for invasion to the epithelial cells of the host ([Darwin and Miller, 1999](#)).

In the present study, PCR was applied on 3 isolates of *Salmonella* representing the recovered serovars to determine the virulence *invA* gene. The results revealed that all the tested isolates of *Salmonella* showed positive results with PCR assay using oligonucleotide primer that amplified a 521 bp fragment of *invA* gene of *Salmonella*. These results were similar to those obtained by [Soliman \(2014\)](#), who determined the virulence *invA* gene in all recovered *Salmonella* serogroups from bovine and ovine; *S. Typhimurium*, *S. Enteritidis*, *S. Virchow*, *S. Montividio* and *S. Rubislow*, by PCR.

Concerning *E. coli*, PCR was applied on 10 isolates representing all the recovered serogroups. Results revealed that 7 serogroups (70%) possessed ETEC virulent gene (K99) including O<sub>26</sub>, O<sub>44</sub>, O<sub>78</sub>, O<sub>86</sub>, O<sub>111</sub>, O<sub>119</sub> and O<sub>127</sub> while O<sub>103</sub>, O<sub>157</sub> and O<sub>158</sub> not possessed K99 gene. Although [Sherwood et al. \(1983\)](#) confirmed the close correlation between enterotoxigenicity and the K99 antigen presence and [Pourtaghi et al. \(2013\)](#) reported that most bovine ETEC produce K99 fimbriae, although [Moon et al. \(1976\)](#) have reported K99 antigen in non-enterotoxigenic *E. coli*.

Despite the increased availability of vaccines against ETEC and other pathogens associated with NCD and continued emphasis on optimizing colostral transfer of passive immunity, improved treatment protocols for calf diarrhea are necessary. Although the administration of intravenous fluids and oral electrolyte solutions plays the main role in treatment, the efficacy of antimicrobial drugs in treating calf diarrhea is argumentative ([Constable, 2004](#)). Diarrheic calves are more likely to have failure or partial failure of passive transfer, and so they are more likely to be bacteraemic ([Constable, 2004](#)) and this is an additional cause that antimicrobial agents might be indicated in the treatment of calf diarrhea. The type of antibiotic drug should better be selected on the basis of its sensitivity, which could be detected by laboratory examination, and the antimicrobial treatment of diarrheic calves should therefore be focused against bacteria in the two sites of infection: the small intestine and the blood ([Constable, 2004](#)).

In the present study, the results of *in-vitro* antibiotic sensitivity test showed that the different serotypes of *Salmonella* showed different degrees of sensitivity against the tested antibiotics but in general they all showed high sensitivities against enrofloxacin, spectinomycin and neomycin. On contrary, they were resistant to gentamycin, erythromycin and amoxicillin. On the other hand, the different *E. coli* serogroups as well as the untyped group showed different degrees of sensitivity against the tested antibiotics. Mostly, *E. coli* isolates showed high resistances against the majority of antimicrobials, meanwhile, high sensitivities were observed against marbofloxacin, spectinomycin and neomycin only. The high occurrence of resistance can be predicted since a large proportion of the animals are probably treated with antimicrobial drugs ([de Verdier et al., 2012](#)). [Call et al. \(2008\)](#) discussed the epidemiology of resistant *E. coli* in calves as multifactorial, complex and e.g. influenced by co-selection due to linkage of resistance genes. [Walk et al. \(2007\)](#) supposed that regardless of use of antimicrobials, antibiotic resistance in *E. coli* is co-selected in calves by an unknown "beneficial mutation."

## 5. Conclusion

Neonatal calf diarrhea remains one of the most important problems faced by livestock, causing great economic losses. The prevalences of *Salmonella* and *E. coli* in diarrheic calves were 18.1% and 75.6%, respectively. *S. Enteritidis* and *S. Typhimurium* were the most prevalent *Salmonella* serotypes while 70% of *E. coli* was ETEC.

## REFERENCES

- Abd-Elrahman AH. Colibacillosis in newly born buffalo calves and role of lacteal fort in preventing recurrence of calf diarrhea. *Life Sci J* 2013;8(4).
- Achá SJ, Kuhn I, Jonsson P, Mbazima G, Katouli M, Mollby RS. Studies on calf diarrhoea in Mozambique: prevalence of bacterial pathogens. *Acta Vet Scand* 2004;45:27-36.
- Acres SD. Enterotoxigenic *Escherichia coli* infections in newborn calves: a review. *J Dairy Sci* 1985;68:229-56.
- Ahmed AA. Calf sources in Egyptian buffalo-cows. Egyptian German Seminar on the Mortality of Newly-born Calves, 1980; Pp. 19-21.
- Akam A, Khelef D, Kaidi R, Othmani A, Lafri M, Tali-Maamar H, et al. Frequency of *Cryptosporidium parvum*, *Escherichia coli* K99 and *Salmonella* spp. isolated from healthy and unhealthy calves in six breeding farms from Mitidja, Algeria (preliminary results). *Rev Sci Parasitol* 2004;5:13-21.
- Awad FI, Farrag I, Shawkat ME, Abeid MH. Studies on enterotoxaemia in young buffalo calves. *Egypt J Vet Sci* 1979;14(1):24-9.
- Azzam RA, Hassan WH, Ibrahim MA, Khaled MS. Prevalence of verocytotoxigenic *E. coli* O157: H7 in cattle and man in Beni-Suef Government. *Alex J Vet* 2006;24(1):111-22.
- Badouei MA, Salehi TZ, Khorasgani MR, Tadjbakhsh H, Brujeni GN, Nadalian MG. Virulence gene profiles and intimin subtypes of Shiga toxin-producing *Escherichia coli* isolated from healthy and diarrhoeic calves. *Vet Rec* 2010;167(22):858-61.
- Bazeley K. Investigation of diarrhoea in the neonatal calf. *In Pract* 2003;25:152-9.
- Bendali F, Bichet H, Schelcher F, Sanaa M. Pattern of diarrhoea in new born calves in South west France. *Vet Res* 1999;30:61-74.
- Call DR, Davis MA, Sawant AA. Antimicrobial resistance in beef and dairy cattle production. *Anim Health Res Rev* 2008;9:159-67.
- Cho Y, Yoon KJ. An overview of calf diarrhea-infectious etiology, diagnosis, and intervention. *J Vet Sci* 2014;15(1):1-17.
- Cho Y, Kim W, Liu S, Kinyon JM, Yoon KJ. Development of a panel of multiplex real-time polymerase chain reaction assays for simultaneous detection of major agents causing calf diarrhea in feces. *J Vet Diagn Invest* 2010;22:509-17.
- Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial disk susceptibility tests, M02-A11 vol. 32. Approved Standard - 11th ed. 2012.
- Collee JG, Fraser AG, Marmion BP, Simmons A. Practical medical microbiology. 14th ed. 1996.
- Constable PD. Review: antimicrobial use in the treatment of calf diarrhea. *J Vet Intern Med* 2004;18:8-17.
- de Verdier K, Nyman A, Greko C, Bengtsson B. Antimicrobial resistance and virulence factors in *Escherichia coli* from Swedish dairy calves. *Acta Vet Scand* 2012;54:2.
- Darwin KH, Miller VL. Molecular basis of the interaction of *Salmonella* with the intestinal mucosa. *Clin Microbiol Rev* 1999;12:405-28.
- Donaldson SC, Straley BA, Hegde NV, Sawant AA, DebRoy C, Jayarao BM. Molecular epidemiology of ceftiofur-resistant *Escherichia coli*: isolates from dairy calves. *Appl Environ Microbiol* 2006;72(6):3940-8.
- Edwards PR, Ewing NH. Identification of enterobacteriaceae. 3rd ed. Atlanta, GA: Burgeon Publishing, Co.; 1972. p. 208-339.
- El-Shehedi MA, Eraqi MM, Ali AR. Characterization of *Escherichia coli* from diarrheic calves with special reference to plasmid profile. *J Am Sci* 2013;9(7).
- Fossler CP, Wells SJ, Kaneene JB, Ruegg PL, Warnick LD, Bender JB, et al. Herd-level factors associated with isolation of *Salmonella* in a multi-state study of conventional and organic dairy farms II. *Salmonella* shedding in calves. *Prev Vet Med* 2005;70:279-91.
- Frank SM, Bosworth BT, Moon HW. Multiplex PCR for enterotoxigenic, attaching and effacing, and Shiga toxin-producing *Escherichia coli* strains from calves. *J Clin Microbiol* 1998;36:1795-7.
- Fricker CR. The isolation of salmonellas and campylobacters. *J Appl Bacteriol* 1987;63:99-116.
- García A, Ruiz-Santa-Quiteria JA, Orden JA, Cid D, Sanz R, Gómez-Bautista M, et al. Rotavirus and concurrent infections with other enteropathogens in neonatal diarrheic dairy calves in Spain. *Comp Immunol Microbiol Infect Dis* 2000;23:175-83.
- Haggag YN, Khaliel SA. Public health importance of certain bacteria isolated from calves and small ruminants. 2<sup>nd</sup> Vet Cong, Fac Vet Med, Minufiya Univ, Egypt 2002;2(1):173-84.
- Hassan AM. Some studies on bacteriological causes of enteritis in calves. *J Vet Adv* 2014;4(5):503-7.
- Hemashenpagam N, Kiruthiga B, Selvaraj T, Panneerselvam A. Isolation, identification and characterization of bacterial pathogens causing calf diarrhea with special reference to *Escherichia coli*. *Internet J Microbiol* 2009;7(2).
- Ibrahim AM. Sanitary studies on newly born calves. [Ph. D. thesis (Animal Hygiene)], Faculty of Veterinary Medicine, Suez Canal University, Egypt; 1995.
- Ibrahim MS. Bacteriological studies of colibacillosis in calves. [M.V.Sc. thesis (Microbiology)], Faculty of Veterinary Medicine, Cairo University, Egypt; 1972.
- Izzo MM, Kirkland PD, Mohler VL, Perkins NR, Gunna AA, House JK. Prevalence of major enteric pathogens in Australian dairy calves with diarrhea. *Aust Vet J* 2011;89(5):167-73.
- Jay JM. Foodborne gastroenteritis caused by *Salmonella* and *Shigella*. In: *Modern food microbiology*. 2000. p. 511-28.
- Joon DS, Kaura YK. Isolation and characterization of some of the enterobacteria from diarrhoeic and non-diarrhoeic calves. *Ind J Anim Sci* 1993;63:373-83.
- Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2004;2:123-40.
- Kauffmann F. Kauffmann White scheme WHO-BD/72, 1, Rev. 1. *Acta Path Microbiol Scand* 1974;61:385-7.
- Khan A, Khan M. Bacteria isolated from natural cases of buffalo and bovine neonatal calf diarrhoea, pneumonia and pneumoenteritis. *Vet Arch* 1997;67(4):161-7.
- Lance SE, Miller GY, Hancock DD, Bartlett PC, Heider LE. *Salmonella* infections in neonatal dairy calves. *J Am Vet Assoc* 1992;201(6):864-8.
- Malorny B, Tassios PT, Rådström P, Cook N, Wagner M, Hoorfar J. Standardization of diagnostic PCR for the detection of food borne pathogens. *Int J Food Microbiol* 2003;83:39-48.
- Moon HW, Whipp SC, Skartvedt SM. Etiologic diagnosis of diarrheal diseases of calves: frequency and methods for detecting enterotoxin and K99 production by *Escherichia coli*. *Am J Vet Res* 1976;37:1025-9.
- Moussa IM, Ashgan MH, Mahmoud MH, Mohamed KFH, Al-Doss AA. Rapid detection of *Salmonella* species in new-borne calves by polymerase chain reaction. *Int J Gen Mol Biol* 2010;62-6.

- Murray CJ. *Salmonella* serovars and phage types in humans and animals in Australia 1987–1992. *Aust Vet J* 1994;71: 78–81.
- Myers LL, Firehammer BD, Border MM, Shop DS. Prevalence of enteric pathogens in the feces of healthy beef calves. *Am J Vet Res* 1984;45:1544–8.
- Oporto B, Esteban JI, Aduriz G, Juste RA, Hurtado A. *Escherichia coli* O<sub>157</sub>:H<sub>7</sub> and non- O157 Shiga toxin-producing *E. coli* in healthy cattle, sheep and swine herds in Northern Spain. *Zoonoses Pub Health* 2008;52:411–550.
- Osman KM, Mustafa AM, Elhariri M, Abdelhamed GS. The distribution of *Escherichia coli* serovars, virulence genes, gene association and combinations and virulence genes encoding serotypes in pathogenic *E. coli* recovered from diarrhoeic calves, sheep and goat. *Transbound Emerg Dis* 2013;60(1):69–78.
- Pourtaghi H, Dahpahlavan V, Momtaz H. Virulence genes in *Escherichia coli* isolated from calves with diarrhoea in Iran. *Comp ClinPathol* 2013;22:513–15.
- Quinn PJ, Markey BK, Carter ME, Donnelly WJC, Leonard FC, Maguire D. *Veterinary microbiology and microbial disease*. Blackwell; 2002. p. 113–16.
- Rahn K, Grandis SA, Clarke RC, McEwen SA, Galan JE, Ginocchio C, et al. Amplification of an *invA* gene sequence of *Salmonella* Typhimurium by polymerase chain reaction as specific method of detection of *Salmonella*. *Mol Cell Probes* 1992; 6:271–9.
- Riad EM, Tanios AI, El-Moghny AFA. Antigen capture ELISA technique for rapid detection of *Salmonella typhimurium* in fecal samples of diarrheic cow calves. *Assiut Vet Med J* 1998;39(78):324–33.
- Seleim RS, Sahar RM, Novert MH, Gobran RA. *Salmonella* infection in calves: virulence proteins and its immunogenic properties. *J Vet online* 2004; <<http://www.priory.com/vet/salmonella.htm>>.
- Sherwood D, Snodgrass DR, Lawson GHK. Prevalence of enterotoxigenic *Escherichia coli* in calves in Scotland and northern England. *Vet Rec* 1983;113:208–12.
- Smith-Palmer A, Stewart WC, Mather H, Greig A, Cowden JM, Reilly WJ. Epidemiology of *Salmonella enterica* serovars enteritidis and typhimurium in animals and people in Scotland between 1990 and 2001. *Vet Rec* 2003;153: 517–20.
- Soliman HA. Conventional and advanced techniques for identification of bovine and ovine salmonellae. [Ph.D. thesis (Microbiology)], Faculty of Veterinary Medicine, Beni-Suef University, Egypt; 2014.
- Steiner I, Busato A, Burnen SA, Gaillard C. Frequency and aetiology of calf losses and calf diseases before weaning in cow calf farms. II. Microbiological and Parasitological diagnosis in diarrhoeic calves. *Dtch Tieraztl Wochenschr* 1997;104(5):169–73.
- Swamy SC, Barnhard HM, Lee HD, Dresen DW. Virulence determinant *invA* and *spvC* in salmonella isolated from poultry product, waste water, and human sources. *App Environ Microbiol* 1996;62:3768–71.
- Tamaki Y, Narimatsu H, Miyazato T, Nakasone N, Toma C, Iwanaga M. The relationship between O antigens and pathogenic genes of diarrhea-associated *E. coli*. *Jpn J Infect Dis* 2005;58:65–9.
- USDA Dairy. Part II: changes in the U.S. dairy cattle industry, 1991–2007. Pp. 57–61, USDA-APHIS-VS, CEAH, Fort Collins, February 2008. <<http://nahms.aphis.usda.gov/dairy/index.htm>>; 2007.
- Valdivia-Andy G, Rosales C, Soriano-becerrili DM, Alba-Hurtado F, Montaraz-Crespo JA, Tortora-Prez JL. Interaction of *E. coli* verocytotoxin strains and rotavirus in outbreak of calf's diarrhoea. *Vet Mex* 2000;31:293–300.
- Veling J, Barkema HW, van der Schans J, van Zijderveld F, Verhoeff J. Herd level diagnosis for *Salmonella enterica* subsp. Enterica serovars Dublin infection in bovine dairy herds. *Prev Vet Med* 2002;53:31–42.
- Viring S, Olsson SO, Alenius S, Emanuelsson U, Jacobsson SO, Larsson B, et al. Studies of enteric pathogens and gamma-globulin levels of neonatal calves in Sweden. *Acta Vet Scand* 1993;34(3):271–9.
- Walk ST, Mladonicky JM, Middleton JA, Heidt AJ, Cunningham JR, Bartlett P, et al. Influence of antibiotic selection on genetic composition of *Escherichia coli* populations from conventional and organic dairy farms. *Appl Environ Microbiol* 2007;73:5982–9.
- Warnick LD, Kanistanon K, McDonough PL, Power L. Effect of previous antimicrobial treatment on fecal shedding of *Salmonella enterica* subsp. Enterica serogroup b in New York dairy herds with recent clinical salmonellosis. *Prev Vet Med (Praha)* 2003;56:285–97.
- Younis EE, Ahmed AM, El-Khodery SA, Osman SA, El-Naker YF. Molecular screening and risk factors of enterotoxigenic *Escherichia coli* and *Salmonella* spp. in diarrheic neonatal calves in Egypt. *Res Vet Sci* 2009;87:373–9.
- Youssef AI, El-Haig MM. Herd problems and occupational zoonoses of *Salmonella enterica* serovars Typhimurium and Enteritidis infection in diarrheic cattle and buffalo calves in Egypt. *Int J Bioflux Soc* 2012;4(3):118–23.