




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com/en



ORIGINAL ARTICLE

Aminopeptidase N (APN/CD13) as a target molecule for scirrhous gastric cancer



Shigeo Nohara^a, Kazunori Kato^{b,c,*}, Daisuke Fujiwara^a,
Naoya Sakuragi^b, Kazuyoshi Yanagihara^d, Yoshimi Iwanuma^a,
Yoshiaki Kajiyama^a

^a Department of Esophageal and Gastroenterological Surgery, Juntendo University School of Medicine, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan

^b Department of Biomedical Engineering, Toyo University, 2100 Kujirai, Kawagoe, Saitama 350-8585, Japan

^c Atopy Research Center, Juntendo University School of Medicine, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan

^d Division of Translational Research, Exploratory Oncology and Clinical Trial Center, National Cancer Center Hospital East, 6-5-1 Kashiwano-ha, Kashiwa, Chiba 277-8577, Japan

Available online 13 January 2016

Summary

Background: Scirrhous gastric cancer is associated with peritoneal dissemination and advanced lymph node metastasis from an early stage, and the prognosis is still poor. In this study, we aimed to analyze candidate molecules for targeted therapy of scirrhous gastric cancer. We searched for molecules/metabolic activity that might be predominantly expressed in a subpopulation of scirrhous gastric cancer cells and might function as cancer stem cell markers.

Results: For this purpose, we investigated the expression of various cell surface markers and of aldehyde dehydrogenase (ALDH) activity. These analyses showed that the scirrhous gastric cancer cell lines HSC-58 and HSC-44PE heterogeneously expressed CD13, while CD44, CD133, EpCAM and ABCG2 were expressed uniformly. Moreover, 10% of the total HSC-58 cell population expressed ALDH enzyme activity. A subpopulation of cells strongly positive for ALDH also expressed high levels of CD13, both of which are known as cancer stem cell markers. HSC-58 cells expressing high levels of CD13 showed lower sensitivity to a cancer drug cisplatin than cells with low levels of CD13. In contrast, CD13^{high} subpopulation of HSC-58 was more sensitive to an aminopeptidase N inhibitor bestatin. In terms of antibody-drug therapy, anti-CD13-immunotoxin was highly cytotoxic towards HSC-58 cells and was more cytotoxic than anti-EpCAM-immunotoxin.

* Corresponding author. Department of Biomedical Engineering, Toyo University, 2100 Kujirai, Kawagoe, Saitama 350-8585, Japan.
E-mail address: kzkato@juntendo.ac.jp (K. Kato).

Conclusion: These data suggest that CD13 is a suitable cell surface candidate for targeted antibody-drug therapy of scirrhous gastric cancer.

© 2015 Elsevier Masson SAS. All rights reserved.

Introduction

Although the prevalence of gastric carcinoma has recently shown a gradual decrease, this cancer still accounts for a significant proportion of cancer-related deaths in Japan. Early detection and prevention of metastasis of this cancer is essential in order to improve the cure rate. Scirrhous gastric cancer, known as diffusely infiltrative carcinoma, Borrmann's type-IV carcinoma, or linitis plastica-type carcinoma, is characterized clinically as having the worst prognosis among the various types of gastric cancer, because it is frequently associated with metastases to lymph nodes and peritoneal dissemination. Scirrhous gastric cancer, which represents approximately 10% of gastric carcinomas, shows a 5-year survival rate of less than 17%, compared with 35 to 70% for other types of gastric cancers [1]. Various treatments, such as chemotherapy, hormonal therapy, hyperthermia, and immunotherapy have been tested for effectiveness against peritoneal metastasis from scirrhous gastric cancer, but the results have been unsatisfactory [2]. Chemotherapy plays a significant role in the treatment of such gastric cancer. However, the prognosis of advanced gastric cancer is still poor and treatment is usually unsuccessful. There is therefore an urgent need for development of an effective treatment for such patients.

Stem cells constitute the source of cell populations in the tissues of each organ and exhibit self-renewal ability and multi-differentiation potential. In the case of cancer, a subpopulation of cancer cells that exhibit the properties of stem cells are also present in cancer tissues. These so-called cancer stem cells (CSCs) were first reported in acute leukemia [3], and it later became clear that CSCs are also present in solid tumors such as breast cancer and brain tumors [4,5]. CSCs have been proposed to be the root cause of cancer growth and to be drug-resistant. Attempts to identify CSCs have been made based on their expression of cell surface molecules or on their intracellular metabolic activity, by analysis of both primary tumor specimens and established cell lines. Currently, there are reports of definitive CSC markers in digestive cancers such as colorectal cancer, liver cancer, and pancreatic cancer [6–8]. Among various cell surface markers for the identification of CSCs, Haraguchi et al. identified CD13 as functional marker that can be used to identify potentially dormant liver CSCs [9]. However, there have been few clear definitive CSC markers in upper gastrointestinal cancers such as in esophagus or stomach cancers.

The first aim of this study was therefore to determine the contribution of CD13 in scirrhous gastric cancer cell lines. We analyzed these cells for CD13 and other cell surface antigens and metabolic enzyme activity that were considered as possible candidate CSC markers by using flow cytometry

[10]. The second aim of this study was to assess whether antibody-mediated drug/toxin conjugates (immunotoxins) targeted towards CD13 on scirrhous gastric cancers, would show enhanced antitumor effect against scirrhous gastric cancer.

Materials and methods

Cell lines and cell culture

The scirrhous gastric cancer cell lines HSC-58, HSC-44PE, 58As1 and 44As3 were previously established from the ascitic fluid of patients with scirrhous gastric carcinoma [11]. The stomach adenocarcinoma MKN-7, MKN-45 and signet-ring cell carcinoma KATO-III were obtained from JCRB Cell Bank (Osaka, Japan) [12]. These cells were maintained in RPMI1640 medium (Sigma, St. Louis, MO) supplemented with 10% fetal calf serum (Gibco Life Technologies; Grand Island, NY), antibiotics and sodium pyruvate (Gibco). The cultures were maintained at 37 °C in an atmosphere of 5% CO₂ in air, and were propagated as low adhesion cultures at the bottom of plastic dishes.

Flow cytometric analysis

The following antibodies were used for flow cytometric analysis of cells: anti-CD24-FITC, anti-CD44-FITC, anti-CD90-FITC, anti-CD133-PE, anti-MDR1-PE, anti-EpCAM-PE, anti-CD13-PE (all from eBioscience, San Diego, CA) and anti-CDCP1-FITC and anti-ABCG2-PE (both from BioLegend, San Diego, CA). Cells were incubated with the indicated antibodies for 60 min and were then washed twice with PBS containing 2% FCS. Flow cytometric analysis was performed using a FACS-Calibur (BD Immunocytometry Systems, Franklin Lakes, NJ).

Aldehyde dehydrogenase assay

Aldehyde dehydrogenase (ALDH) activity in the cancer cells was assayed using the ALDEFLUOR[®] substrate according to the manufacturer's protocol (StemCell technologies, Vancouver, Canada). Specimens that were analyzed for ALDH activity were counter-stained with anti-CD13-PE at the appropriate dilution. Non-viable cells were eliminated using propidium iodide (Sigma). The specific flow gates for ALDH positive cells were set using a control sample of the isolated tumor cells in which ALDH activity was inhibited with diethylaminobenzaldehyde (DEAB). Subsequent flow cytometry runs were used to identify populations of cells with




Figure 1 Flow cytometric analysis of the expression of various cell surface molecules on human scirrhous and adenocarcinoma gastric cancer cell lines. A. Expression of CD24, CD90, CD44 and CDCP1 (upper panels) and CD133, MDR1, ABCG2, and CD13 (lower panels) on scirrhous gastric cancer HSC-58 are analyzed by flow cytometry and shown as histograms. Open histograms indicate the cells stained with mouse isotype control IgG1. And shaded histograms show the cells stained with various antibodies as indicated. B. Expression of CD13 on gastric cancer cell lines, HSC-58, 58As1, HSC-44PE and 44As3 (scirrhous cancers; upper panels) and KATO-III, MKN-7 and MKN-45 (adenocarcinomas; lower panels). C. Expression of ABCG2 on scirrhous gastric cancer cell lines are shown as histograms.




Figure 2 Characterization of aldehyde dehydrogenase (ALDH) activity and CD13 expression of HSC-58 cells. ALDH expression was assayed in the human gastric cancer cell lines HSC-58 (left panels) and HSC-44PE (middle panels), and in the stomach signet-ring cell carcinoma cell line KATO-III (right panels) by staining with the ALDH substrate ALDEFLUOR® (BAAA; BODIPY® aminoacetaldehyde) in the presence or absence of the ALDH inhibitor diethylaminobenzaldehyde (DEAB), followed by FACS analysis. Horizontal and vertical axes denote expression intensity.

towards HSC-58 cells that had bound anti-CD13 Ab, in a saporin-conjugated antibody dose-dependent manner. However, the saporin-conjugated second antibody was not cytotoxic towards HSC-58 cells to which the anti-EpCAM Ab was bound. These data show that CD13 is a suitable cell surface candidate for targeted antibody-drug therapy of scirrhus gastric cancer.

Discussion

Numerous reports have shown that a small subset of cancer cells bears drug-resistance and stem cell properties and that these cells are referred to as cancer stem or stem-like cells (CSCs), or as tumor-initiating cells (TICs). CSC identification was first attempted in acute leukemia, and was then later attempted for many types of cancers, however, no CSCs have been validated in scirrhus gastric cancer cell lines. Li et al. reported that CSCs could be identified in pancreatic cancers based on the high expression of CD44, CD24 and ESA on their cell surface [8]. Other

studies have shown that CD133 is a marker for CSCs of breast, colon, lung, ovary, pancreas and prostate cancers [6,23–27]. In addition to these markers, overexpression of cell surface transport proteins including MDR-1 and ABCG2 are related to the multidrug-resistance property of CSCs [28,29]. We therefore tried to identify CSCs in the scirrhus gastric cancer cell line HSC-58 using antibodies against various cell surface CD antigens such as CD24, CD44, CD90, CD133, CD13, ABCG2 and MDR-1, using flow cytometry. Unexpectedly, most of these cell surface markers were either not expressed or were uniformly expressed, except for CD13 that was heterogeneously expressed. The heterogeneous expression of CD13 was also found in HSC-58 and HSC-44PE cells. Interestingly, highly metastatic cell line 58As1 (derived from parental HSC-58) caused fatal cancerous peritonitis and bloody ascites expressed extremely high levels of CD13. In contrast, KATO-III with weak tumorigenicity in nude mice lacked the expression of CD13. It is anticipated that tumorigenic properties of scirrhus gastric cancer cell lines might be related with the CD13^{high} population.




Figure 3 ALDH activity and CD13 expression of HSC-58 cells. A. HSC-58 cells were analyzed for ALDH activity as in the legend of Fig. 2, and were also analyzed by FACS for CD13 expression. Horizontal axes denote the expression intensity of ALDH activity, the vertical axes denote CD13 expression (upper panels). The shaded histogram indicates CD13 expressing cells with low ALDH activity. The open histogram (thick line) indicates CD13 expressing cells with high ALDH activity. B, C. Drug sensitivity test of HSC-58 cells sorted into CD13^{high} (open circles) and CD13^{low} populations (closed circles). HSC-58 cells were separated by AutoMACS and sorted cells are exposed to the indicated concentrations of cisplatin (B) or bestatin (C) for 5 days following which viable cells were counted by fluorescence analysis of reduction of the AlamarBlue[®] reagent. Data are analyzed and represented by GraphPad Prism ver. 4.0 software.




Figure 4 Targeted delivery of an immunotoxin into HSC-58 cells. A. Expression of CD13 and EpCAM on HSC-58 cells as analyzed by flow cytometry. B. Cytotoxicity of a saporin-conjugated second antibody targeted against anti-CD13, anti-EpCAM antibody or control IgG bound to HSC-58 cells was assayed using fluorescence analysis of reduction of the dye AlamarBlue®. The horizontal axis denotes the concentration of the saporin immunotoxin. The vertical axis indicates cell survival. C. Phase-contrast images of representative cells treated with isotype control IgG (negative control; top right), isotype control IgG plus saporin-conjugated second antibody (bottom left) and anti-CD13 IgG plus saporin-conjugated second antibody (bottom right) for 5 days.

Another commonly-used cancer and normal hematopoietic stem cell marker is ALDH, which is an intracellular enzyme that is involved not only in retinoic acid and ethanol metabolism but also in resistance to alkylating cancer drugs [30]. It has been shown that high ALDH expression correlates with tumorigenic potential and poor prognosis in breast cancer patients [10]. We therefore determined expression of the metabolic marker ALDH in HSC-58 cells. Previous reports have shown that expression of CD44, or of an alternatively spliced variant of CD44 (CD44R), and of ABCG2 strongly contribute to cellular resistance against various anticancer drugs [31,32]. Based on our data, it is unlikely that ALDH activity is correlated with the expression of these cell surface molecules in the scirrhous gastric cancer cell line HSC-58. However, we observed that approximately 10% of the HSC-58 cells expressed high levels of ALDH activity. This cell population that was enriched for ALDH activity was almost identical to the CD13^{high} population. A significantly greater number of the CD13^{high} and ALDH^{high} cell population of HSC-58 survived chemotherapy with CDDP relative to the CD13^{low} and ALDH^{low} cell population. In our preliminary data, CD13 expression on HSC-58 cells was increased by treatment with anticancer agents including doxorubicin and decitabine (data not shown). These data suggest that CD13 is one of candidates that can be used as a marker to enrich a cell population containing putative CSCs.

CD13 is an abundant myeloid differentiation antigen that is also expressed on non-hematopoietic cells. Several reports have shown a relationship between CD13 and cancer [9,33]. CD13 is thought to play an important role in protein digestion and also to decompose type-4 collagen and confer an extra-cellular matrix metastasizing ability on tumor cells. Moreover, CD13 is expressed by endothelial cells in the process of neovascularization and can contribute to enhancement of tumor metastasis and invasion, indicating that CD13 expression is a candidate for contribution to the poor prognosis of cancers [34,35]. Indeed, some inhibitors with CD13-inhibiting properties have entered anti-cancer clinical trials. Thus, Bestatin (ubenimex), which was originally described as an immunomodulant, inhibits aminopeptidase activity, and several clinical trials targeted towards hematological malignancies have been conducted with this agent. A survival benefit of ubenimex has been demonstrated in acute myeloid leukemia and lymphomas [36]. Additionally, various clinical studies have been conducted with ubenimex in non-hematopoietic cancers such as lung, bladder, head and neck, esophagus and skin [37]. As shown in Fig. 3C, bestatin exhibited antiproliferative effect of HSC-58 cells in vitro and CD13^{high} cell population of HSC-58 was more sensitive to bestatin than CD13^{low} HSC-58 cells. Furthermore, curcumin is a natural phenolic product isolated from turmeric that has been reported to inhibit CD13 [38]. The inhibition of CD13 by curcumin is greater than that by bestatin, consequently curcumin is currently being investigated for its effect on patients with breast, prostate and pancreatic cancer [39–42]. These results indicate that CD13 is associated with cell growth, invasion and metastasis of various cancers and suggest that CD13 is a suitable target for anticancer therapy.

Antibody targeting of cell surface molecules that are overexpressed in cancer cells compared to normal cells is effective for cancer therapy. Currently, more than 30

therapeutic monoclonal antibodies have been approved for clinical applications against cancers [43,44]. Novel strategies of conjugation of drugs or radioactive agents to such antibodies are particularly promising therapeutic approaches. Several drug-conjugated antibodies specific for CD33, CD20, HER2 and CD22 are currently being tested in clinical trials [45,46]. However, not all antibodies that bind to cell surface molecules are suitable for use as therapeutic drug conjugates. This is because, in order for delivery of the drug/cytotoxic agent that is bound to the antibody into the cells, it is essential that the drug-conjugated antibody be internalized together with the cytotoxic agent into the cancer cells. There has so far been no investigation regarding the usefulness of CD13 for drug-conjugated antibody therapy. In order to investigate this possibility, we analyzed the cytotoxicity of immunotoxin (saporin) conjugated anti-CD13 antibody treatment of HSC-58 cells in vitro. As shown in Fig. 4, the efficiency of in vitro toxin delivery into HSC-58 cells was higher for anti-CD13 antibody in combination with saporin-conjugated anti-mouse IgG than for the control IgG, and its efficacy increased in a saporin-conjugated antibody dose-dependent manner. As compared with anti-CD13/APN inhibitors such as bestatin and curcumin, the advantage of anti-CD13 antibody conjugate therapy is more specific and effective against scirrhous gastric cancer HSC-58.

The efficacy of the anti-CD13 antibody for toxin delivery is in contrast to that of the anti-EpCAM antibody. Thus, although EpCAM is highly expressed in HSC-58 cells, improvement of toxin delivery compared to control IgG using the anti-EpCAM antibody is quite low. Low efficiency of EpCAM-targeted immunotoxin might be due to the lack of internalization property of anti-EpCAM antibody. The higher efficacy of the anti-CD13 antibody may be due to the fact that CD13 has been characterized as a cellular receptor for coronavirus and cytomegalovirus infection, indicating that CD13 can efficiently deliver the virus protein into the cells [47]. The antibody to CD13 can function as an artificial ligand and that can mimic a natural viral CD13 ligand against both CD13^{high} and CD13^{low} cell population of HSC-58 cells, indicating that anti-CD13 antibody-mediated intracellular delivery of immunotoxin may be therapeutically effective against scirrhous gastric cancer. We are also planning to investigate the therapeutic effect of anti-CD13 immunoliposome containing curcumin against scirrhous gastric cancer xenograft.

In conclusion, the data of this study suggest the possibility that a certain cell population of scirrhous gastric cancers with drug-resistance express CD13 that it may also function as a target for cancer therapy. However, further investigation is needed to clarify the contribution of CD13 to CSC function and to corroborate the role of CD13 in drug-resistance or tumor activity by analysis of clinical specimens of scirrhous gastric cancers.

Disclosure of interest

The authors declare that they have no competing interest.

Acknowledgments

The authors wish to thank Ritsuko Harigai and Noriyuki Ohmori for performing cell culture, flow cytometry, cell

sorting and chemosensitivity assay. This research was supported in part by grant-in-aid for Scientific Research "B" (grant no. 25293133) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- [1] Yashiro M, Hirakawa K. Cancer-stromal interactions in scirrhous gastric carcinoma. *Cancer Microenviron* 2010;3:127–35.
- [2] Kitamura K, Beppu R, Anai H, Ikejiri K, Yakabe S, Sugimachi K, et al. Clinicopathologic study of patients with Borrmann type IV gastric carcinoma. *J Surg Oncol* 1995;58:112–7.
- [3] Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med* 1997;3:730–7.
- [4] Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci U S A* 2003;100:3983–8.
- [5] Singh SK, Hawkins C, Clarke ID, Squire JA, Bayani J, Hide T, et al. Identification of human brain tumour initiating cells. *Nature* 2004;432:396–401.
- [6] Ricci-Vitiani L, Lombardi DG, Pilozzi E, Biffoni M, Todaro M, Peschle C, et al. Identification and expansion of human colon-cancer-initiating cells. *Nature* 2007;445:111–5.
- [7] Ma S, Chan KW, Hu L, Lee TK, Wo JY, Ng IO, et al. Identification and characterization of tumorigenic liver cancer stem/progenitor cells. *Gastroenterology* 2007;132:2542–56.
- [8] Li C, Heidt DG, Dalerba P, Burant CF, Zhang L, Adsay V, et al. Identification of pancreatic cancer stem cells. *Cancer Res* 2007;67:1030–7.
- [9] Haraguchi N, Ishii H, Mimori K, Tanaka F, Ohkuma M, Kim HM, et al. CD13 is a therapeutic target in human liver cancer stem cells. *J Clin Invest* 2010;120:3326–39.
- [10] Ginestier C, Hur MH, Charafe-Jauffret E, Monville F, Dutcher J, Brown M, et al. ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell Stem Cell* 2007;1:555–67.
- [11] Yanagihara K, Tanaka H, Takigahira M, Ino Y, Yamaguchi Y, Toge T, et al. Establishment of two cell lines from human gastric scirrhous carcinoma that possess the potential to metastasize spontaneously in nude mice. *Cancer Sci* 2004;95:575–82.
- [12] Motoyama T, Hojo H, Watanabe H. Comparison of seven cell lines derived from human gastric carcinomas. *Acta Pathol Jpn* 1986;36:65–83.
- [13] Rocco A, Liguori E, Pirozzi G, Tirino V, Compare D, Franco R, et al. CD133 and CD44 cell surface markers do not identify cancer stem cells in primary human gastric tumors. *J Cell Physiol* 2012;227:2686–93.
- [14] Buhning HJ, Kuci S, Conze T, Rathke G, Bartolovic K, Grunebach F, et al. CD133 identifies a broad spectrum of normal and malignant stem/progenitor cell subsets of hematopoietic and nonhematopoietic origin. *Stem Cells* 2004;22:334–43.
- [15] Keshet GI, Goldstein I, Itzhaki O, Cesarkas K, Shenhav L, Yakirevitch A, et al. MDR1 expression identifies human melanoma stem cells. *Biochem Biophys Res Commun* 2008;368:930–6.
- [16] Zhou S, Schuetz JD, Bunting KD, Colapietro AM, Sampath J, Morris JJ, et al. The ABC transporter Bcrp1/ABCG2 is expressed in a wide variety of stem cells and is a molecular determinant of the side-population phenotype. *Nat Med* 2001;7:1028–34.
- [17] Britton KM, Eyre R, Harvey IJ, Stemke-Hale K, Browell D, Lennard TW, et al. Breast cancer, side population cells and ABCG2 expression. *Cancer Lett* 2012;323:97–105.
- [18] Leccia F, Del Vecchio L, Mariotti E, Di Noto R, Morel AP, Puisieux A, et al. ABCG2, a novel antigen to sort luminal progenitors of BRCA1- breast cancer cells. *Mol Cancer* 2014;13:213.
- [19] Zhang G, Wang Z, Luo W, Jiao H, Wu J, Jiang C. Expression of potential cancer stem cell marker ABCG2 is associated with malignant behaviors of hepatocellular carcinoma. *Gastroenterol Res Pract* 2013;2013:782581.
- [20] Abbott BL. ABCG2 (BCRP) expression in normal and malignant hematopoietic cells. *Hemat Oncol* 2003;21:115–30.
- [21] Chen W, Zhang X, Chu C, Cheung WL, Ng L, Lam S, et al. Identification of CD44⁺ cancer stem cells in human gastric cancer. *Hepatogastroenterology* 2013;60:949–54.
- [22] Nishikawa S, Konno M, Hamabe A, Hasegawa S, Kano Y, Ohta K, et al. Aldehyde dehydrogenase high gastric cancer stem cells are resistant to chemotherapy. *Int J Oncol* 2013;42:1437–42.
- [23] Liu Q, Li JG, Zheng XY, Jin F, Dong HT. Expression of CD133, PAX2, ESA, and GPR30 in invasive ductal breast carcinomas. *Chin Med J* 2009;122:2763–9.
- [24] Eramo A, Lotti F, Sette G, Pilozzi E, Biffoni M, Di Virgilio A, et al. Identification and expansion of the tumorigenic lung cancer stem cell population. *Cell Death Differ* 2008;15:504–14.
- [25] Curley MD, Therrien VA, Cummings CL, Sergeant PA, Koulouris CR, Friel AM, et al. CD133 expression defines a tumor initiating cell population in primary human ovarian cancer. *Stem Cells* 2009;27:2875–83.
- [26] Hermann PC, Huber SL, Herrler T, Aicher A, Ellwart JW, Guba M, et al. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem Cell* 2007;1:313–23.
- [27] Richardson GD, Robson CN, Lang SH, Neal DE, Maitland NJ, Collins AT. CD133, a novel marker for human prostatic epithelial stem cells. *J Cell Sci* 2004;117:3539–45.
- [28] Nakai E, Park K, Yawata T, Chihara T, Kumazawa A, Nakabayashi H, et al. Enhanced MDR1 expression and chemoresistance of cancer stem cells derived from glioblastoma. *Cancer Invest* 2009;27:901–8.
- [29] Shigeta J, Katayama K, Mitsunashi J, Noguchi K, Sugimoto Y. BCRP/ABCG2 confers anticancer drug resistance without covalent dimerization. *Cancer Sci* 2010;101:1813–21.
- [30] Chute JP, Muramoto GG, Whitesides J, Colvin M, Safi R, Chao NJ, et al. Inhibition of aldehyde dehydrogenase and retinoid signaling induces the expansion of human hematopoietic stem cells. *Proc Natl Acad Sci U S A* 2006;103:11707–12.
- [31] Hong SP, Wen J, Bang S, Park S, Song SY. CD44-positive cells are responsible for gemcitabine resistance in pancreatic cancer cells. *Int J Cancer* 2009;125:2323–31.
- [32] Prochazka L, Tesarik R, Turanek J. Regulation of alternative splicing of CD44 in cancer. *Cell Signal* 2014;26:2234–9.
- [33] Wickstrom M, Larsson R, Nygren P, Gullbo J. Aminopeptidase N (CD13) as a target for cancer chemotherapy. *Cancer Sci* 2011;102:501–8.
- [34] Saiki I, Fujii H, Yoneda J, Abe F, Nakajima M, Tsuruo T, et al. Role of aminopeptidase N (CD13) in tumor-cell invasion and extracellular matrix degradation. *Int J Cancer* 1993;54:137–43.
- [35] Fujii H, Nakajima M, Saiki I, Yoneda J, Azuma I, Tsuruo T. Human melanoma invasion and metastasis enhancement by high expression of aminopeptidase N/CD13. *Clin Exp Metastasis* 1995;13:337–44.
- [36] Urabe A, Mutoh Y, Mizoguchi H, Takaku F, Ogawa N. Ubenimex in the treatment of acute nonlymphocytic leukemia in adults. *Ann Hematol* 1993;67:63–6.
- [37] Ota K, Uzuka Y. Clinical trials of bestatin for leukemia and solid tumors. *Biotherapy* 1992;4:205–14.
- [38] Shim JS, Kim JH, Cho HY, Yum YN, Kim SH, Park HJ, et al. Irreversible inhibition of CD13/aminopeptidase N by the antiangiogenic agent curcumin. *Chem Biol* 2003;10:695–704.
- [39] Liu D, Chen Z. The effect of curcumin on breast cancer cells. *J Breast Cancer* 2013;16:133–7.
- [40] Ryan JL, Heckler CE, Ling M, Katz A, Williams JP, Pentland AP, et al. Curcumin for radiation dermatitis: a randomized, double-blind, placebo-controlled clinical trial of thirty breast cancer patients. *Radiat Res* 2013;180:34–43.

- [41] Lin TH, Lee SO, Niu Y, Xu D, Liang L, Li L, et al. Differential androgen deprivation therapies with anti-androgens casodex/bicalutamide or MDV3100/enzalutamide versus anti-androgen receptor ASC-J9(R) lead to promotion versus suppression of prostate cancer metastasis. *J Biol Chem* 2013;288:19359–69.
- [42] Kanai M, Otsuka Y, Otsuka K, Sato M, Nishimura T, Mori Y, et al. A phase I study investigating the safety and pharmacokinetics of highly bioavailable curcumin (theracurmin) in cancer patients. *Cancer Chemother Pharmacol* 2013;71:1521–30.
- [43] Scott AM, Allison JP, Wolchok JD. Monoclonal antibodies in cancer therapy. *Cancer Immun* 2012;12:14.
- [44] Stern M, Herrmann R. Overview of monoclonal antibodies in cancer therapy: present and promise. *Crit Rev Oncol Hematol* 2005;54:11–29.
- [45] Borthakur G, Rosenblum MG, Talpaz M, Daver N, Ravandi F, Faderl S, et al. Phase 1 study of an anti-CD33 immunotoxin, humanized monoclonal antibody M195 conjugated to recombinant gelonin (HUM-195/rGEL), in patients with advanced myeloid malignancies. *Haematologica* 2013;98:217–21.
- [46] Lambert JM. Drug-conjugated antibodies for the treatment of cancer. *Br J Clin Pharmacol* 2013;76:248–62.
- [47] Kasman LM. CD13/aminopeptidase N and murine cytomegalovirus infection. *Virology* 2005;334:1–9.