

Erratum

Hispidulin prevents hypoxia-induced epithelial-mesenchymal transition in human colon carcinoma cells: Am J Cancer Res. 2015; 5(3): 1047-1061

Jing Xie^{1*}, Hui Gao^{1*}, Jianjun Peng^{2*}, Yantao Han^{1*}, Xuehong Chen¹, Qixiao Jiang¹, Chunbo Wang¹

¹Medical College, Qingdao University, Qingdao 266071, Shandong, China; ²College of Life Sciences, Chongqing Normal University, Chongqing 401331, China. *Equal contributors.

Received February 27, 2020; Accepted March 2, 2020; Epub April 1, 2020; Published April 15, 2020

Recently, we rechecked all our papers and realized that some data in **Figures 2** and **3** were wrongly presented. The data presented came from the wrong folder. Therefore, we would like to make corrections. Hope you can help me with this process. The revised **Figures 2** and **3** are presented as following:

Address correspondence to: Dr. Jianjun Peng, College of Life Sciences, Chongqing Normal University, Chongqing 401331, China. E-mail: jianjunpeng@126.com; Dr. Yantao Han, Medical College, Qingdao University, Qingdao 266071, Shandong, China. E-mail: hanyt19@126.com

Hispidulin prevents hypoxia-induced EMT in cancer cells




Figure 2. Hispidulin suppresses hypoxia-induced migration (A) and invasion (B) in HT-29 cells. HT-29 cells growing in hypoxia were treated with hispidulin (25 μM) for 24 hours. Cell migration and invasion were assessed by wound scratch and Transwell assays, respectively. The results represent mean \pm SD from three independent experiments. *P < 0.05 vs. control, ^P < 0.05 vs. hypoxia.

Hispidulin prevents hypoxia-induced EMT in cancer cells




Figure 3. Hispidulin prevents hypoxia-induced EMT. HT-29 cells growing in hypoxia were treated with hispidulin (25 μ M) for 24 hours. A. Effect of hispidulin on cell morphology. The images represent three independent experiments. B. The effect of hispidulin on the expression of E-cadherin examined by fluorescence staining, western blotting and qRT-PCR. The images and blot are representative of three independent experiments. C. The effect of hispidulin on the expressions of N-cadherin, vimentin, fibronectin and α -SMA determined by western blotting and qRT-PCR. The immunoblot represents three independent experiments. Data are presented as mean \pm SD. *P < 0.05 vs. control, ^P < 0.05 vs. hypoxia.