



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Recently, a small number of patients have presented with systemic intra-arterial thrombosis and a diagnosis of SARS-CoV-2. There is some evidence to suggest that this may occur, with a small series reporting ischemic strokes in patients in intensive care units.⁵ In our experience, patients presenting with limb-threatening ischemia and SARS-CoV-2 have had no other clear reason for development of intra-arterial thrombosis. Management has been dictated primarily by the severity of the organ dysfunction. A conservative approach with the prescription of therapeutic low-molecular-weight-heparin with subsequent conversion to a direct-acting oral anticoagulant has been an effective strategy.

Vascular surgeons have to recognize the potential for limb- and life-threatening arterial occlusion events in patients with SARS-CoV-2. Further research is required to try to elucidate the mechanisms by which this disease process activates the coagulation cascade.

Kirsten Victoria Hamilton, MBChB, MRCS Ed
Keith Kelso Hussey, MBChB, FRCS Ed, MD

Queen Elizabeth University Hospital
Glasgow, Scotland

REFERENCES

1. Dolhnikoff M, Duarte-Neto AN, de Almeida Monteiro RA, Ferraz da Silva LF, Pierre de Oliveira E, Nascimento Saldiva PH, et al. Pathological evidence of pulmonary thrombotic phenomena in severe COVID-19. *J Thromb Haemost* 2020;18:1517-9.
2. Liu PP, Blet A, Smyth D, Li H. The science underlying COVID-19: implications for the cardiovascular system. *Circulation* 2020;142:68-78.
3. Terpos E, Ntanasis-Stathopoulos I, Elalamy I, Kastritis E, Sergentanis TN, Politou M, et al. Hematological findings and complications of COVID-19. *Am J Hematol* 2020;95:834-47.
4. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost* 2020;18:844-7.
5. Klok F, Kruijff M, Meer NV, Arbous M, Gommers D, Kant K, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020;191:145-7.

<https://doi.org/10.1016/j.jvs.2020.05.019>

An increased severity of peripheral arterial disease in the COVID-19 era



The recent coronavirus 2019 (COVID-19) pandemic has significantly increased the pressure on our healthcare system around the world. The health emergency has changed the organization of health institutions and focused attention on pandemic management. This has led to important changes in the treatment of patients without COVID-19 and has resulted in the most difficult access to care with delays in diagnosis and treatment. Vascular diseases, including peripheral arterial disease,

will require rapid treatment in most cases. The severity of peripheral arterial disease is assessed using a functional classification system (Fontaine or Rutherford). The most severe grades of classifications include critical limb ischemia (CLI). CLI is defined by limb pain at rest, nonhealing wounds or ulcers, and/or gangrene in one or both legs.¹ CLI requires rapid revascularization to avoid tissue loss and amputation. The introduction of the lockdown in Italy from March 9, 2020, led to the closure of outpatient activities and prevented early observation of patients with CLI. Furthermore, the fear of contagion led to an underestimation of the symptoms and delayed access to treatment through the emergency room. Thus, patients with severe CLI with significant septic ulcers and gangrene arrived at our vascular surgery department. For most of these patients, it was not possible to save the limbs, and, therefore, they underwent amputation surgery. We observed that the number of amputations performed in our department from March 9 to April 20, 2020, was significantly greater than the number performed in the same period in 2019. Specifically, we performed 9 amputations during this period compared with 5 amputations performed in 2019 in the same period; an increase of almost 50%. We believe that the COVID-19 pandemic has led to the poor treatment of patients with other pathologic entities; thus, it is necessary to adopt more suitable measures to avoid other serious consequences on the health of citizens. Consequently, it is necessary to identify paths that will allow these patients to have rapid access to treatment with marked improvements in outcome.

Giuseppe Sena, MD
Giuseppe Gallelli, MD

Department of Vascular Surgery
"Pugliese-Ciaccio" Hospital
Catanzaro, Italy

REFERENCE

1. Gerhard-Herman MD, Gornik HL, Barratt C, Barshes NR, Corriere MA, Drachman DE, et al. 2016 AHA/ACC guideline on the management of patients with lower extremity peripheral artery disease: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. *Circulation* 2017;135:726-79.

<https://doi.org/10.1016/j.jvs.2020.04.489>

Cardiovascular examination should also include peripheral arterial evaluation for COVID-19 patients



The global impact of coronavirus disease (COVID-19) on vascular surgical services has been initially addressed by Ng et al.¹ This change in how vascular surgery