

RESEARCH PAPER



Cell wall/vacuolar inhibitor of fructosidase 1 regulates ABA response and salt tolerance in Arabidopsis

Wenting Yang^a, Siyu Chen^a, Yuxin Cheng^a, Na Zhang^a, Yanxing Ma^a, Wei Wang^a, Hainan Tian^a, Yingying Li^a, Saddam Hussain^a, and Shucai Wang^{a,b}

^aKey Laboratory of Molecular Epigenetics of MOE, Institute of Genetics and Cytology, Northeast Normal University, Changchun, China; ^bCollege of Life Sciences, Linyi University, Linyi, China

ABSTRACT

ABA regulates abiotic stress tolerance in plants via activating/repressing gene expression. However, the functions of many ABA response genes remained unknown. *C/VIFs* are proteinaceous inhibitors of the CWI and VI invertases. We report here the involvement of *C/VIF1* in regulating ABA response and salt tolerance in Arabidopsis. We found that the expression level of *C/VIF1* was increased in response to ABA treatment. By using CRISPR/Cas9 gene editing, we generated transgene-free *c/vif1* mutants. We also generated *C/VIF1* overexpression plants by expressing *C/VIF1* under the control of the 35S promoter. We examined ABA response of the 35S:*C/VIF1* transgenic plants and the *c/vif1* mutants by using seed germination and seedling greening assays, and found that the 35S:*C/VIF1* transgenic plants showed an enhanced sensitivity to ABA treatment in both assays. On the other hand, the *c/vif1* mutants showed slight enhanced tolerance to ABA only at the early stage of germination. We also found that salt tolerance was reduced in the 35S:*C/VIF1* transgenic plants in seed germination assays, but slightly increased in the *c/vif1* mutants. Taken together, our results suggest that *C/VIF1* is an ABA response gene, and *C/VIF1* is involved in the regulation of ABA response and salt tolerance in Arabidopsis.

ARTICLE HISTORY

Received 11 February 2020
Revised 5 March 2020
Accepted 6 March 2020

KEYWORDS

Cell wall/vacuolar inhibitor of fructosidase 1; abscisic acid; salt tolerance; gene editing; Arabidopsis

Introduction

Abiotic stresses including drought, cold, salinity and extreme temperatures reduce the yield of crops and other plants.^{1,2} ABA (abscisic acid) is the key plant hormone that regulates plant responses to abiotic stresses.³⁻⁶ ABA signaling through the PYR1/PYL/RCAR (Pyrabactin resistance 1/PYR1-like/Regulatory component of ABA receptor) receptors,⁷⁻⁹ the PP2Cs (A-group PROTEIN PHOSPHATASE 2Cs) phosphatases,^{10,11} the SnRKs (NONFERMENTING 1 (SNF1)-RELATED PROTEIN KINASES) protein kinases,¹² and a few ABF/AREB/ABI5-type bZIP (basic region leucine zipper) transcription factors¹³⁻¹⁵ regulate plant response to abiotic stresses via activating/repressing ABA response genes.^{3,5,6,10-12} Consistent with this, mutation in the ABA signaling genes affected plant tolerance to abiotic stresses. For example, overexpression of *PYL* genes enhanced drought tolerance in plants,^{16,17} whereas loss-of-function of *SnRK* genes or ABF/AREB/ABI5-type bZIP transcription factor genes led to reduced drought tolerance in plant.^{18,19} However, functions of a lot of ABA response genes in ABA signaling and/or plant abiotic stress responses remained unknown.

Cell wall invertase (CWI) and vacuolar invertase (VI) are acid invertases, they can irreversibly cleave sucrose, a primary form of carbon assimilates in the non-photosynthetic sink organs, into glucose and fructose.²⁰⁻²³ The glucose and fructose then serve as metabolites and nutrient sources of the plants, and as signaling molecules to modulate gene expression during plant growth and development as well as plant

responding to environmental cues.^{24,25} As a result, up- or down-regulation of *CWI* and *VI* expression levels affected plant growth and development and/or plants response to stresses. For example, overexpression of *CWI* gene delayed leaf aging and enhanced drought tolerance in tobacco and tomato,^{26,27} suppression of *VI* gene decreased cold-induced sweetening in potato,^{28,29} and ectopic expression of a tea *VI* gene enhanced cold tolerance in Arabidopsis.³⁰

Activities of CWI and VI invertases are regulated by the low-molecular-weight proteinaceous inhibitors, cell wall/vacuolar inhibitor of fructosidases (*C/VIFs*), who can physically interact with CWI and VI invertases and inhibit their activities.³¹⁻³³ Consistent with their biological functions, changes in the expression of *C/VIF* genes also affected plant growth and development and/or plants response to stresses. For example, silencing of *CIF* gene led to improved seed filling, prolonged leaf green, and enhanced cold tolerance in tomato,^{34,35} loss-of-function of *CIF* gene resulted in increased seed production and germination in Arabidopsis,³⁶ and ectopic expression of tobacco *VIF* enhanced drought tolerance in Arabidopsis.³⁷ However, the relationship between ABA and *C/VIF* inhibitors in regulating plant abiotic stress tolerance remained largely unclear.

We report here the roles of *C/VIF1* in regulating ABA response and salt tolerance in Arabidopsis. We found that the expression level of *C/VIF1* was upregulated by ABA, and transgenic plant overexpressing *C/VIF1* showed enhanced sensitivity to ABA and reduced tolerance to salt.

Materials and methods

Plant materials and growth conditions

The Columbia-0 (Col) wild-type *Arabidopsis thaliana* was used as a control for ABA sensitivity and salt tolerance assays, and for plant transformation to generate *c/vif1* mutants and 35S:*C/VIF1* transgenic plants.

For plant transformation, seeds of the Col wild type were germinated and grown in soil pots in a growth room. For total RNA isolation, ABA response and salt tolerance assays, seeds of the Col wild type, the 35S:*C/VIF1* transgenic plants and the *c/vif1* mutants were sterilized with 25% (v/v) bleach for 10 min, washed with sterilized water four times, and placed on vitamins (PlantMedia)-containing 1/2 MS (Murashige & Skoog) plates. The medium contains 1% (w/v) sucrose and was solidified with 0.6% (w/v) phytoagar (PlantMedia). The plates were kept in darkness at 4°C for 2 d and then transferred to a growth room.

The growth conditions in the growth room were set at 22°C, and a 16 h light/8 h dark cycle with photon density at $\sim 125 \mu\text{mol m}^{-2} \text{s}^{-1}$.

ABA treatment, RNA isolation and RT-PCR

To examine the expression of *C/VIF1* and *C/VIF2* in response to ABA, 7-day-old Col wild-type seedlings were treated with 50 μM ABA or solvent methanol as a control for 4 h, frozen in liquid N_2 . Total RNA was then isolated from the seedlings, and cDNA was synthesized as previously described.³⁸ RT-PCR was used to examine the expression of *C/VIF1* and *C/VIF2* in response to ABA treatment, and qRT-PCR was used to quantitative the fold changes of the expression levels. The expression of *ACT2* (*ACTIN2*) was used as a control. The primers used for PCR have been described previously.³⁶

Constructs

To generate the 35 S:*C/VIF1* construct for plant transformation, the full-length ORF sequence of *C/VIF1* was amplified by RT-PCR, double digested with *Nde1* and *Sac1* enzymes and cloned in frame with an N-terminal HA tag into the *pUC19* vector. The *pUC19*-35S:*C/VIF1* construct was then digested with *Pst1* and *Sac1* enzymes and cloned into the binary vector *pPZP211*,³⁹ to generate *pPZP*-35S:*C/VIF1* construct. The primers used to amplify *C/VIF1* were *C/VIF1-Nde1F*, 5'-CAACATATGATGAAGATGATGAAGGTGATGATG-3', and *C/VIF1-Sac1R*, 5'-CGAGCTCTTACCAACAAGTTCTTCTCTATTGAA-3'.

To generate *CRISPR/Cas9* constructs for *C/VIF1* gene editing, potential target sequences within the exons of *C/VIF1* were selected by using CRISPRscan (<http://www.crisprscan.org/?page=sequence>). Target specificity was then evaluated on Cas-OFFinder (<http://www.rgenome.net/cas-offinder/>). The specific target sequences selected for editing *C/VIF1* were 5'-GGTTCCTCTGCCGACACTTC(TGG)-3', 5'-GTCCCCAAATTCGGCGAAGA(CGG)-3' and 5'-GAAGACGGTGTGATC GACGC(CGG)-3'. We also selected one target for *C/VIF2*, i.e., 5'-AGTCCCACAGCCGACACAAA(AGG)-3'. The target sequences were inserted into the *pHEE-FT* vector,⁴⁰ by using

a method as described by Wang et al.⁴¹ The primers used to generate *pHEE-FT-C/VIF1* construct were *DT1-BsF* (*C/VIF1sgR1*), 5'-ATATATGGTCTCGATTGGTTCCTCTGCCGACACTTCGTT-3', *DT1-F0* (*C/VIF1sgR1*), 5'-TGGTTCCTCGCCGACACTTCGTTTATAGAGCTAGAAATAGC-3', *DT2-R0* (*C/VIF1sgR2*), 5'-AACGCGTCGATCACACCGTCTTCAATCTCTTAGTTCGACTCTAC-3', and *DT2-BsR* (*C/VIF1sgR2*), 5'-ATTATTGGTCTCGAAACGCGTCGATCACACCGTCTTC-3'. The primers used to generate *pHEE-FT-C/VIF1-C/VIF2* construct were *DT1-BsF*(*C/VIF1sgR3*), 5'-ATATATGGTCTCGATTGTCGCCCAAATTCGGCGAAGAGTT-3', *DT1-F0* (*C/VIF1sgR3*), 5'-TGTCCCCAAATTCGGCGAAGAGTTTATAGAGCTAGAAATAGC-3', *DT2-R0* (*C/VIF2sgR1*), 5'-AACTTTGTGTCGGCTGTGGGACCAATCTCTTAGTTCGACTCTAC-3', and *DT2-BsR* (*C/VIF2sgR1*), 5'-ATTATTGGTCTCGAAACTTTGTGTCGGCTGTGGGACC-3'. The *U626-IDF* and *U629-IDR* primers⁴² were used for colony PCR and sequencing of the *CRISPR/Cas9* constructs generated.

Plant transformation, transgenic plant selection and transgene-free mutant isolation

The Col wild-type plants with several mature flowers (~ 5 -week-old under our growth conditions) were transformed via *GV3101* mediated transformation by using floral dip method.⁴³ The plants were transformed with the *pPZP*-35S:*C/VIF1* construct to generate 35S:*C/VIF1* overexpression plants, and the *CRISPR/Cas9* constructs to generate *c/vif1* mutants.

The 35S:*C/VIF1* transgenic plants were selected by germinating the T1 seeds on 1/2 MS plates containing 50 $\mu\text{g/ml}$ Kanamycin and 100 $\mu\text{g/ml}$ Carbenicillin. The 35S:*C/VIF1* transgenic plants with a single T-DNA insertion were selected by germinating T2 seeds, whereas homozygous 35S:*C/VIF1* transgenic plants were selected by germinating T3 seeds on 1/2 MS plates containing 30 $\mu\text{g/ml}$ Kanamycin. Multiple homozygous lines were obtained and two with high expression levels of *C/VIF1* were used for the experiments.

Transgenic plants generated with the *CRISPR/Cas9* constructs were selected by germinating the T1 seeds on 1/2 MS plates containing 30 $\mu\text{g/ml}$ Hygromycin and 100 $\mu\text{g/ml}$ Carbenicillin, and transgene-free mutants were isolated as described previously.⁴⁰ In brief, early flowering T1 plants were selected and gene editing status was examined by amplifying and sequencing the genomic sequence of *C/VIF1*. T2 plants with normal flowering time were selected, and transgene-free status was confirmed by PCR amplification of *Cas9* gene fragment.

DNA isolation and PCR

To examine *C/VIF1* editing status, DNA was isolated from leaves of early flowering T1 or normal flowering T2 transgenic plants generated using the *CRISPR/Cas9* constructs, genomic sequences of *C/VIF1* were amplified by PCR and sequenced. To isolate transgene-free mutants, DNA was isolated from leaves of normal flowering T2 transgenic plants and used for PCR amplification of the *Cas9* gene fragment. The primers for amplifying *Cas9* fragment have been described previously.⁴²

In planta invertase activity analysis

Invertase activity was analyzed by NBT (nitroblue tetrazolium) staining as described previously.³⁶ Briefly, seeds of the Col wild type and the *c/vif1* mutants were fixed with fixation buffer, rinsed with water to remove soluble sugar, and stained with staining solution. After staining, the seeds were rinsed with water and photographed under a dissection microscopy equipped with a digital camera.

ABA and salt sensitivity assays

ABA inhibited seed germination and cotyledon greening assays were performed as described previously.⁴⁴ Briefly, seeds of the Col wild type, the 35S:*C/VIF1* transgenic plants and the *c/vif1* mutants were sterilized and germinated on 1/2 MS plates in the presence or absence of 0.75 μ M ABA. The plates were kept in darkness at 4°C for 2 d and then transferred to a growth room. Seeds germinated were counted every 12 h, photographs were taken and seedlings with green cotyledons were counted 8 d after the transfer. Percentage of germination and green seedlings was calculated.

For salt tolerance assays, seeds of the Col wild type, the 35S:*C/VIF1* transgenic plants and the *c/vif1* mutants were sterilized and germinated on 1/2 MS plates in the presence or absence of 200 mM NaCl. The plates were transferred to a growth room after kept for 2 d in darkness at 4°C. Seeds germinated were counted every 12 h, and the percentage of germination was calculated. All the experiments were repeated at least three times.

Results

Expression of *C/VIF1* is upregulated by ABA treatment

ABA plays an important role in regulating plant response to abiotic stresses. In order to identify novel players in ABA signaling and/or abiotic stress tolerance, we attempted to characterize the ABA response genes identified in our previously transcriptome assays, and with unknown functions or functions in ABA signaling and/or abiotic stresses remained unknown.⁴⁵ By using this strategy, we have previously

successfully identified bHLH129 transcription factor as an ABA response regulator,^{6,46} and AITRs as a novel family of transcription repressors that play a feed-forward regulating role in ABA signaling and function as negative regulators of plant response to abiotic stresses such as drought and salt.⁶

In the previous transcriptome data set, we found that the expression level of the invertase inhibitor gene *C/VIF1* was greatly increased in response to ABA treatment, with an RPKM of 22 in ABA treated seedlings compared with 2 in control samples. In another report, it has also been shown that the expression level of *C/VIF1* was slightly increased in response to ABA treatment.³⁶

Therefore, we examined the expression of *C/VIF1* in response to ABA in Arabidopsis seedlings. The Col wild-type seedlings were treated with 50 μ M ABA and RT-PCR was used to examine the expression of *C/VIF1*. We found that the expression level of *C/VIF1* was clearly increased in Arabidopsis seedlings treated with ABA (Figure 1a). Quantitative RT-PCR results showed that the expression level of *C/VIF1* in Arabidopsis seedlings increased about 10 folds in response to ABA (Figure 1b). On the other hand, the expression level of another known invertase inhibitor gene, *C/VIF2* remained largely unchanged (Figure 1).

Functions of *C/VIF1* is affected in the gene edited *c/vif1* mutants

Having shown that expression of *C/VIF1* was greatly upregulated by ABA treatment (Figure 1), we wanted to explore the functions of *C/VIF1* in ABA response and abiotic tolerance in Arabidopsis by generating and analyzing overexpression plants and loss-of-function mutants of the *C/VIF1*. Transgenic plants overexpressing *C/VIF1* were generated by expressing *C/VIF1* under the control of the 35S promoter (35S:*C/VIF1*) in the Col wild-type plants, and CRISPR/Cas9 gene editing was used to generate *c/vif1* mutants.

Since we have used the *FT* expression cassette-containing CRISPR/Cas9 construct *pHEE-FT* to generate transgene-free Arabidopsis mutants in a much fast and easier way,⁴⁰ the *pHEE-FT* vector was used to generate CRISPR/Cas9 constructs for *C/VIF1* gene editing. Two *pHEE-FT* vector constructs were generated. The *pHEE-FT-C/VIF1* construct was




Figure 1. Expression of *C/VIF1* in response to ABA treatment. (a) Expression of *C/VIF1* was induced by ABA. Seven-day-old Col seedlings were treated with 50 μ M ABA or mock-treated with solvent methanol for 4 h. Total RNA was isolated, cDNA was synthesized and used as a template for RT-PCR. The expression of *ACT2* was used as a control. (b) Fold changes of *C/VIF1* in response to ABA. Expression level of *C/VIF1* was examined by qRT-PCR. The expression of *ACT2* was used as an inner control. The fold change was calculated by comparing the expression levels of *C/VIF1* in ABA treated and mock-treated seedlings. Data represent the mean \pm SD of three replicates.




Figure 2. Generation of *pHEE-FT* CRISPR/Cas9 constructs for *C/VIF1* gene editing. (a) Diagram of the *pHEE-FT* vector with a *GmFT2a* expression cassette. (b) The *sgRNA* expression cassettes in the *pHEE-FT-C/VIF1* and *pHEE-FT-C/VIF1-C/VIF2* constructs. The target sequences of *C/VIF1* and *C/VIF2* were introduced into the *sgRNA* expression cassettes by PCR amplification, and the *sgRNA* expression cassettes were cloned into the *pHEE-FT* vector by using Golden gate cloning. (c) Target sequences in *C/VIF1* and *C/VIF2*. Numbers indicate the positions of the nucleotides in the coding sequences of *C/VIF1* or *C/VIF2*. PAM sites were indicated in the brackets.

made to target *C/VIF1* gene at two different sites, aimed to generate fragment deletion mutants. The *pHEE-FT-C/VIF1-C/VIF2* construct was made to target one target in *C/VIF1* gene, and the other in *C/VIF2*, intended to generate double mutants (Figure 2).

We obtained *c/vif1* single mutants with both constructs. The mutant generated by using *pHEE-FT-C/VIF1-C/VIF2* construct, *c/vif1-c1*, has a single nucleotide insertion occurred in *C/VIF1* (Figure 3a), led to a few amino acid substitutions and premature stop in *C/VIF1* (Figure 3b). The premature stop occurred after PKF (Figure 3b), a motif has been shown to be critical for invertase-inhibitor interaction.³² We also obtained another *c/vif1* mutant by using the *pHEE-FT-C/VIF1* construct, *c/vif1-c2*, which has 216 nucleotides fragment deletion in the coding sequence of *C/VIF1* (Figure 3a), resulted in 73 amino acids deletion spanning the PKF domain in *C/VIF1* (Figure 3b). NBT (nitroblue tetrazolium) staining of the seeds showed that the invertase activities were increased in both *c/vif1-c1* and *c/vif1-c2* mutants (Figure 3c), suggest that the function of *C/VIF1* was affected in these *c/vif1* mutants.

The 35S:*C/VIF1* transgenic plants are hypersensitive to ABA

After generated the *C/VIF1* overexpression and the *c/vif1* mutant plants, we examined their ABA response by using seed germination and cotyledon greening assays.

As shown in Figure 4a, reduced germination rate was observed for the 35S:*C/VIF1* transgenic plant seeds at nearly all the time points examined, indicating that the 35S:*C/VIF1*

transgenic plants are more sensitive to ABA. On the other hand, a little difference on germination rate between the seeds of Col wild type and the *c/vif1-c2* mutant plants was observed at the beginning, however, reduced germination rate was observed for the *c/vif1* mutant plant seeds at the other time points examined (Figure 4a). The results suggest that disruption of the functions of *C/VIF1* affected plant response to ABA.

In the cotyledon greening assays, the 35S:*C/VIF1* plants also showed enhanced sensitivity to ABA treatment, but a little, if any difference was observed for the *c/vif1* mutants when compared with the Col wild-type plants (Figure 4b). Quantitative analysis showed that the green cotyledon rate of the 35S:*C/VIF1* plants was only about half of the Col wild-type plants, whereas a slight decreased green cotyledon rate was observed for the *c/vif1* mutants (Figure 4c).

Salt tolerance is decreased in the 35S:*C/VIF1* transgenic plants

ABA is a key hormone that affects plant response to abiotic stresses, having shown that *C/VIF1* regulates ABA response in Arabidopsis, we further examined if *C/VIF1* may also regulate plant response to abiotic stresses. We found that reduced germination rate was observed for the 35S:*C/VIF1* transgenic plant seeds at nearly all the time points examined, suggest that salt tolerance is reduced in the 35S:*C/VIF1* transgenic plants in the seed germination assays, whereas germination rate was slightly increased in the *c/vif1-c2* mutant seeds, but the *c/vif1-c1* mutants showed a near wild-type response to salt treatments (Figure 5).




Figure 3. Isolation of transgene-free *c/vif1* mutants. (a) Editing status of *C/VIF1* in the *c/vif1* mutants. The *c/vif1-c1* and *c/vif1-c2* mutants were obtained by transforming Col wild-type plants with *pHEE-FT-C/VIF1-C/VIF2* and *pHEE-FT-C/VIF1* constructs, respectively. Editing status in T1 generation was examined, and transgene-free homozygous mutants were isolated in T2 generation. Star indicates the single T nucleotide insertion in the *c/vif1-c1* mutant. Arrowhead indicates the 216 bp fragment deletion in the *c/vif1-c2* mutant. Underlines indicate the PAM sites. (b) Alignment of *C/VIF1* amino acid sequences in the Col wild type and the *c/vif1* mutants. ORF of *C/VIF1* in the *c/vif1* mutants was identified by using ORFfinder (<https://www.ncbi.nlm.nih.gov/orffinder/>) and predicted amino acid sequences were used for alignment with the amino acid sequence of *C/VIF1*. Underline indicates the PKF motif. (c) NBT staining of the seeds. Seeds of the Col wild type and the *c/vif1* mutants were fixed with fixation buffer for 1 h at 4°C, rinsed overnight with water to remove soluble sugar, and stained with staining buffer in a water bath at 30°C. Stained seeds were rinsed with water and photographed under a dissection microscopy equipped with a digital camera.

Discussion

C/VIFs are low-molecular-weight proteinaceous inhibitors that can inhibit the activities of CWI and VI,³¹ both are acid invertases that can irreversibly cleave sucrose, into glucose and fructose.²⁰⁻²³ Consistent with their roles in sucrose metabolism, accumulated evidence suggests that changes in *C/VIF* gene expression affect plant growth and development, as well as plants response to stresses including biotic stresses and abiotic stresses in several different plants such as potato,⁴⁷⁻⁵⁰ tomato,^{34,35,51} poplar,³³ and Arabidopsis.^{36,37} Evidences provided in this study suggest that *C/VIF1* plays a role in the regulation of ABA response and salt tolerance in Arabidopsis. First, the expression of *C/VIF1* was increased in response to ABA treatment (Figure 1). Second, the 35S:*C/VIF1* transgenic plants showed enhanced sensitivity to ABA in both seed germination and seedling greening assays (Figure 4). Third,

the 35S:*C/VIF1* transgenic plants showed a reduced salt tolerance in seed germination assay (Figure 5).

In addition to serve as metabolites and nutrient sources of the plants, glucose and fructose also serve as signaling molecules to modulate gene expression during plant growth and development, and plant response to environmental stimuli,^{24,25} whereas CWI and VI are able to cleave sucrose into glucose and fructose.²⁰⁻²³ Therefore it is reasonable to speculate that *C/VIF1*'s functions in regulating salt response is, at least partially, due to its ability in inhibiting CWI and VI activities, therefore affected the cleaving of sucrose into glucose and fructose. On the other hand, considering that ABA sensitivity in the 35S:*C/VIF1* transgenic plants was affected (Figure 4), it will be of great interest to examine if and how ABA signaling may also play a role in *C/VIF1* mediated salt tolerance in Arabidopsis. Considering that salinity affects ion content in plants, it will be also of




Figure 4. Effects of ABA on seed germination and cotyledon greening of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants. (a) Effects of ABA on seed germination of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants. Sterilized seeds of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants were germinated on 1/2 MS plates in the presence or absence of 0.75 μ M ABA. The plates were transferred to a growth room after kept in darkness for 2 d at 4°C. Seeds germinated were counted every 12 h and the percentage of germination was then calculated. All the seeds on the control plates were generated 24 h after the transfer. Data represent the mean \pm SD of three replicates. (b) Effects of ABA on cotyledon greening of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants. Sterilized seeds of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants were germinated on 1/2 MS plates in the presence or absence of 0.75 μ M ABA. The plates were transferred to a growth room after kept in darkness for 2 d at 4°C, and pictures were taken 8 d after the transfer. (c) Percentage of green seedlings of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants in response to ABA treatment. Seedlings with green cotyledons were counted 8 d after the transfer and used to calculate the percentage of green cotyledons. Data represent the mean \pm SD of three replicates.

interest to examine if there is any difference in ion contents in the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants in response to salt treatment.

We noted that slightly reduced sensitivity to ABA was observed only in the *c/vif1-c2*, but not the *c/vif1-c1* mutants (Figure 4), similar, slightly enhanced salt tolerance was also observed only in the *c/vif1-c2* mutants (Figure 5). It has been reported that CIF interacts with CWI via its PKF motif, suggesting that C/VIFs inhibit invertase activities via direct physical interactions.^{32,52} Considering that the single nucleotide insertion in the *c/vif1-c1* mutant resulted in a few amino acid substitutions and premature stop for C/VIF1 after the PKF motif, whereas the 216 nucleotides fragment deletion in the *c/vif1-c2* mutant resulted in 73 amino acids deletion of the PKF domain in C/VIF1 (Figure 3), it is very likely that the *c/vif1-c2* is a loss-of-function mutant, but the *c/vif1-c1* is not. However, because the invertase activities were also increased in the *c/vif1-c1* mutant (Figure 3), it is possible that C/VIF1 in

the *c/vif1-c1* mutant may still able to interact with invertases, but its activities in inhibiting invertases were reduced. Even though additional evidence may be required to support that tolerance to ABA and salt was increased in loss-of-function mutants of C/VIF1, our evidence with the *35S:C/VIF1* transgenic plants indeed indicate that C/VIF1 plays a negative role in ABA response and salt tolerance.

It should be noted that C/VIF may have opposite effects in regulating abiotic stress responses in different plants. For example, silencing of *CIF* enhanced cold tolerance in tomato,³⁵ but ectopic expression of *VIF* enhanced drought tolerance in Arabidopsis.³⁷ However, silencing of C/VIF can improve agronomic traits in all plants studied so far. For example, silencing of *CIF* improved seed filling in tomato,³⁴ loss-of-function of *CIF1* increased seed production in Arabidopsis,³⁶ and suppression of C/VIF genes by *RNAi* improved seed weight in soybean.⁵³ Considering that CRISPR/Cas9 gene editing is able to generate transgene-free




Figure 5. Effects of NaCl on seed germination of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants. Sterilized seeds of the Col wild type, the *35S:C/VIF1* transgenic plants and the *c/vif1* mutants were germinated on 1/2 MS plates in the presence or absence of 200 mM NaCl. The plates were transferred to a growth room after kept in darkness for 2 d at 4°C. Seeds germinated were counted every 12 h and the percentage of germination was then calculated. All the seeds on the control plates generated 24 h after the transfer. Data represent the mean \pm SD of three replicates.

mutants and has been used to improve important agronomic traits in crops,⁵⁴⁻⁵⁶ editing of *C/VIF* genes together with other agronomic trait controlling genes by CRISPR/Cas9 may enable to integrate desired agronomic traits in plants.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

This work was supported by the National Key R&D Program of China (2016YFD0101900) and a startup funding from Linyi University (LYDX2019BS039). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

SW conceived the study. SW, WY designed the experiments. WY, SC, YC, NZ, YM, WW, HT, YL and SH performed the experiments. WY and SW analyzed the data. SW drafted the manuscript. All the authors participated in the revision of the manuscript.

ORCID

Shucai Wang  <http://orcid.org/0000-0001-7619-2385>

References

- Wang W, Vinocur B, Altman A. Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta*. 2003;218:1-14. doi:10.1007/s00425-003-1105-5.
- Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, Yamaguchi-Shinozaki K, Shinozaki K. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Curr Opin Plant Biol*. 2006;9:436-442. doi:10.1016/j.pbi.2006.05.014.
- Umezawa T, Nakashima K, Miyakawa T, Kuromori T, Tanokura M, Shinozaki K, Yamaguchi-Shinozaki K. Molecular basis of the core regulatory network in ABA responses: sensing, signaling and transport. *Plant Cell Physiol*. 2010;51:1821-1839. doi:10.1093/pcp/pcq156.
- Yoshida T, Mogami J, Yamaguchi-Shinozaki K. ABA-dependent and ABA-independent signaling in response to osmotic stress in plants. *Curr Opin Plant Biol*. 2014;21:133-139. doi:10.1016/j.pbi.2014.07.009.
- Song L, Huang SC, Wise A, Castanon R, Nery JR, Chen H, Watanabe M, Thomas J, Bar-Joseph Z, Ecker JR. A transcription factor hierarchy defines an environmental stress response network. *Science*. 2016;354:aag1550. doi:10.1126/science.aag1550.
- Tian H, Chen S, Yang W, Wang T, Zheng K, Wang Y, Cheng Y, Zhang N, Liu S, Li D, et al. A novel family of transcription factors conserved in angiosperms is required for ABA signalling. *Plant Cell Environ*. 2017;40:2958-2971.
- Ma Y, Szostkiewicz I, Korte A, Moes D, Yang Y, Christmann A, Grill E. Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science*. 2009;324:1064-1068. doi:10.1126/science.1172408.
- Park SY, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A, Chow TF, et al. Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science*. 2009;324:1068-1071. doi:10.1126/science.1173041.
- Santiago J, Rodrigues A, Saez A, Rubio S, Antoni R, Dupeux F, Park SY, Marquez JA, Cutler SR, Rodriguez PL. Modulation of drought resistance by the abscisic acid receptor PYL5 through inhibition of clade A PP2Cs. *Plant J*. 2009;60:575-588. doi:10.1111/j.1365-3113.2009.03981.x.
- Rodriguez PL, Leube MP, Grill E. Molecular cloning in *Arabidopsis thaliana* of a new protein phosphatase 2C (PP2C) with homology to ABI1 and ABI2. *Plant Mol Biol*. 1998;38:879-883. doi:10.1023/A:1006012218704.
- Gosti F, Beaudoin N, Serizet C, Webb AA, Vartanian N, Giraudat J. ABI1 protein phosphatase 2C is a negative regulator of abscisic acid signaling. *Plant Cell*. 1999;11:1897-1910. doi:10.1105/tpc.11.10.1897.
- Fujii H, Verslues PE, Zhu JK. Identification of two protein kinases required for abscisic acid regulation of seed germination, root growth, and gene expression in *Arabidopsis*. *Plant Cell*. 2007;19:485-494. doi:10.1105/tpc.106.048538.

13. Stone SL, Williams LA, Farmer LM, Vierstra RD, Callis J. KEEP ON GOING, a RING E3 ligase essential for Arabidopsis growth and development, is involved in abscisic acid signaling. *Plant Cell*. 2006;18:3415–3428. doi:10.1105/tpc.106.046532.
14. Liu H, Stone SL. Abscisic acid increases Arabidopsis ABI5 transcription factor levels by promoting KEG E3 ligase self-ubiquitination and proteasomal degradation. *Plant Cell*. 2010;22:2630–2641. doi:10.1105/tpc.110.076075.
15. Chen YT, Liu HX, Stone S, Callis J. ABA and the ubiquitin E3 ligase KEEP ON GOING affect proteolysis of the Arabidopsis thaliana transcription factors ABF1 and ABF3. *Plant J*. 2013;75:965–976. doi:10.1111/tpj.12259.
16. Park SY, Peterson FC, Mosquana A, Yao J, Volkman BF, Cutler SR. Agrochemical control of plant water use using engineered abscisic acid receptors. *Nature*. 2015;520:545–548. doi:10.1038/nature14123.
17. Zhao Y, Chan Z, Gao J, Xing L, Cao M, Yu C, Hu Y, You J, Shi H, Zhu Y, et al. ABA receptor PYL9 promotes drought resistance and leaf senescence. *Proc Natl Acad Sci USA*. 2016;113:1949–1954. doi:10.1073/pnas.1522840113.
18. Fujita Y, Nakashima K, Yoshida T, Katagiri T, Kidokoro S, Kanamori N, Umezawa T, Fujita M, Maruyama K, Ishiyama K, et al. Three SnRK2 protein kinases are the main positive regulators of abscisic acid signaling in response to water stress in Arabidopsis. *Plant Cell Physiol*. 2009;50:2123–2132. doi:10.1093/pcp/pcp147.
19. Yoshida T, Fujita Y, Maruyama K, Mogami J, Todaka D, Shinozaki K, Yamaguchi-Shinozaki K. Four Arabidopsis AREB/ABF transcription factors function predominantly in gene expression downstream of SnRK2 kinases in abscisic acid signaling in response to osmotic stress. *Plant Cell Environ*. 2015;38:35–49. doi:10.1111/pce.12351.
20. Koch K. Sucrose metabolism: regulatory mechanisms and pivotal roles in sugar sensing and plant development. *Curr Opin Plant Biol*. 2004;7:235–246. doi:10.1016/j.pbi.2004.03.014.
21. Sturm A. Invertases. Primary structures, functions, and roles in plant development and sucrose partitioning. *Plant Physiol*. 2002;121:1–8. doi:10.1104/pp.121.1.1.
22. Van den Ende W, Lammens W, Van Laere A, Schroeven L, Le Roy K. Donor and acceptor substrate selectivity among plant glycoside hydrolase family 32 enzymes. *FEBS J*. 2009;276:5788–5798. doi:10.1111/j.1742-4658.2009.07316.x.
23. Wan H, Wu L, Yang Y, Zhou G, Ruan YL. Evolution of sucrose metabolism: the dichotomy of invertases and beyond. *Trends Plant Sci*. 2018;23:163–177. doi:10.1016/j.tplants.2017.11.001.
24. Rolland F, Baena-Gonzalez E, Sheen J. Sugar sensing and signaling in plants: conserved and novel mechanisms. *Annu Rev Plant Biol*. 2006;57:675–709. doi:10.1146/annurev-arplant.57.032905.105441.
25. Ruan Y-L. Sucrose metabolism: gateway to diverse carbon use and sugar signaling. *Annu Rev Plant Biol*. 2014;65:33–67. doi:10.1146/annurev-arplant-050213-040251.
26. Balibrea Lara ME, Gonzalez Garcia M-C, Fatima T, Ehneß R, Lee TK, Proels R, Tanner W, Roitsch T. Extracellular invertase is an essential component of cytokinin-mediated delay of senescence. *Plant Cell*. 2004;16:1276–1287. doi:10.1105/tpc.018929.
27. Albacete A, Cantero-Navarro E, Großkinsky DK, Arias CL, Balibrea ME, Bru R, Fragner L, Ghanem ME, González Mde L, Hernández JA, et al. Ectopic overexpression of the cell wall invertase gene *CINI* leads to dehydration avoidance in tomato. *J Exp Bot*. 2015;66:863–878. doi:10.1093/jxb/eru448.
28. Bhaskar PB, Wu L, Busse JS, Whitty BR, Hamernik AJ, Jansky SH, Buell CR, Bethke PC, Jiang J. Suppression of the vacuolar invertase gene prevents cold-induced sweetening in potato. *Plant Physiol*. 2010;154:939–948. doi:10.1104/pp.110.162545.
29. Zhu X, Gong H, He Q, Zeng Z, Busse JS, Jin W, Bethke PC, Jiang J. Silencing of vacuolar invertase and asparagine synthetase genes and its impact on acrylamide formation of fried potato products. *Plant Biotechnol J*. 2016;14:709–718. doi:10.1111/pbi.12421.
30. Qian W, Xiao B, Wang L, Hao X, Yue C, Cao H, Wang Y, Li N, Yu Y, Zeng J, et al. CsINV5, a tea vacuolar invertase gene enhances cold tolerance in transgenic Arabidopsis. *BMC Plant Biol*. 2018;18:228. doi:10.1186/s12870-018-1456-5.
31. Rausch T, Greiner S. Plant protein inhibitors of invertases. *Biochim Biophys Acta-Proteins Proteomics*. 2004;1696:253–261. doi:10.1016/j.bbapap.2003.09.017.
32. Hothorn M, Van den Ende W, Lammens W, Rybin V, Scheffzek K. Structural insights into the pH-controlled targeting of plant cell-wall invertase by a specific inhibitor protein. *Proc Natl Acad Sci*. 2010;107:17427–17432. doi:10.1073/pnas.1004481107.
33. Su T, Han M, Min J, Zhou H, Zhang Q, Zhao J, Fang Y. Functional characterization of invertase inhibitors PtC/VIF1 and 2 revealed their involvements in the defense response to fungal pathogen in *Populus Trichocarpa*. *Front Plant Sci*. 2020;10:1654. doi:10.3389/fpls.2019.01654.
34. Jin Y, Ni D-A, Ruan Y-L. Posttranslational elevation of cell wall invertase activity by silencing its inhibitor in tomato delays leaf senescence and increases seed weight and fruit hexose level. *Plant Cell*. 2009;21:2072–2089. doi:10.1105/tpc.108.063719.
35. Xu X, Hu Q, Yang W, Jin Y. The roles of cell wall invertase inhibitor in regulating chilling tolerance in tomato. *BMC Plant Biol*. 2017;17:195. doi:10.1186/s12870-017-1145-9.
36. Su T, Wolf S, Han M, Zhao H, Wei H, Greiner S, Rausch T. Reassessment of an Arabidopsis cell wall invertase inhibitor AtCIF1 reveals its role in seed germination and early seedling growth. *Plant Mol Biol*. 2016;90:137–155. doi:10.1007/s11103-015-0402-2.
37. Chen SF, Liang K, Yin DM, Ni DA, Zhang ZG, Ruan YL. Ectopic expression of a tobacco vacuolar invertase inhibitor in guard cells confers drought tolerance in Arabidopsis. *J Enzyme Inhib Med Chem*. 2016;31:1381–1385. doi:10.3109/14756366.2016.1142981.
38. Wang X, Wang X, Hu Q, Dai X, Tian H, Zheng K, Wang X, Mao T, Chen JG, Wang S. Characterization of an activation-tagged mutant uncovers a role of GLABRA2 in anthocyanin biosynthesis in Arabidopsis. *Plant J*. 2015;83:300–311. doi:10.1111/tpj.12887.
39. Hajdukiewicz P, Svab Z, Maliga P. The small, versatile pPZP family of Agrobacterium binary vectors for plant transformation. *Plant Mol Biol*. 1994;25:989–994. doi:10.1007/BF00014672.
40. Cheng Y, Zhang N, Hussain S, Ahmed S, Yang W, Wang S. Integration of a FT expression cassette into CRISPR/Cas9 construct enables fast generation and easy identification of transgene-free mutants in Arabidopsis. *PLoS One*. 2019;14:e0218583. doi:10.1371/journal.pone.0218583.
41. Wang ZP, Xing HL, Dong L, Zhang HY, Han CY, Wang XC, Chen QJ. Egg cell-specific promoter-controlled CRISPR/Cas9 efficiently generates homozygous mutants for multiple target genes in Arabidopsis in a single generation. *Genome Biol*. 2015;16:144. doi:10.1186/s13059-015-0715-0.
42. Chen S, Zhang N, Zhang Q, Zhou G, Tian H, Hussain S, Ahmed S, Wang T, Wang S. Genome editing to integrate seed size and abiotic stress tolerance traits in Arabidopsis reveals a role for DPA4 and SOD7 in the regulation of inflorescence architecture. *Int J Mol Sci*. 2019;20:2695. doi:10.3390/ijms20112695.
43. Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J*. 1998;16:735–743. doi:10.1046/j.1365-313x.1998.00343.x.
44. Guo J, Wang J, Xi L, Huang WD, Liang J, Chen JG. RACK1 is a negative regulator of ABA responses in Arabidopsis. *J Exp Bot*. 2009;60:3819–3833. doi:10.1093/jxb/erp221.
45. Guo J, Wang S, Valerius O, Hall H, Zeng Q, Li JF, Weston DJ, Ellis BE, Chen JG. Involvement of Arabidopsis RACK1 in protein translation and its regulation by abscisic acid. *Plant Physiol*. 2011;155:370–383. doi:10.1104/pp.110.160663.

46. Tian H, Guo H, Dai X, Cheng Y, Zheng K, Wang X, Wang S. An ABA down-regulated bHLH transcription repressor gene, bHLH129 regulates root elongation and ABA response when overexpressed in Arabidopsis. *Sci Rep.* 2015;5:17587. doi:10.1038/srep17587.
47. Greiner S, Rausch T, Sonnewald U, Herbers K. Ectopic expression of a tobacco invertase inhibitor homolog prevents cold-induced sweetening of potato tubers. *Nat Biotechnol.* 1999;17:708–711. doi:10.1038/10924.
48. Brummell DA, Chen RKY, Harris JC, Zhang H, Hamiaux C, Kralicek AV, McKenzie MJ. Induction of vacuolar invertase inhibitor mRNA in potato tubers contributes to cold-induced sweetening resistance and includes spliced hybrid mRNA variants. *J Exp Bot.* 2011;62:3519–3534. doi:10.1093/jxb/err043.
49. Liu X, Lin Y, Liu J, Song B, Ou Y, Zhang H, Li M, Xie C. StInvInh2 as an inhibitor of StvacINV1 regulates the cold-induced sweetening of potato tubers by specifically capping vacuolar invertase activity. *Plant Biotechnol J.* 2013;11:640–647. doi:10.1111/pbi.12054.
50. Mckenzie MJ, Chen RKY, Harris JC, Ashworth MJ, Brummell DA. Post-translational regulation of acid invertase activity by vacuolar invertase inhibitor affects resistance to cold-induced sweetening of potato tubers. *Plant Cell Environ.* 2013;36:176–185. doi:10.1111/j.1365-3040.2012.02565.x.
51. Qin G, Zhu Z, Wang W, Cai J, Chen Y, Li L, Tian S. A tomato vacuolar invertase inhibitor mediates sucrose metabolism and influences fruit ripening. *Plant Physiol.* 2016;172:1596–1611. doi:10.1104/pp.16.01269.
52. Hothorn M, Wolf S, Aloy P, Greiner S, Scheffzek K. Structural insights into the target specificity of plant invertase and pectin methylesterase inhibitory proteins. *Plant Cell.* 2004;16:3437–3447. doi:10.1105/tpc.104.025684.
53. Tang X, Su T, Han M, Wei L, Wang W, Yu Z, Xue Y, Wei H, Du Y, Greiner S, et al. Suppression of extracellular invertase inhibitor gene expression improves seed weight in soybean (*Glycine max*). *J Exp Bot.* 2017;68:469–482. doi:10.1093/jxb/erw425.
54. Ma X, Zhang Q, Zhu Q, Liu W, Chen Y, Qiu R, Wang B, Yang Z, Li H, Lin Y, et al. A robust CRISPR/Cas9 system for convenient, high-efficiency multiplex genome editing in monocot and dicot plants. *Mol Plant.* 2015;8:1274–1284. doi:10.1016/j.molp.2015.04.007.
55. Li X, Xie Y, Zhu Q, Liu YG. Targeted genome editing in genes and cis-regulatory regions improves qualitative and quantitative traits in crops. *Mol Plant.* 2017;10:1368–1370. doi:10.1016/j.molp.2017.10.009.
56. He Y, Zhu M, Wang L, Wu J, Wang Q, Wang R, Zhao Y. Programmed self-elimination of the CRISPR/Cas9 construct greatly accelerates the isolation of edited and transgene-free rice plants. *Mol Plant.* 2018;11:1210–1213. doi:10.1016/j.molp.2018.05.005.