



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Antihypertensive drugs and risk of COVID-19?

Lei Fang and colleagues¹ postulate that because severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) binds to the angiotensin-converting enzyme 2 (ACE2) receptor to facilitate host cell entry,^{2,3} disease severity and mortality of coronavirus disease 2019 (COVID-19) might be increased in patients taking angiotensin converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) during the COVID-19 pandemic, because the ACE2 receptor might be upregulated with use of ACEIs and ARBs.

Although it is true that many patients with severe COVID-19 have hypertension and diabetes, as reported by Guan and colleagues in a study of more than 1000 patients,⁴ in that study other factors were also notably unevenly distributed among groups, including age, sex, smoking status (eg, current smoker), and chronic pulmonary disease. Even though patients with diabetes and hypertension frequently use ACEIs and ARBs, many confounders must be accounted for when interpreting findings of observational studies.⁵ In another study looking at risk factors for mortality among patients with COVID-19,⁶ despite hypertension and diabetes being identified as the most frequently associated comorbidities, after multivariate regression the association between these disorders and COVID-19 mortality was no longer significant.

In theory, use of ACEIs and ARBs over time would result in an upregulation of ACE2 receptors. In children and young adults, ACE2 receptors are present at

a much higher density in lung tissue than in older individuals. Thus, the condition in people taking ACEIs and ARBs emulates that in young people. It is possible that having more ACE2 receptors provides a reserve against target-mediated destruction by SARS-CoV-2.⁷ ACE2 functions to produce the heptapeptide angiotensin (1-7), which has favourable effects on the pulmonary endothelium and might provide resilience from development of pulmonary failure in COVID-19.⁷

We caution against indiscriminate discontinuation of ACEIs and ARBs in patients who rely on these drugs for treatment of heart failure and who, additionally, might benefit from the postulated positive effects during overwhelming infection with SARS-CoV-2. Discontinuation of ACEIs or ARBs is associated with readmission to hospital and mortality among patients with heart failure.⁸ A surge of admissions to hospital for heart failure because of indiscriminate cessation of these important agents could overload already burdened health-care systems with vulnerable patients and cause diagnostic problems in view of the range of symptoms shared between acute heart failure and COVID-19, such as cough and shortness of breath.

The quick developing nature of the COVID-19 pandemic, and uncertainty around best practice, calls for caution when putting forward hypotheses that suggest a change in drugs. More evidence is needed before any clinical decisions are made.

We declare no competing interests.


*Kevin Bryan Lo, Peter A McCullough, Janani Rangaswami
lokevinb@einstein.edu

Department of Medicine, Einstein Medical Center, Philadelphia, PA 19141, USA (KBL, JR); Baylor Heart and Vascular Institute, Dallas, TX, USA (PAM); and Sidney Kimmel College of Thomas Jefferson University, Philadelphia, PA, USA

- 1 Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* 2020; published online March 11. [https://doi.org/10.1016/S2213-2600\(20\)30116-8](https://doi.org/10.1016/S2213-2600(20)30116-8).
- 2 Ferrario CM, Jessup J, Chappell MC, et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation* 2005; **111**: 2605–10.
- 3 Hoffmann M, Kleine-Weber H, Krueger N, Mueller MA, Drosten C, Pöhlmann S. The novel coronavirus 2019-nCoV uses the SARS-coronavirus receptor ACE2 and the cellular protease TMPRSS2 for entry into target cells. Jan 31, 2020. <https://www.biorxiv.org/content/10.1101/2020.01.31.929042v1> (accessed March 23, 2020).
- 4 Guan W, Ni Z, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020; published online Feb 28. DOI:10.1056/NEJMoa2002032.
- 5 Jepsen P, Johnsen SP, Gillman MW, Sørensen HT. Interpretation of observational studies. *Heart* 2004; **90**: 956–60.
- 6 Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; published online March 11. [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3).
- 7 Gurwitz D. Angiotensin receptor blockers as tentative SARS-CoV-2 therapeutics. *Drug Dev Res* 2020; published online March 4. DOI:10.1002/ddr.21656.
- 8 Oliveros E, Oni ET, Shahzad A, et al. Benefits and risks of continuing angiotensin-converting enzyme inhibitors, angiotensin II receptor antagonists, and mineralocorticoid receptor antagonists during hospitalizations for acute heart failure. *Cardiorenal Med* 2020; **10**: 69–84.

Related links

- The latest guidance from WHO on ibuprofen and COVID-19 (dated: 19.03.2020)
- Statement from Prof Michael Roth on how to interpret the original letter



Ramon Andrade 3D/Science Photo Library

Published Online
March 26, 2020
[https://doi.org/10.1016/S2213-2600\(20\)30156-9](https://doi.org/10.1016/S2213-2600(20)30156-9)