



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.




Figure 1 Three-dimensional point cloud surface reconstruction.

## Funding

Research grant for funding of projects in biomedicine, health care management and social work and health care services. Regional Health Care Administration. Government of Castilla y Leon, 2017. Project file 1571/A/17).

## References

1. Poncet P, Kravarusic D, Richart T, Evison R, Ronsky JL, Alassiri A, et al. Clinical impact of optical imaging with 3-D reconstruction of torso topography in common anterior chest wall anomalies. *J Pediatr Surg.* 2007;42(5):898–903.
2. Riphagen JM, van Neck JW, van Adrichem LN. 3D surface imaging in medicine: a review of working principles and implications for imaging the unsedated child. *J Craniofac Surg.* 2008;19(2):517–24.
3. Kravarusic D, Dicken BJ, Dewar R, Harder J, Poncet P, Schneider M, et al. The Calgary protocol for bracing of pectus carinatum: a preliminary report. *J Pediatr Surg.* 2006;41(5):923–6.
4. Haecker FM. The vacuum bell for conservative treatment of pectus excavatum: the Basle experience. *Pediatr Surg Int.* 2011;27(6):623–7.

Sara Fuentes<sup>a,\*</sup>, Francisco Damián<sup>b</sup>, Mario Berlioz<sup>b</sup>, José-María Pradillos<sup>a</sup>, Teresa Lorenzo<sup>b</sup>

<sup>a</sup> Servicio de Cirugía Pediátrica, Complejo Asistencial Universitario de León, León, Spain

<sup>b</sup> Servicio de Radiología, Complejo Asistencial Universitario de León, León, Spain

\*Corresponding author.

E-mail address: [sfuentesc@saludcastillayleon.es](mailto:sfuentesc@saludcastillayleon.es) (S. Fuentes).

23 January 2019 20 February 2019

<https://doi.org/10.1016/j.anpede.2019.02.006>  
2341-2879/

© 2020 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## First case of neonatal infection due to COVID-19 in Spain<sup>☆</sup>



### Primer caso de infección neonatal por COVID-19 en España

Dear Editor:

Infection by new coronavirus (2019-nCoV, later renamed SARS-CoV-2), a virus first identified in 2019 in Wuhan,

China, causes the disease known as coronavirus disease 2019 (COVID-19). This disease is currently spreading in what the World Health Organization (WHO) has now declared a global pandemic.<sup>1,2</sup>

At present, the data on COVID-19 in newborns are scarce. There is no clear evidence of vertical transmission, with few descriptions of cases of newborns of mothers infected in the third trimester of pregnancy, in whom viral detection tests have been negative. The risk of horizontal transmission through contact with an infected individual seems to be the same compared to the general population, and 3 such cases have been reported to date.<sup>3,4</sup>

Given the current lack of information, we are describing a case of neonatal infection by SARS-CoV-2 in Spain.

<sup>☆</sup> Please cite this article as: Alonso Díaz C, López Maestro M, Moral Pumarega MT, Flores Antón B, Pallás Alonso C. Primer caso de infección neonatal por COVID-19 en España. *An Pediatr (Barc).* 2020;92:237–238.

The mother was aged 41 years and underwent an urgent caesarean section due to severe preeclampsia on week 38 + 4 of gestation. The pregnancy was achieved by in vitro fertilization, and the mother had a history of hypothyroidism for which she was receiving ongoing treatment.

The newborn was a girl with low birth weight for gestational age (2500 g), a 1-min Apgar score of 7 and a 5-min Apgar score of 9 who required resuscitation with oropharyngeal suctioning. She was transferred to the neonatal unit due to immediate respiratory distress under respiratory support with continuous positive airway pressure (CPAP) without supplemental oxygen. Respiratory support was discontinued at 2 h post birth and the findings of the physical examination at 9 h post birth were normal. These problems were attributed to stress during the transition to extrauterine life, and the patient was subsequently transferred to the maternity ward to stay with her mother.

On day 3 from admission, the mother developed a low-grade fever, and obstetric complications were ruled out. Two days later the mother had peaks of fever and respiratory symptoms, and a chest X-ray revealed severe bilateral pneumonia. The mother reported a negative history of travel to coronavirus risk areas or contact with infected individuals, although her partner had exhibited fever associated with gastroenteritis since the day of the caesarean delivery. After common viral illnesses had been ruled out, a real-time reverse transcription polymerase chain reaction (RT-PCR) test for detection of coronavirus 2019 was ordered for the mother. This test detects 2 regions of viral genome. The result was positive. Her partner was tested subsequently, and the result was also positive.

Up to this point, the girl had remained asymptomatic as she stayed with her mother in the maternity ward and received mixed breastfeeding. Due to the clinical condition of the mother, which required care in a special unit, mother and child were separated and the newborn also underwent testing for COVID-19 by RT-PCR on a sample of nasopharyngeal aspirate (day 6 post birth).

Despite the negative result, the girl was kept in isolation in the maternity ward in the care of the hospital staff until a second sample was obtained 36 h later (nasopharyngeal swab) and testing turned out positive (day 8 post birth). The newborn was then transferred to the neonatal unit without performance of additional diagnostic tests, as she remained asymptomatic. She was closely monitored, and on day 9 post birth she exhibited intermittent hyperpnoea with mild intercostal retractions and 2 self-limited episodes of oxygen desaturation during deep sleep and feeding. A capillary sample was obtained for blood gas analysis, which revealed transient mild acidosis (pH, 7.27;  $p\text{CO}_2$ , 49 mmHg; bicarbonate, 22 mEq/L; base excess -4; normal lactate), a chest radiograph detected ground glass opacities mainly in the right perihilar region and the serum level of C-reactive protein was normal (0.06 mg/dL). After 24 h, the symptoms resolved, and the patient has remained asymptomatic to date (day 13 post birth), when the COVID-19 RT-PCR has been repeated and continues to be positive. Her mother is in the intensive care unit under mechanical ventilation.

We suspect this is a case of horizontal transmission, as the initial COVID-19 test was negative. The clinical presentation is consistent with the current literature,<sup>5,6</sup> which describes disease in newborns as mild and possibly associated with abnormal radiographic findings.

On the other hand, since the incubation period in newborns is unknown, we worry that testing a single sample may not suffice in cases like the one we have just presented. According to the WHO,<sup>2</sup> a negative result does not rule out infection, so if infection by SARS-CoV-2 is highly suspected in a patient, especially in cases in which only samples of the upper airway have been tested, additional tests should be performed. This is important for the purpose of preventing community transmission through the children of mothers that have tested positive for coronavirus.

## References

1. Sociedad Española de Neonatología. Recomendaciones para el manejo del recién nacido en relación con la infección por SARS-CoV-2 (08/03/2020) [consultado 12 Mar 2020]. Disponible en: [https://www.seneo.es/images/site/noticias/home/Recomendaciones\\_SENeo\\_SARS-CoV-2Version.2.pdf](https://www.seneo.es/images/site/noticias/home/Recomendaciones_SENeo_SARS-CoV-2Version.2.pdf)
2. World Health Organization. Laboratory testing for coronavirus disease 2019 (COVID-19) in suspected human cases (2 March 2020). Available from: file:///C:/Users/02545809P/Downloads/WHO-COVID-19-laboratory-2020.4-eng.pdf [accessed 12.03.20].
3. Zhu H, Wang L, Fang C, Peng S, Zhang L, Chang G, et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Transl Pediatr.* 2020;9:51–60.
4. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet.* 2020;395:809–15.
5. Wang L, Shi Y, Xiao T, Fu J, Feng X, Mu D, et al., Working Committee on Perinatal and Neonatal Management for the Prevention and Control of the 2019 Novel Coronavirus Infection. Chinese expert consensus on the perinatal and neonatal management for the prevention and control of the 2019 novel coronavirus infection (first edition). *Ann Transl Med.* 2020;8:47.
6. Lu Q, Shi Y. Coronavirus disease (COVID-19) and neonate: what neonatologist need to know. *J Med Virol.* 2020, <http://dx.doi.org/10.1002/jmv.25740>.

Clara Alonso Díaz\*, María López Maestro,  
María Teresa Moral Pumarega, Beatriz Flores Antón,  
Carmen Pallás Alonso

*Hospital Universitario 12 de Octubre, Madrid, Spain*

\* Corresponding author.

*E-mail address:* [claraalonsodiaz@gmail.com](mailto:claraalonsodiaz@gmail.com)  
(C. Alonso Díaz).

<https://doi.org/10.1016/j.anpede.2020.03.002>  
2341-2879/

© 2020 Published by Elsevier España, S.L.U. on behalf of Asociación Española de Pediatría. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).