

**Corrigendum to “Luyckx VA, Cherney DZI, Bello AK. Preventing CKD in Developed Countries.”  
*Kidney Int Rep.* 2020;5:263–277**




*Kidney Int Rep* (2020) 5, 758; <https://doi.org/10.1016/j.ekir.2020.03.021> © 2020 International Society of Nephrology. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The authors regret an error in Figure 1. A preventable/treatable risk factor for chronic kidney disease in adult women is *preeclampsia* (incorrectly written as *pregnancy* in the figure).

The authors would like to apologize for any inconvenience caused.

Prevention	Fetus/neonate	Child	Adolescent	Adult	Elderly
<b>Primordial*</b>	<ul style="list-style-type: none"> <li>• Achieve all SDGs</li> <li>• Preconception care</li> <li>• Healthy mother, father</li> <li>• Healthy pregnancy</li> </ul>	<ul style="list-style-type: none"> <li>• Achieve all SDGs</li> <li>• Educate girls</li> <li>• Public health measures</li> </ul>	<ul style="list-style-type: none"> <li>• Achieve all SDGs</li> <li>• Educate girls</li> <li>• Prevent teenage pregnancy</li> <li>• Public health measures</li> </ul>	<ul style="list-style-type: none"> <li>• Achieve all SDGs</li> <li>• Gender equality</li> <li>• Public health measures</li> </ul>	<ul style="list-style-type: none"> <li>• Achieve all SDGs</li> <li>• Healthy aging</li> <li>• Public health measures</li> </ul>
<b>Primary</b>	<ul style="list-style-type: none"> <li>• UHC/ANC</li> <li>• Lifestyle education, mother, infant</li> <li>• Preeclampsia prophylaxis</li> <li>• Identify mothers at future risk of CKD, CVD, DM</li> <li>• Avoid AKI</li> </ul>	<ul style="list-style-type: none"> <li>• Lifestyle education</li> <li>• Good nutrition</li> <li>• UHC</li> <li>• Avoid overweight/obesity</li> <li>• Avoid AKI</li> </ul>	<ul style="list-style-type: none"> <li>• Lifestyle education</li> <li>• Sexual health</li> <li>• Family planning</li> <li>• Preeclampsia prophylaxis</li> <li>• Avoid nephrotoxins</li> <li>• Avoid smoking, alcohol, drugs</li> <li>• Avoid AKI</li> <li>• UHC</li> </ul>	<ul style="list-style-type: none"> <li>• Lifestyle education</li> <li>• Preeclampsia prophylaxis</li> <li>• Avoid nephrotoxins</li> <li>• Avoid smoking, alcohol</li> <li>• Weight control</li> <li>• Avoid AKI</li> <li>• Screen high-risk groups</li> <li>• Treat HT, DM, infections</li> <li>• UHC</li> </ul>	<ul style="list-style-type: none"> <li>• Lifestyle education</li> <li>• Avoid nephrotoxins</li> <li>• Avoid smoking, alcohol</li> <li>• UHC</li> <li>• Screen high-risk groups</li> <li>• Treat HT, DM, infections</li> <li>• Weight control</li> <li>• Avoid AKI</li> </ul>
<b>Secondary</b>	<ul style="list-style-type: none"> <li>• Adjust medications in neonatal ICU</li> <li>• Manage obstruction</li> </ul>	<ul style="list-style-type: none"> <li>• Screen LBW, pre-term, SGA</li> </ul>	<ul style="list-style-type: none"> <li>• Treat HT, DM</li> <li>• Weight loss if overweight/obese</li> </ul>	<ul style="list-style-type: none"> <li>• Early diagnosis and treatment of CKD</li> <li>• Avoid AKI</li> </ul>	<ul style="list-style-type: none"> <li>• Early diagnosis and treatment of CKD</li> <li>• Avoid AKI</li> </ul>



**Figure 1.** Illustration of the spectrum of strategies for chronic kidney disease (CKD) prevention across the life course. \*Primordial prevention refers to strategies to optimize upstream factors that may lead to increased risk of CKD at an individual- or population-level. AKI, acute kidney injury; ANC, antenatal care; BMI, body mass index; BP, blood pressure; CVD, cardiovascular disease; DM, diabetes mellitus; GFR, glomerular filtration rate; HT, hypertension; ICU, intensive care unit; LBW, low birth weight; SDGs, Sustainable Development Goals<sup>37</sup>; SGA, small for gestational age at birth; UHC, Universal Health Coverage.