



FLOWERING NEWSLETTER REVIEW

Gynoecium size and ovule number are interconnected traits that impact seed yield

Mara Cucinotta¹, Maurizio Di Marzo¹, Andrea Guazzotti¹, Stefan de Folter², Martin M. Kater¹ and Lucia Colombo^{1,*}

¹ Dipartimento di Bioscienze, Università degli Studi di Milano, Via Celoria 26, 20133 Milan, Italy

² Unidad de Genómica Avanzada (UGA-Langebio), Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional (CINVESTAV-IPN), Km. 9.6 Libramiento Norte, Carretera Irapuato-Leon, CP 36824 Irapuato, Gto., Mexico

* Correspondence: lucia.colombo@unimi.it

Received 13 October 2019; Editorial decision 13 January 2020; Accepted 24 January 2020

Editor: Frank Wellmer, Trinity College Dublin, Ireland

Abstract

Angiosperms form the largest group of land plants and display an astonishing diversity of floral structures. The development of flowers greatly contributed to the evolutionary success of the angiosperms as they guarantee efficient reproduction with the help of either biotic or abiotic vectors. The female reproductive part of the flower is the gynoecium (also called pistil). Ovules arise from meristematic tissue within the gynoecium. Upon fertilization, these ovules develop into seeds while the gynoecium turns into a fruit. Gene regulatory networks involving transcription factors and hormonal communication regulate ovule primordium initiation, spacing on the placenta, and development. Ovule number and gynoecium size are usually correlated and several genetic factors that impact these traits have been identified. Understanding and fine-tuning the gene regulatory networks influencing ovule number and pistil length open up strategies for crop yield improvement, which is pivotal in light of a rapidly growing world population. In this review, we present an overview of the current knowledge of the genes and hormones involved in determining ovule number and gynoecium size. We propose a model for the gene regulatory network that guides the developmental processes that determine seed yield.

Keywords: Gynoecium, hormones, organ boundary, ovule number, ovule primordia, pistil, seed yield.

Introduction

Life on earth is affected by plants in varied ways. Of the estimated 400 000 extant plant species, approximately 94% are seed plants (Govaerts, 2001; Willis, 2017). This demonstrates that seed development and dispersion strategies greatly contributed to the success of this organismal group. The vast majority of seed plants are angiosperms and only a comparatively small number are gymnosperms. Both plant divisions produce ovules; however, only angiosperm species produce flowers, and as another selective advantage, each flower produces one or

more gynoecia that protect and nourish the ovules. Following fertilization, the gynoecium (or pistil) generally develops into a fruit and ovules develop into seeds.

Depending on the species, the gynoecium consists of one or more carpels, which can be fused or unfused (Endress and Igersheim, 2000). The Arabidopsis gynoecium consists of two fused carpels (Smyth *et al.*, 1990; Alvarez-Buylla *et al.*, 2010). Along the margins where the carpels fuse, a meristematic tissue, termed the carpel margin meristem (CMM), is formed.

Table 1. Genes involved in determining gynoecium size and/or ovule number

Gene name	Family or protein type	Gynoecium size	Ovule number	Reference
<i>ANT</i>	AP2/EREBP transcription factor	<i>ant-9</i> ↓ <i>ant-4</i> ↓ 35S:: <i>ANT</i> ↑	<i>ant-1</i> ↓ <i>ant-3</i> ↓ <i>ant-4</i> ↓ <i>ant-9</i> ↓	Elliott <i>et al.</i> (1996), Liu <i>et al.</i> (2000), Azhakanandam <i>et al.</i> (2008), Krizek (2009), Wynn <i>et al.</i> (2014)
<i>ARGOS</i>	ARGOS protein	35S:: <i>ARGOS</i> ↑		Hu <i>et al.</i> (2003)
<i>CRC</i>	YABBY transcription factor	<i>crc-1</i> ↓		Gross <i>et al.</i> (2018)
<i>SPT</i>	bHLH transcription factor	<i>spt-2</i> ↓	<i>spt-2</i> ↓	Heisler <i>et al.</i> (2001), Alvarez and Smyth (2002), Nahar <i>et al.</i> (2012)
<i>ETT (ARF3)</i>	ARF transcription factor	<i>ett-1</i> ↓ <i>ett-2</i> ↓		Sessions <i>et al.</i> (1997), Nemhauser <i>et al.</i> (2000)
<i>HEC1, HEC2, HEC3</i>	bHLH transcription factor	<i>hec1 hec2 hec3</i> ↓		Gremski <i>et al.</i> (2007)
<i>ARR1, ARR10, ARR12</i>	Type-B ARR transcription factor	<i>arr1 arr10 arr12</i> ↓	<i>arr1 arr10 arr12</i> ↓	Reyes-Olalde <i>et al.</i> (2017)
<i>CRF2, CRF3, CRF6</i>	ERF transcription factor	<i>crf2 crf3 crf6</i> ↓	<i>crf2 crf3 crf6</i> ↓	Cucinotta <i>et al.</i> (2016)
<i>PIN1</i>	PIN auxin efflux carrier	<i>pin1</i> ↓	<i>pin1</i> ↓ <i>pin1-5</i> ↓	Okada <i>et al.</i> (1991), Bencivenga <i>et al.</i> (2012), Cucinotta <i>et al.</i> (2016)
<i>CKX3, CKX5</i>	CKX cytokinin oxidase/dehydrogenase protein	<i>ckx3 ckx5</i> ↑	<i>ckx3 ckx5</i> ↑	Bartrina <i>et al.</i> (2011)
<i>UGT85A3, UGT73C1</i>	UDP-glucosyl transferase	35S:: <i>UGT85A3</i> ↓ 35S:: <i>UGT73C1</i> ↓	35S:: <i>UGT85A3</i> ↓ 35S:: <i>UGT73C1</i> ↓	Cucinotta <i>et al.</i> (2018)
<i>SAUR8, SAUR10, SAUR12</i>	SAUR-like auxin-responsive protein family	35S:: <i>SAUR8</i> ↑ 35S:: <i>SAUR10</i> ↑ 35S:: <i>SAUR12</i> ↑		van Mourik <i>et al.</i> (2017)
<i>BZR1</i>	Brassinosteroid signalling regulatory protein	<i>bzr1-1D</i> ↑	<i>bzr1-1D</i> ↑	Huang <i>et al.</i> (2013)
<i>BIN2</i>	ATSK (shaggy-like kinase) family	<i>bin2</i> ↓	<i>bin2</i> ↓	Huang <i>et al.</i> (2013)
<i>DET2</i>	3-Oxo-5- α -steroid 4-dehydrogenase protein	<i>det2</i> ↓	<i>det2</i> ↓	Huang <i>et al.</i> (2013)
<i>BRI1</i>	Leucine-rich receptor-like protein kinase protein	<i>bri1-5</i> ↓	<i>bri1-5</i> ↓	Huang <i>et al.</i> (2013)
<i>CYP85A2</i>	Cytochrome p450 enzyme		<i>cyp85a2-1</i> ↓ <i>cyp85a2-2</i> ↓	Nole-Wilson <i>et al.</i> (2010b)
<i>SEU</i>	Transcriptional adaptor	<i>seu-1</i> ↓	<i>seu-1</i> ↓	Nole-Wilson <i>et al.</i> (2010b)
<i>CTR1</i>	RAF homologue of serine/threonine kinase	<i>ctr1-1</i> ↓		Carbonell-Bejerano <i>et al.</i> (2011)
<i>REV</i>	Homeobox-leucine zipper protein		<i>ant rev</i> ↓	Nole-Wilson <i>et al.</i> (2010a)
<i>L-UG</i>	WD40/YVTN repeat-like-containing domain transcription factor		<i>lug-1</i> ↓ <i>lug-3</i> ↓	Azhakanandam <i>et al.</i> (2008)
<i>PAN</i>	bZIP transcription factor	<i>ant pan</i> ↓ <i>seu pan</i> ↓	<i>ant pan</i> ↓ <i>seu pan</i> ↓	Wynn <i>et al.</i> (2014)
<i>HLL</i>	Ribosomal protein L14p/L23e	<i>hll</i> ↓	<i>hll</i> ↓	Schneitz <i>et al.</i> (1998), Skinner <i>et al.</i> (2001)
<i>SIN2</i>	P-loop containing nucleoside triphosphate hydrolase superfamily protein	<i>sin-2</i> ↓	<i>sin-2</i> ↓	Broadhvest <i>et al.</i> (2000)
<i>YUC1, YUC4</i>	Flavin-binding monooxygenase protein		<i>yuc1 yuc4</i> ↓	Cheng <i>et al.</i> (2006)
<i>AHK2, AHK3, CRE1</i>	Histidine kinase		<i>cre1-12 ahk2-2 ahk3-3</i> ↓	Bencivenga <i>et al.</i> (2012)
<i>CUC1, CUC2</i>	NAC transcription factor		<i>cuc1 cuc2</i> ↓ <i>pSTK::CUC1/RNAi</i> <i>cuc2-1</i> ↓ 35S:: <i>MIR164A</i> ↓	Galbiati <i>et al.</i> (2013)
<i>MIR164A</i>	microRNA		35S:: <i>MIR164A</i> ↓	Gonçalves <i>et al.</i> (2015)
<i>GAI, RGA, RGL2</i>	GRAS transcription factor	<i>gaiT6 rgaT2 rgl2-1</i> ↓	<i>gaiT6 rgaT2 rgl2-1</i> ↓	Gomez <i>et al.</i> (2018)
<i>GID1A, GID1B</i>	α/β -Hydrolase superfamily protein		<i>gid1ab</i> ↑	Gomez <i>et al.</i> (2018)
<i>REM22</i>	B3 protein transcription factor		<i>rem22-1</i> ↑	Gomez <i>et al.</i> (2018)
<i>UNE16</i>	Homeodomain-like superfamily protein		<i>une16-1</i> ↓	Gomez <i>et al.</i> (2018)
<i>NERD1</i>	GW repeat- and PHD-finger-containing protein NERD		<i>nerd1-2</i> ↓ <i>nerd1-4</i> ↓	Yuan and Kessler (2019)
<i>ONA2</i>	Unknown protein		<i>ona2</i> ↓	Yuan and Kessler (2019)
<i>ASHH2</i>	Histone-lysine N-methyltransferase		<i>ashh2</i> ↓	Grini <i>et al.</i> (2009)

Up- and down-pointing arrows represent how the mutant phenotype impacts either gynoecium size or ovule number.

Subsequently, three layers of primordium cells form a finger-like structure during stage 10, which then differentiates into three regions along the proximal–distal axis: the funiculus, the chalaza, and the nucellus (Schneitz *et al.*, 1995). These three regions undergo distinct but interdependent developmental processes. The nucellus is the site of megasporogenesis, where the megaspore mother cell differentiates and locates to the up-most, central, and subepidermal position of the digit-shaped ovule primordium (reviewed in Pinto *et al.*, 2019). The chalaza is the region from which the inner and outer integuments develop, and these finally envelop and protect the embryonic sac. The funiculus remains attached to the gynoecium via the placental tissue and this connection is required for the transport of nutrients to the ovule (Fig. 1). For this reason, the placental tissue is fundamental for ovule primordia formation, and for determining their number and maintenance.

In Arabidopsis, placental tissue differentiates from the CMM, which is the central ridge of cells that fuse and give rise to the septum. Placental tissue differentiates along the length of the septum adjacent to the lateral walls (Alvarez and Smyth, 2002; Nole-Wilson *et al.*, 2010a; Reyes-Olalde *et al.*, 2013). Communication between transcription factors and hormones is essential to maintain the meristematic activity of the placenta, to determine the sites of ovule initiation and ovule identity, and to establish the distance between two adjacent ovules (Cucinotta *et al.*, 2014). Several genes that are important for placenta development have been described in the literature (reviewed by Cucinotta *et al.*, 2014; Reyes-Olalde and de Folter, 2019), including *AINTEGUMENTA* (*ANT*), *CUP-SHAPED COTYLEDON 1* (*CUC1*) and *CUC2*, *LEUNIG* (*LUG*), *MONOPTEROS* (*MP*), and *PERIANTHIA* (*PAN*) (Fig. 1; Table 1).

AINTEGUMENTA encodes an AP2 transcription factor (Klucher *et al.*, 1996) and positively regulates organ size via determining cell number and meristematic competence. *Ant* mutants have fewer and smaller floral organs than the wild-type. In particular, the *ant-9* mutant is characterized by unfused carpels at the tip of the pistil (Elliott *et al.*, 1996), whereas in *ant-4*, the size of floral organs is reduced (Krizek, 2009). In contrast to these mutant phenotypes, Arabidopsis plants that overexpress *ANT* possess larger floral organs than the wild-type (Mizukami and Fischer, 2000). Expression of *ANT* is controlled by *AUXIN-REGULATED GENE INVOLVED IN ORGAN SIZE* (*ARGOS*), an auxin-inducible gene (Hu *et al.*, 2003). When *ARGOS* is overexpressed, floral organs become enlarged, resulting in longer siliques than those of wild-type (Hu *et al.*, 2003). This was one of the first pieces of evidence that implicated a key role for auxin in pistil development.

ANT expression initiates in the placenta and is maintained throughout all stages of ovule development, in particular in the chalaza region and in the integuments. The reduced ovule number phenotype of the *ant* mutant is exacerbated when it is combined with other mutations that affect CMM and placenta development, such as *revoluta* (*rev*), suggesting that the activity of the *REV* gene, which encodes a class III homeodomain leucine zipper transcription factor, is also required for placenta formation (Nole-Wilson *et al.*, 2010a). *ANT* interacts with the transcriptional repressor SEUSS (*SEU*) and simultaneous

loss of both protein activities severely affects placenta development and leads to a complete loss of ovule formation. When a weaker *ant-3* allele was combined with *seu-3*, placenta development was maintained but the number of ovules that initiated was reduced to approximately half of that observed in Col-0 wild-type plants (Azhakanandam *et al.*, 2008). Another transcriptional co-regulator involved in gynoecium patterning is *LEUNIG* (*LUG*). Strong *lug-1* and intermediate *lug-3* alleles show a failure in ridge fusion and a reduction in the amount of placental tissue, with a consequent decrease in the number of ovules formed (Liu *et al.*, 2000). The combination of *lug* and *ant* mutations results in gynoecia that are unable to develop ovules (Liu *et al.*, 2000). The loss of ovules in the *ant* and *seu* backgrounds is strongly enhanced by mutations in the *PERIANTHIA* (*PAN*) gene, which encodes a bZIP transcription factor that is expressed in the gynoecium medial ridge, placenta, and ovules, where it promotes ovule formation (Wynn *et al.*, 2014).

Similar to *ANT*, factors important for integument growth often affect ovule primordium formation. Two examples are *HUELLENLOS* (*HLL*) and *SHORT INTEGUMENTS 2* (*SIN2*). *HLL* encodes a mitochondrial ribosomal protein and its mutation is associated with smaller gynoecia and a 10% reduction in the number of ovules (Schneitz *et al.*, 1998; Skinner *et al.*, 2001). Shorter gynoecia that bear fewer ovules are also observed in the *sin2* mutant; however, more interestingly, the absence of *SIN2* function leads to an abnormal distribution of ovules along the placenta (Broadhvest *et al.*, 2000), in which the distance between ovules is greater than in the wild-type; thus, a reduction in ovule number is caused by a reduction in gynoecium size and by the reduced ability of the placental tissue to initiate ovule primordia. *SIN2* encodes a mitochondrial DAR GTPase and, similar to *HLL*, is hypothesized to function in mitochondrial ribosome assembly (Hill *et al.*, 2006). Notably, these two ribosomal proteins, which are targeted to the mitochondria, are necessary for ovule primordium formation, and it has been suggested that impaired mitochondrial function might cause cell-cycle arrest in the placenta and subsequently in the ovule integuments (Broadhvest *et al.* 2000).

Complex hormonal communication promotes ovule initiation and determines pistil size

Plant organogenesis requires cells to proliferate, grow, and differentiate in a coordinated way. The intercellular communication required during organ initiation is mediated by different phytohormones (Davies, 2004; Vanstraelen and Benková, 2012; Schaller *et al.*, 2015; Marsch-Martínez and de Folter, 2016). As will be discussed in this review, auxins, cytokinins (CKs), gibberellins (GAs), and brassinosteroids (BRs) all play fundamental roles in ovule primordium formation (Fig. 1).

In most auxin-related mutants, defects in gynoecium formation lead to the reduction or absence of placental tissue and the corresponding absence of ovules (reviewed in Balanzá *et al.*, 2006; Larsson *et al.*, 2013; Cucinotta *et al.*, 2014). This phenotype is common to all mutants in which auxin synthesis

or transport pathways are compromised, such as *yucca1 yucca4* (*yuc1 yuc4*) (Cheng *et al.*, 2006) and *pin1-1* (Okada *et al.*, 1991) or is similar to that following treatment with the polar auxin transport inhibitor 1-naphthyl phthalamic acid (NPA) (Nemhauser *et al.*, 2000).

Polar auxin transport is mediated by the PINFORMED1 (PIN1) efflux transporter and is required to create a zone with an auxin concentration maximum in the placenta, where the founder cells of the ovule primordia will be specified (Benková *et al.*, 2003; Ceccato *et al.*, 2013; Galbiati *et al.*, 2013). Subsequently, the orientation of PIN1 within the membrane relocalizes and redirects auxin flow, establishing a gradient with a maximum at the apices of the formed primordia. In developing organs, auxin distribution can be monitored *in vivo* by imaging a synthetic auxin-inducible promoter, *DR5*. In plants that express green fluorescent protein (GFP) from the *DR5* promoter, green fluorescence is detected at the apices of the ovule primordia, consistent with PIN1-mediated auxin flow directed to the apex (Benková *et al.*, 2003; Galbiati *et al.*, 2013). The weak *pin1-5* mutant allele can produce some flowers in which the pistils have slightly reduced valves, which on average contain only nine ovules (Bennett *et al.*, 1995; Sohlberg *et al.*, 2006; Bencivenga *et al.*, 2012).

CKs occupy a central role in the regulation of cell division and cell differentiation. They are positive regulators of ovule formation, as demonstrated by the phenotype of mutants in which CK pathways are altered. In the *ckx3 ckx5* double mutant, the degradation of CKs is compromised and the consequent increase in the levels of these hormones leads to an increased activity of the reproductive meristem (Bartrina *et al.*, 2011). Moreover, the longer than normal gynoecia of *ckx3 ckx5* double mutants contain about twice as many ovules as those of the wild-type, indicating an increase in the meristematic capacity of placental tissue (Bartrina *et al.*, 2011). By contrast, reduced ovule formation is observed in mutants in which the synthesis or perception of CKs is compromised. Plants that carry mutations in genes that encode all three CK receptors, *cytokinin response 1* (*cre1-12*), *histidine kinase2* (*ahk2-2*), and *ahk3*, develop five ovules per pistil on average, in addition to showing pleiotropic growth defects (Higuchi *et al.*, 2004; Bencivenga *et al.*, 2012). The AHK2 and AHK3 receptors are expressed throughout ovule development, from the early stages until maturity, whereas *CRE1/AHK4* is expressed in the chalaza region and subsequently in the integuments, suggesting that AHK2 and AHK3 preferentially contribute to ovule primordium formation (Bencivenga *et al.*, 2012). The ovule and gynoecium phenotype of the *cre1-12 ahk2-2 ahk3-3* triple mutant resembles that of the weak *pin1-5* mutant allele (Bencivenga *et al.*, 2012). This similarity is due to the downregulation of *PIN1* expression in the triple mutant, suggesting that during the early stages of ovule development, CK activates *PIN1* expression. Bencivenga *et al.* (2012) showed that treating inflorescences with the synthetic CK 6-benzylaminopurine (BAP) increases *PIN1* expression in the gynoecium. Strikingly, treatment with BAP causes the formation of on average 20 additional ovule primordia in each gynoecium, which are positioned between the existing primordia formed before the treatment. This suggests that placental tissue at the boundaries between ovules

maintains meristematic competence. During root development, CK affects auxin polar transport via PIN1 both at the transcriptional and post-transcriptional levels. In contrast to the situation in the gynoecium, CK negatively regulates the expression of *PIN1* in the root and controls the endorecycling of PIN1 from the membrane to direct it to vacuoles for lytic degradation (Ruzicka *et al.*, 2009; Marhavý *et al.*, 2011). In roots, CYTOKININ RESPONSE FACTORS (CRFs), especially CRF2, CRF3, and CRF6, transcriptionally regulate *PIN1* by binding to its promoter at the *cis*-regulatory *PIN* CYTOKININ RESPONSE ELEMENT (PCRE) (Šimášková *et al.*, 2015) and modulate its expression in response to CK. Similarly, CRFs also mediate *PIN1* expression in ovules in response to CK (Cucinotta *et al.*, 2016). Indeed, *PIN1* expression is reduced in the *arf2 arf3 arf6* (*arf2/3/6*) triple mutant and cannot be increased by CK treatment. The placenta in *arf2/3/6* is also shorter, but this is not sufficient to explain the 30% decrease in ovule number as ovule density is lower in *arf2/3/6* than in the wild-type (Cucinotta *et al.*, 2016). Because *PIN1* expression in *arf2/3/6* was unresponsive to CK application, the mutant was significantly less sensitive to CK treatment than the wild-type with regard to an increase in ovule number and pistil length. Auxin also regulates *CRF2*, which is a direct target of the auxin response factor (ARF) AUXIN RESPONSE FACTOR 5/MONOPTEROS (ARF5/MP) (Schlereth *et al.*, 2010), highlighting another convergence point between auxin and CK.

Another ARF family member that is required for appropriate apical-basal gynoecium patterning is ARF3/ETTIN (ETT). The *ett* mutant is characterized by a shorter ovary with an elongated style and gynophore (Sessions *et al.*, 1997). A similar gynoecium phenotype resulted from treatment with the auxin transport inhibitor (NPA), suggesting that ETT plays a key role in auxin signalling along the apical-basal gynoecium axis (Nemhauser *et al.*, 2000). Moreover, ETT restricts the expression domain of *SPATULA* (*SPT*), which encodes a basic helix-loop-helix (bHLH) transcription factor (Heisler *et al.*, 2001). Mutations in *SPT* causes a split-carpel phenotype in the apical part of the gynoecium, leading to a slight reduction in ovule number (Alvarez and Smyth, 1999; Nahar *et al.*, 2012). *SPT* dimerizes with another bHLH transcription factor, INDEHISCENT (*IND*), to repress the expression of *PINOID* (Girin *et al.*, 2011), which encodes a serine/threonine kinase that regulates PIN1 polarization via phosphorylation (Friml *et al.*, 2004). The repression of *PID* by *SPT* and *IND* allows the formation of a radially symmetric auxin ring in the upper part of the gynoecium that is required for correct style and stigma development (Moubayidin and Østergaard, 2014).

Furthermore, *SPT* interacts with the three closely related bHLH transcription factors, HECATE1 (*HEC1*), *HEC2*, and *HEC3* (Gremski *et al.*, 2007), and similar to *ett*, *hec-1 hec-2 hec-3* triple mutants possess an elongated style and shorter ovaries. The *HEC* proteins and *SPT* promote auxin transport in concert by activating *PIN1* and *PIN3* expression (Schuster *et al.*, 2015) and also transcriptionally activate the type-A ARABIDOPSIS RESPONSE REGULATORS (*ARR-As*), which are negative regulators of CK signalling (Schuster *et al.*, 2015). Via this dual

action on auxin transport and CK response, HECs and SPT regulate wild-type gynoecium fusion at the apex, and style and stigma development. Furthermore, SPT alone in the medial domain activates the type-B ARRs, especially ARR1, which are positive regulators of CK signalling. The *arr1 arr10 arr12* triple mutant possesses a shorter gynoecium and significantly fewer ovules than the wild-type (Reyes-Olalde *et al.*, 2017).

In addition to auxin localization, correct auxin signalling is also required for wild-type gynoecium development, as confirmed by a recent study on members of the Small Auxin-Upregulated RNA (SAUR) family, which were initially identified as short transcripts that were rapidly upregulated in response to auxin (McClure and Guilfoyle, 1987). When SAUR8, SAUR10, and SAUR12 are ectopically overexpressed in Arabidopsis, the gynoecium and resulting siliques are longer than in wild-type, suggesting that auxin positively regulates gynoecium length and, probably indirectly, silique length (van Mourik *et al.*, 2017). Notably, SAUR gene expression increased 100-fold following combined auxin and BR treatment (van Mourik *et al.*, 2017). BRs are clearly involved in pistil growth and ovule number specification; gynoecia of the enhanced BR-signalling mutant *brassinazole-resistant 1-1D (bzi1-1D)* not only contained more ovules than wild-type but they were also longer. By contrast, BR-deficient mutants such as *de-etiolated 2 (det-2)*, *brassinosteroid insensitive 1 (bri1-5)* and *brassinosteroid-insensitive 2 (bin2-1)* developed shorter pistils with fewer ovules (Huang *et al.*, 2013).

The involvement of BRs in gynoecium and ovule development was also confirmed by Nole-Wilson *et al.* (2010b), who observed that a reduction in the expression of *CYP85A2*, which encodes an enzyme involved in the final step of brassinolide biosynthesis (Nomura *et al.*, 2005), enhances the *seuss* mutant phenotypic disruptions in ovules and gynoecia (Nole-Wilson *et al.*, 2010b).

CUP-SHAPED COTYLEDON 1 and 2 function synergistically with auxin and cytokinins

During ovule primordium formation, CK homeostasis requires two NAC-domain transcription factors, CUP-SHAPED COTYLEDON 1 (CUC1) and CUC2. These are expressed in lateral organ boundaries and function redundantly during organ boundary determination. *CUC1* and *CUC2* are expressed in the septum and placenta, and following the emergence of ovule primordia, *CUC2* expression is restricted to the borders between the ovules (Ishida *et al.*, 2000; Galbiati *et al.*, 2013; Gonçalves *et al.*, 2015). The *CUC1* and *CUC2* genes are both post-transcriptionally regulated by *miR164* microRNAs (Laufs *et al.*, 2004; Mallory *et al.*, 2004). Gynoecia of the *in vitro* regenerated *cuc1 cuc2* mutant as well as *cuc2-1 pSTK::CUC1_RNAi* plants have reduced ovule numbers. The *cuc1 cuc2* double mutant has on average fewer than 10 ovules per pistil (Ishida *et al.*, 2000), whereas *cuc2-1 pSTK::CUC1_RNAi* plants, in which *CUC1* was specifically silenced in the placenta and ovules, showed a 20% reduction in ovule number, but gynoecium length was not affected. In pistils of these

plants, ovules were more widely spaced when compared with the wild-type (Galbiati *et al.*, 2013). This result was confirmed by silencing *CUC1* and *CUC2* by overexpressing *MIR164A*, which strongly reduced ovule number, indicating a major contribution of *CUC1* and *CUC2* to ovule initiation (Gonçalves *et al.*, 2015). The analysis of PIN1-GFP expression in *cuc2-1 pSTK::CUC1_RNAi* plants revealed that *CUC1* and *CUC2* redundantly promote *PIN1* expression and *PIN1* membrane localization in ovules. Treatment with BAP increased *PIN1* expression and complemented the reduced ovule number phenotype of *cuc2-1 pSTK::CUC1_RNAi* plants (Galbiati *et al.*, 2013). Therefore, CKs act downstream from or in parallel with *CUC1* and *CUC2* to induce the expression of *PIN1*. Recently, it has been demonstrated that *CUC1* and *CUC2* induce CK responses *in vivo* and function upstream of CK by transcriptionally repressing *UGT73C1* and *UGT85A3*, which encode two enzymes involved in CK inactivation (Cucinotta *et al.*, 2018). Consistent with this result, the concentration of inactive CK glucosides was higher in *cuc2-1 pSTK::CUC1_RNAi* inflorescences than in wild-type plants.

The expression of *CUC1* and *CUC2* is also linked with auxin signalling: their expression pattern coincides with that of the auxin response factor ARF5/MP (see above) and both genes are downregulated in pistils of the weak *mp-S319* mutant allele (Galbiati *et al.*, 2013). During the early stages of placenta development and ovule formation, ARF5/MP directly transcriptionally activates *CUC1* and *CUC2*, but also *ANT*. The observation that BAP treatment did not complement the ovule number phenotype of *ant-4* suggests that *ANT* functions independently of *CUC1* and *CUC2*. This is further supported by the additive effects on the reduction in ovule number observed in *ant-4 cuc2-1 pSTK::CUC1_RNAi* plants (Galbiati *et al.*, 2013). Together these data suggest that *ANT* promotes cell proliferation, whereas *CUC1* and *CUC2* regulate CK homeostasis and auxin transport. Although *CUC3* shares high similarity with *CUC1* and *CUC2*, the *cuc3* mutant was not affected in ovule initiation and number, but together with *CUC2*, *CUC3* promotes ovule separation; this is reflected by the *cuc2 cuc3* double mutant, which produces seeds that result from the fusion of two ovules (Gonçalves *et al.*, 2015). These results suggest that specific *CUC* genes independently promote ovule initiation and ovule separation.

Lee *et al.* (2009) identified LATERAL ORGAN FUSION 1 (LOF1) to be involved in lateral organ separation and to functionally overlap with *CUC2* and *CUC3*. The *LOF1* gene is expressed at the base of ovule primordia and its overexpression results in a wrinkled pistil with an enlarged replum, an amorphous septum and an irregular ovule distribution (Gomez *et al.*, 2011).

The role of gibberellins in ovule primordium formation

GAs are involved in key developmental processes throughout the plant life cycle, from seed germination in particular, to flowering time (reviewed in Hedden and Sponsel, 2015; Rizza and Jones, 2019), but their involvement in ovule initiation

has only recently been demonstrated. Gomez and colleagues (2018) showed that DELLA proteins, which belong to a sub-family of the plant-specific GRAS family of transcriptional regulators that repress GA signalling, positively regulate ovule number in Arabidopsis. In addition to DELLA proteins, the GA signalling core includes the GA receptor *GID1*. When *GID1* binds bioactive GA, the GA-*GID1*-DELLA complex is formed and triggers the polyubiquitination and degradation of DELLA proteins. The *della* triple mutant *gaiT6 rgaT2 rgl2-1* produces fewer ovules than wild-type (Gomez et al., 2018). By contrast, the gain-of-function DELLA mutant *gai-1*, which cannot be degraded upon GA sensing, produced more ovules. Consistent with this observation, the double *gid1a gid1b* mutant, which cannot perceive GA, forms more ovules than the wild-type, demonstrating a negative correlation between GAs and ovule number (Gomez et al., 2018). The *GAI*, *RGA*, *RGL2*, *GID1a*, and *GID1b* genes are expressed in placental tissue and outgrowing ovules. The reduction in ovule number was more dramatic in the *gaiT6 rgaT2 rgl2-1* triple mutant than that in ovary length, resulting in a lower ovule density, whereas the dominant *gai-1* mutant has an increased ovule/placenta ratio, suggesting that GAs predominantly affect ovule initiation and not placenta elongation.

Other evidence to demonstrate that DELLA proteins promote ovule formation derives from an experiment in which the expression of the stable mutant protein *rgaΔ17* under the control of the *ANT* promoter in the placenta resulted in the formation of 20% more ovules than in control lines (Gomez et al., 2018). This effect of GAs on the number of developing ovules was not correlated with auxin signalling or transport, and neither *PIN1* localization nor *DR5* expression was affected by GA treatment or DELLA activity (Gomez et al., 2018).

Confirmation of a positive role for *RGL2* in determining ovule number came from the analysis of transgenic lines in which *RGL2*-dependent GA signalling was blocked by the expression of a dominant version of *RGL2* (*pRGL2:rgl2Δ17*) (Gómez et al., 2019). Pistils of *pRGL2:rgl2Δ17* plants contained 10% more ovules than those of the wild-type, whereas pistil length did not differ, indicating that the main function of *rgl2Δ17* is to positively promote ovule primordium formation but not placenta elongation (Gómez et al., 2019). Furthermore, Gomez et al. (2018) identified *REPRODUCTIVE MERISTEM 22* (*REM22*) and *UNFERTILIZED EMBRYO SAC 16* (*UNE16*) via transcriptomic analysis to be DELLA targets that are positive regulators of ovule initiation. *REM22* is a B3 family transcription factor that is expressed in the placenta (Mantegazza et al., 2014) and increased *REM22* expression in the *rem22-1* enhancer allele significantly increases ovule number. *UNE16* is a transcription factor involved in embryo sac development and the knockdown allele *une16-1* produces fewer ovules. Because *UNE16* expression is regulated by BRs (Pagnussat et al., 2005; Sun et al., 2010), it represents a potential nexus for crosstalk between GAs and BRs in ovule initiation. The establishment of GA as an important additional component of the ovule regulatory network has introduced an additional layer of complexity to the current model for ovule initiation and it remains to be established how GAs integrate into this model. GAs might function antagonistically to CKs

and BRs, which in contrast to GAs, positively regulate pistil size and ovule number.

Finally, the *ctr1-1* constitutive ethylene-responsive mutant possesses a shorter gynoecium at anthesis compared with wild-type and a delay in the response to GA₃ treatment that induces gynoecium senescence, suggesting that ethylene affects gynoecium size, probably by interactions with GA pathways (Carbonell-Bejerano et al., 2011).

In conclusion, there is ample evidence for complex interactions between different hormonal pathways that together determine ovule number and pistil size.

Ovule number: the ecotype matters

It has been known for 20 years that the number of ovules varies hugely among different Arabidopsis ecotypes (diploid accessions) (Alonso-Blanco et al., 1999): for example, the Landsberg *erecta* accession produces 20% more ovules than the Cape Verde Islands (Cvi) accession. Recently, 189 Arabidopsis accessions from the Arabidopsis Biological Resource Center were analysed for differences in ovule number and they display a remarkable degree of variation, ranging from 39 to 82 ovules per pistil (Yuan and Kessler, 2019). The commonly used reference accession Col-0 lies in the middle of the range, with a mean ovule number of 63, which is strongly dependent on experimental growth conditions. Ovule number, in contrast to, for instance, flowering time, does not correlate with geographical origin (Stinchcombe et al., 2004; Yuan and Kessler, 2019). By conducting a genome-wide association study on these 189 accessions, two loci associated with ovule number were identified (Yuan and Kessler, 2019): *NEW ENHANCER OF ROOT DWARFISM* (*NERD1*) and *OVULE NUMBER ASSOCIATED 2* (*ONA2*). Mutation of *NERD1* or *ONA2* leads to a significant reduction in ovule number, with a stronger phenotype in the *nerd1-2* and *nerd1-4* alleles. *ONA2* encodes a protein of unknown function and was not further analysed. In addition to a reduction in ovule number, *nerd* mutants display additional severe male and female fertility defects. *NERD1* encodes an integral membrane protein mainly localized to the Golgi. Notably, *NERD1* expression is lower in Altai-5 and Kas-2 accessions, which have low ovule numbers (Yuan and Kessler, 2019), but high *NERD1* expression in Altai-5 leads to a significant increase in ovule number. However, overexpression of *NERD1* in Col-0 plants did not affect ovule number, indicating that *NERD1* function in determining ovule number is background-dependent (Yuan and Kessler, 2019).

Considerable genetic variation in ovule number was also described for F₁ triploids of different Arabidopsis genotypes by Duszynska et al. (2013), who observed differences in ovule number between genetically identical F₁-hybrid offspring, after crossing parental genome excess lines (2m:1p with 1m:2p). These effects can only be explained by epigenetic mechanisms that affect genes controlling ovule number, for example DNA or histone methylation. The analysis of null alleles of *ASH1 HOMOLOG 2* (*ASH2*), which show a remarkable 80% reduction in ovule number, provided a clear example of

the involvement of histone methylation in determining ovule number (Grini *et al.*, 2009). The transcriptional state of the *ASH2* locus remains active during development via H3K36 trimethylation (Xu *et al.*, 2008). It will be highly relevant to study the effect of epigenetic modifications induced by biotic and abiotic stresses in determining ovule number. Epigenetic responses to stress are fundamental to create the plasticity required for plant survival, especially considering that plants are sessile organisms. These epigenetic changes can be temporally transmitted, even in the absence of the original stress (Iglesias and Cerdán, 2016). Furthermore, variation in ovule number in response to fluctuations in environmental conditions, such as temperature, can be used to understand the plasticity and inheritability of (epigenetic) adaptation and response to temperature stress. Variation in ovule number under stress conditions is, of course, also highly relevant from an ecological, environmental, and evolutionary perspective.

Ovule number decreases with ageing

Ovule number varies throughout inflorescence development: early flowers developing on the main inflorescence (from the fifth to the twenty-fifth flower) of *Arabidopsis Ler* plants produced a relatively invariable number of ovules, whereas flowers that developed later had pistils with fewer ovules (Gomez *et al.*, 2018; Yuan and Kessler, 2019). Loss- and gain-of-function mutants of *DELLA* genes showed an increase in ovule number in early- and late-arising flowers (Gomez *et al.*, 2018). To minimize age-related variation in their genome-wide association studies, Yuan and Kessler (2019) only counted ovules in flowers 6–10 from the main inflorescence.

It has been reported for other plant species that flower position as well as size influences ovule number per flower. For example, in pomegranate, the number of ovules per flower was significantly influenced by flower size, with more ovules being produced in larger flowers (Wetzstein *et al.*, 2013).

Overall, when studying changes in ovule numbers it is important to be aware of the possible variation in the different flowers of the plant. Therefore, large numbers will have to be analysed using thorough statistical analyses, especially for genotypes that show only relatively minor changes.

A 'gold mine' for seed yield improvement within the Brassicaceae

Improving seed yield via the genetic manipulation of crops has historically been a central goal in agricultural research. The enormous body of data, which has been generated and shared by the scientific community over the past decades, represents a true 'gold mine' for translational and applied research. The determination of pistil size and ovule number may be considered one of the most straightforward traits that can be enhanced to improve overall seed yield in species characterized by multi-ovulate ovaries and the increasing amount of literature on this topic evidences an active and prolific research field. Although some questions concerning the networks controlling seed

number and pistil size remain open, comprehensive knowledge of the phytohormone interactions involved in these pathways is already available and applicable (Cucinotta *et al.*, 2014; Zúñiga-Mayo *et al.*, 2019; Reyes-Olalde and de Folter, 2019).

Understanding these developmental processes in *Arabidopsis* can inform promising strategies for knowledge transfer to closely related and agronomically important crops. Rapeseed (*Brassica napus*), another Brassicaceae species, is an important breeding target, since it is a crop widely cultivated in Europe, Asia, Canada, and Australia. It is characterized by an oil-rich seed and its processing provides both rapeseed oil (used as edible vegetable oil or as biodiesel) and a by-product mostly used as cattle fodder (Snowdon *et al.*, 2007).

It has recently been demonstrated that *Arabidopsis* and *B. napus* share well-conserved response mechanisms to CK treatment (Zuñiga-Mayo *et al.*, 2018). Strikingly, exogenous CK application causes a reduction in silique length in *B. napus*. However, these shorter siliques contain increased ovule numbers and upon manual pollination, the plants show an increase in seed yield of 18%. Intriguingly, increases in ovule and seed number have also been observed in the offspring of the treated plants, suggesting that the mechanism has an underlying epigenetic basis (Zuñiga-Mayo *et al.*, 2018).

An increase in CK level has also been reported to beneficially affect seed yield in transgenic *B. napus* lines expressing the CK biosynthetic enzyme isopentenyltransferase (*IPT*) under the *Arabidopsis* promoter of the *AtMYB32* gene. An increase in seed yield of up to 23% was obtained in the transgenic lines that were analysed (Kant *et al.*, 2015).

CK homeostasis is mediated by CYTOKININ OXIDASES/DEHYDROGENASES (CKXs) during pistil and silique development in *Arabidopsis*. Remarkably, the expression level of *CKX* genes in *B. napus* is associated with silique length, and RNA-sequencing and qRT-PCR analyses revealed a significantly different expression level of *BnCKX5-1*, *5-2*, *6-1*, and *7-1* in two distinct cultivated varieties with long versus short siliques (Liu *et al.*, 2018). These findings open up promising strategies with which to modulate silique length in *B. napus* by manipulating *CKX* gene expression.

In addition to phytohormones, genetic knowledge from *Arabidopsis* can be successfully applied to *B. napus* crop improvement. Mutations in the K-box of the *Arabidopsis* orthologue of *APETALA1* in *B. napus* caused a significant increase in the number of seeds per plant (Shah *et al.*, 2018). These generated alleles could conceivably be introduced into a rapeseed breeding programme in field trials.

Germplasm of *B. napus* revealed substantial natural variation with respect to seed number per pod. Current rapeseed cultivars produce on average 20 seeds per pod, which is far lower than the maximum observed among the germplasm resources (Yang *et al.*, 2017). Moreover, genetic improvement promises to deliver a massive improvement in seed yield (Yang *et al.*, 2017). The gold mine of knowledge obtained from the closely related species *Arabidopsis* will certainly be fundamentally important in the exploitation of the encouraging genetic variation potential. Furthermore, it has recently been demonstrated that CRISPR-Cas9 technology can be efficiently applied to precisely induce targeted mutation in rapeseed (Braatz

et al., 2017), making it a powerful tool for future genetic improvement. Similarly, existing knowledge could be used to improve other Brassicaceae species, or even non-phylogenetically related species such as soybean.

Acknowledgements

The authors' work is supported by the H2020 Marie Skłodowska-Curie Actions of European Commission (H2020-MSCA-RISE-2015 EXPOSEED and MSCA-RISE-2014 PROCROP projects) also acknowledges the Mexican National Council of Science and Technology (CONACyT) grants FC-2015-2/1061 and CB2017-2018 A1-S-10126. The authors would like to thank John Chandler for editing the paper.

References

- Alonso-Blanco C, Blankestijn-de Vries H, Hanhart CJ, Koornneef M.** 1999. Natural allelic variation at seed size loci in relation to other life history traits of *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **96**, 4710–4717.
- Alvarez J, Smyth DR.** 1999. *CRABS CLAW* and *SPATULA*, two *Arabidopsis* genes that control carpel development in parallel with *AGAMOUS*. *Development* **126**, 2377–2386.
- Alvarez J, Smyth DR.** 2002. *CRABS CLAW* and *SPATULA* genes regulate growth and pattern formation during gynoecium development in *Arabidopsis thaliana*. *International Journal of Plant Sciences* **163**, 17–41.
- Alvarez-Buylla ER, Benítez M, Corvera-Poiré A, et al.** 2010. Flower development. *The Arabidopsis Book* **8**, e0127.
- Angenent GC, Colombo L.** 1996. Molecular control of ovule development. *Trends in Plant Science* **1**, 228–232.
- Azhakanandam S, Nole-Wilson S, Bao F, Franks RG.** 2008. *SEUSS* and *AINTEGUMENTA* mediate patterning and ovule initiation during gynoecium medial domain development. *Plant Physiology* **146**, 1165–1181.
- Balanzá V, Navarrete M, Trigueros M, Ferrándiz C.** 2006. Patterning the female side of *Arabidopsis*: the importance of hormones. *Journal of Experimental Botany* **57**, 3457–3469.
- Bartrina I, Otto E, Strnad M, Werner T, Schmülling T.** 2011. Cytokinin regulates the activity of reproductive meristems, flower organ size, ovule formation, and thus seed yield in *Arabidopsis thaliana*. *The Plant Cell* **23**, 69–80.
- Bencivenga S, Colombo L, Masiero S.** 2011. Cross talk between the sporophyte and the megagametophyte during ovule development. *Sexual Plant Reproduction* **24**, 113–121.
- Bencivenga S, Simonini S, Benková E, Colombo L.** 2012. The transcription factors *BEL1* and *SPL* are required for cytokinin and auxin signaling during ovule development in *Arabidopsis*. *The Plant Cell* **24**, 2886–2897.
- Benková E, Michniewicz M, Sauer M, Teichmann T, Seifertová D, Jürgens G, Friml J.** 2003. Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* **115**, 591–602.
- Bennett SRM, Alvarez J, Bossinger G, Smyth DR.** 1995. Morphogenesis in *pinoid* mutants of *Arabidopsis thaliana*. *The Plant Journal* **8**, 505–520.
- Bowman JL, Baum SF, Eshed Y, Putterill J, Alvarez J.** 1999. Molecular genetics of gynoecium development in *Arabidopsis*. *Current Topics in Developmental Biology* **45**, 155–205.
- Braatz J, Harloff HJ, Mascher M, Stein N, Himmelbach A, Jung C.** 2017. CRISPR-Cas9 targeted mutagenesis leads to simultaneous modification of different homoeologous gene copies in polyploid oilseed rape (*Brassica napus*). *Plant Physiology* **174**, 935–942.
- Broadhvest J, Baker SC, Gasser CS.** 2000. *SHORT INTEGUMENTS 2* promotes growth during *Arabidopsis* reproductive development. *Genetics* **155**, 899–907.
- Carbonell-Bejerano P, Urbez C, Granell A, Carbonell J, Perez-Amador MA.** 2011. Ethylene is involved in pistil fate by modulating the onset of ovule senescence and the GA-mediated fruit set in *Arabidopsis*. *BMC Plant Biology* **11**, 84.
- Ceccato L, Masiero S, Sinha Roy D, Bencivenga S, Roig-Villanova I, Ditengou FA, Palme K, Simon R, Colombo L.** 2013. Maternal control of *PIN1* is required for female gametophyte development in *Arabidopsis*. *PLoS ONE* **8**, e66148.
- Cheng Y, Dai X, Zhao Y.** 2006. Auxin biosynthesis by the *YUCCA* flavin monooxygenases controls the formation of floral organs and vascular tissues in *Arabidopsis*. *Genes & Development* **20**, 1790–1799.
- Colombo L, Battaglia R, Kater MM.** 2008. *Arabidopsis* ovule development and its evolutionary conservation. *Trends in Plant Science* **13**, 444–450.
- Cucinotta M, Colombo L, Roig-Villanova I.** 2014. Ovule development, a new model for lateral organ formation. *Frontiers in Plant Science* **5**, 117.
- Cucinotta M, Manrique S, Cuesta C, Benkova E, Novak O, Colombo L.** 2018. *CUP-SHAPED COTYLEDON1* (*CUC1*) and *CUC2* regulate cytokinin homeostasis to determine ovule number in *Arabidopsis*. *Journal of Experimental Botany* **69**, 5169–5176.
- Cucinotta M, Manrique S, Guazzotti A, Quadrelli NE, Mendes MA, Benkova E, Colombo L.** 2016. Cytokinin response factors integrate auxin and cytokinin pathways for female reproductive organ development. *Development* **143**, 4419–4424.
- Davies PJ.** 2004. *Plant hormones: biosynthesis, signal, transduction, action!* Dordrecht, The Netherlands: Kluwer Academic Publishers.
- Drews GN, Koltunow AM.** 2011. The female gametophyte. *The Arabidopsis Book* **9**, e0155.
- Duszynska D, McKeown PC, Juenger TE, Pietraszewska-Bogiel A, Geelen D, Spillane C.** 2013. Gamete fertility and ovule number variation in selfed reciprocal F1 hybrid triploid plants are heritable and display epigenetic parent-of-origin effects. *New Phytologist* **198**, 71–81.
- Elliott RC, Betzner AS, Huttner E, Oakes MP, Tucker WQ, Gerentes D, Perez P, Smyth DR.** 1996. *AINTEGUMENTA*, an *APETALA2*-like gene of *Arabidopsis* with pleiotropic roles in ovule development and floral organ growth. *The Plant Cell* **8**, 155–168.
- Endress PK.** 2011. Angiosperm ovules: diversity, development, evolution. *Annals of Botany* **107**, 1465–1489.
- Endress PK, Igersheim A.** 2000. Gynoecium structure and evolution in basal angiosperms. *International Journal of Plant Sciences* **161**, S211–S213.
- Friml J, Yang X, Michniewicz M, et al.** 2004. A PINOID-dependent binary switch in apical-basal PIN polar targeting directs auxin efflux. *Science* **306**, 862–865.
- Galbiati F, Sinha Roy D, Simonini S, et al.** 2013. An integrative model of the control of ovule primordia formation. *The Plant Journal* **76**, 446–455.
- Gasser CS, Broadhvest J, Hauser BA.** 1998. Genetic analysis of ovule development. *Annual Review of Plant Physiology and Plant Molecular Biology* **49**, 1–24.
- Gasser CS, Skinner DJ.** 2019. Development and evolution of the unique ovules of flowering plants. *Current Topics in Developmental Biology* **131**, 373–399.
- Girin T, Paicu T, Stephenson P, et al.** 2011. *INDEHISCENT* and *SPATULA* interact to specify carpel and valve margin tissue and thus promote seed dispersal in *Arabidopsis*. *The Plant Cell* **23**, 3641–3653.
- Gomez MD, Barro-Trastoy D, Escoms E, et al.** 2018. Gibberellins negatively modulate ovule number in plants. *Development* **145**, dev163865.
- Gómez MD, Fuster-Almunia C, Ocaña-Cuesta J, Alonso JM, Pérez-Amador MA.** 2019. *RGL2* controls flower development, ovule number and fertility in *Arabidopsis*. *Plant Science* **281**, 82–92.
- Gomez MD, Urbez C, Perez-Amador MA, Carbonell J.** 2011. Characterization of *constricted fruit* (*ctf*) mutant uncovers a role for *AtMYB117/LOF1* in ovule and fruit development in *Arabidopsis thaliana*. *PLoS ONE* **6**, e18760.
- Gonçalves B, Hasson A, Belcram K, et al.** 2015. A conserved role for *CUP-SHAPED COTYLEDON* genes during ovule development. *The Plant Journal* **83**, 732–742.
- Govaerts R.** 2001. How many species of seed plants are there? *Taxon* **50**, 1085–1090.
- Gremski K, Ditta G, Yanofsky MF.** 2007. The *HECATE* genes regulate female reproductive tract development in *Arabidopsis thaliana*. *Development* **134**, 3593–3601.

- Grini PE, Thorstensen T, Alm V, Vizcay-Barrena G, Windju SS, Jørstad TS, Wilson ZA, Aalen RB.** 2009. The ASH1 HOMOLOG 2 (ASHH2) histone H3 methyltransferase is required for ovule and anther development in *Arabidopsis*. *PLoS ONE* **4**, e7817.
- Gross T, Broholm S, Becker A.** 2018. CRABS CLAW acts as a bifunctional transcription factor in flower development. *Frontiers in Plant Science* **9**, 835.
- Grossniklaus U, Schneitz K.** 1998. The molecular and genetic basis of ovule and megagametophyte development. *Seminars in Cell & Developmental Biology* **9**, 227–238.
- Hedden P, Sponsel V.** 2015. A century of gibberellin research. *Journal of Plant Growth Regulation* **34**, 740–760.
- Heisler MG, Atkinson A, Bylstra YH, Walsh R, Smyth DR.** 2001. *SPATULA*, a gene that controls development of carpel margin tissues in *Arabidopsis*, encodes a bHLH protein. *Development* **128**, 1089–1098.
- Higuchi M, Pischke MS, Mähönen AP, et al.** 2004. *In planta* functions of the *Arabidopsis* cytokinin receptor family. *Proceedings of the National Academy of Sciences, USA* **101**, 8821–8826.
- Hill TA, Broadhvest J, Kuzoff RK, Gasser CS.** 2006. *Arabidopsis* *SHORT INTEGUMENTS 2* is a mitochondrial DAR GTPase. *Genetics* **174**, 707–718.
- Hu Y, Xie Q, Chua NH.** 2003. The *Arabidopsis* auxin-inducible gene *ARGOS* controls lateral organ size. *The Plant Cell* **15**, 1951–1961.
- Huang HY, Jiang WB, Hu YW, Wu P, Zhu JY, Liang WQ, Wang ZY, Lin WH.** 2013. BR signal influences *Arabidopsis* ovule and seed number through regulating related genes expression by BZR1. *Molecular Plant* **6**, 456–469.
- Iglesias FM, Cerdán PD.** 2016. Maintaining epigenetic inheritance during DNA replication in plants. *Frontiers in Plant Science* **7**, 38.
- Ishida T, Aida M, Takada S, Tasaka M.** 2000. Involvement of *CUP-SHAPED COTYLEDON* genes in gynoecium and ovule development in *Arabidopsis thaliana*. *Plant & Cell Physiology* **41**, 60–67.
- Kant S, Burch D, Badenhorst P, Palanisamy R, Mason J, Spangenberg G.** 2015. Regulated expression of a cytokinin biosynthesis gene IPT delays leaf senescence and improves yield under rainfed and irrigated conditions in canola (*Brassica napus* L.). *PLoS ONE* **10**, e0116349.
- Klucher KM, Chow H, Reiser L, Fischer RL.** 1996. The *AINTEGUMENTA* gene of *Arabidopsis* required for ovule and female gametophyte development is related to the floral homeotic gene *APETALA2*. *The Plant Cell* **8**, 137–153.
- Krizek B.** 2009. *AINTEGUMENTA* and *AINTEGUMENTA-LIKE6* act redundantly to regulate *Arabidopsis* floral growth and patterning. *Plant Physiology* **150**, 1916–1929.
- Larsson E, Franks RG, Sundberg E.** 2013. Auxin and the *Arabidopsis thaliana* gynoecium. *Journal of Experimental Botany* **64**, 2619–2627.
- Laufs P, Peaucelle A, Morin H, Traas J.** 2004. MicroRNA regulation of the CUC genes is required for boundary size control in *Arabidopsis* meristems. *Development* **131**, 4311–4322.
- Lee D-K, Geisler M, Springer PS.** 2009. *LATERAL ORGAN FUSION1* and *LATERAL ORGAN FUSION2* function in lateral organ separation and axillary meristem formation in *Arabidopsis*. *Development* **136**, 2423–2432.
- Liu P, Zhang C, Ma JQ, et al.** 2018. Genome-wide identification and expression profiling of cytokinin oxidase/dehydrogenase (*CKX*) genes reveal likely roles in pod development and stress responses in oilseed rape (*Brassica napus* L.). *Genes* **9**, 168.
- Liu Z, Franks RG, Klink VP.** 2000. Regulation of gynoecium marginal tissue formation by LEUNIG and AINTEGUMENTA. *The Plant Cell* **12**, 1879–1892.
- Mallory AC, Reinhart BJ, Jones-Rhoades MW, Tang G, Zamore PD, Barton MK, Bartel DP.** 2004. MicroRNA control of *PHABULOSA* in leaf development: importance of pairing to the microRNA 5' region. *The EMBO Journal* **23**, 3356–3364.
- Mantegazza O, Gregis V, Mendes MA, Morandini P, Alves-Ferreira M, Patreze CM, Nardeli SM, Kater MM, Colombo L.** 2014. Analysis of the *Arabidopsis* *REM* gene family predicts functions during flower development. *Annals of Botany* **114**, 1507–1515.
- Marhavý P, Bielach A, Abas L, et al.** 2011. Cytokinin modulates endocytic trafficking of PIN1 auxin efflux carrier to control plant organogenesis. *Developmental Cell* **21**, 796–804.
- Marsch-Martínez N, de Folter S.** 2016. Hormonal control of the development of the gynoecium. *Current Opinion in Plant Biology* **29**, 104–114.
- McClure BA, Guilfoyle T.** 1987. Characterization of a class of small auxin-inducible soybean polyadenylated RNAs. *Plant Molecular Biology* **9**, 611–623.
- Mizukami Y, Fischer RL.** 2000. Plant organ size control: *AINTEGUMENTA* regulates growth and cell numbers during organogenesis. *Proceedings of the National Academy of Sciences, USA* **97**, 942–947.
- Moubayidin L, Østergaard L.** 2014. Dynamic control of auxin distribution imposes a bilateral-to-radial symmetry switch during gynoecium development. *Current Biology* **24**, 2743–2748.
- Nahar MA, Ishida T, Smyth DR, Tasaka M, Aida M.** 2012. Interactions of *CUP-SHAPED COTYLEDON* and *SPATULA* genes control carpel margin development in *Arabidopsis thaliana*. *Plant & Cell Physiology* **53**, 1134–1143.
- Nemhauser JL, Feldman LJ, Zambryski PC.** 2000. Auxin and *ETTIN* in *Arabidopsis* gynoecium morphogenesis. *Development* **127**, 3877–3888.
- Nole-Wilson S, Azhakanandam S, Franks RG.** 2010a. Polar auxin transport together with *AINTEGUMENTA* and *REVOLUTA* coordinate early *Arabidopsis* gynoecium development. *Developmental Biology* **346**, 181–195.
- Nole-Wilson S, Rueschhoff EE, Bhatti H, Franks RG.** 2010b. Synergistic disruptions in *seuss cyp85A2* double mutants reveal a role for brassinolide synthesis during gynoecium and ovule development. *BMC Plant Biology* **10**, 198.
- Nomura T, Kushiro T, Yokota T, Kamiya Y, Bishop GJ, Yamaguchi S.** 2005. The last reaction producing brassinolide is catalyzed by cytochrome P-450s, CYP85A3 in tomato and CYP85A2 in *Arabidopsis*. *The Journal of Biological Chemistry* **280**, 17873–17879.
- Okada K, Ueda J, Komaki MK, Bell CJ, Shimura Y.** 1991. Requirement of the auxin polar transport system in early stages of *Arabidopsis* floral bud formation. *The Plant Cell* **3**, 677–684.
- Pagnussat GC, Yu HJ, Ngo QA, Rajani S, Mayalagu S, Johnson CS, Capron A, Xie LF, Ye D, Sundaresan V.** 2005. Genetic and molecular identification of genes required for female gametophyte development and function in *Arabidopsis*. *Development* **132**, 603–614.
- Pinto SC, Mendes MA, Coimbra S, Tucker MR.** 2019. Revisiting the female germline and its expanding toolbox. *Trends in Plant Science* **24**, 455–467.
- Reiser L, Fischer RL.** 1993. The ovule and the embryo sac. *The Plant Cell* **5**, 1291–1301.
- Reyes-Olalde JI, de Folter S.** 2019. Control of stem cell activity in the carpel margin meristem (CMM) in *Arabidopsis*. *Plant Reproduction* **32**, 123–136.
- Reyes-Olalde JI, Zuñiga-Mayo VM, Chávez Montes RA, Marsch-Martínez N, de Folter S.** 2013. Inside the gynoecium: at the carpel margin. *Trends in Plant Science* **18**, 644–655.
- Reyes-Olalde JI, Zuñiga-Mayo VM, Serwatowska J, et al.** 2017. The bHLH transcription factor *SPATULA* enables cytokinin signaling, and both activate auxin biosynthesis and transport genes at the medial domain of the gynoecium. *PLoS Genetics* **13**, e1006726.
- Rizza A, Jones AM.** 2019. The makings of a gradient: spatiotemporal distribution of gibberellins in plant development. *Current Opinion in Plant Biology* **47**, 9–15.
- Roeder AH, Yanofsky MF.** 2006. Fruit development in *Arabidopsis*. *The Arabidopsis Book* **4**, e0075.
- Ruzicka K, Simásková M, Duclercq J, Petrášek J, Zazimalová E, Simon S, Friml J, Van Montagu MC, Benková E.** 2009. Cytokinin regulates root meristem activity via modulation of the polar auxin transport. *Proceedings of the National Academy of Sciences, USA* **106**, 4284–4289.
- Schaller GE, Bishopp A, Kieber JJ.** 2015. The yin-yang of hormones: cytokinin and auxin interactions in plant development. *The Plant Cell* **27**, 44–63.
- Schlereth A, Möller B, Liu W, Kientz M, Flipse J, Rademacher EH, Schmid M, Jürgens G, Weijers D.** 2010. *MONOPTEROS* controls embryonic root initiation by regulating a mobile transcription factor. *Nature* **464**, 913–916.
- Schneitz K, Baker SC, Gasser CS, Redweik A.** 1998. Pattern formation and growth during floral organogenesis: *HUELLENLOS* and *AINTEGUMENTA* are required for the formation of the proximal region of the ovule primordium in *Arabidopsis thaliana*. *Development* **125**, 2555–2563.

- Schneitz K, Hulskamp M, Pruitt RE.** 1995. Wild-type ovule development in *Arabidopsis thaliana*: a light microscope study of cleared whole-mount tissue. *The Plant Journal* **7**, 731–749.
- Schuster C, Gaillochet C, Lohmann JU.** 2015. *Arabidopsis* *HECATE* genes function in phytohormone control during gynoecium development. *Development* **142**, 3343–3350.
- Sessions A, Nemhauser JL, McColl A, Roe JL, Feldmann KA, Zambryski PC.** 1997. *ETTIN* patterns the *Arabidopsis* floral meristem and reproductive organs. *Development* **124**, 4481–4491.
- Shah S, Karunarathna NL, Jung C, Emrani N.** 2018. An *APETALA1* ortholog affects plant architecture and seed yield component in oilseed rape (*Brassica napus* L.). *BMC Plant Biology* **18**, 380.
- Shi DQ, Yang WC.** 2011. Ovule development in *Arabidopsis*: progress and challenge. *Current Opinion in Plant Biology* **14**, 74–80.
- Shirley NJ, Aubert MK, Wilkinson LG, Bird DC, Lora J, Yang X, Tucker MR.** 2019. Translating auxin responses into ovules, seeds and yield: Insight from *Arabidopsis* and the cereals. *Journal of Integrative Plant Biology* **61**, 310–336.
- Šimášková M, O'Brien JA, Khan M, et al.** 2015. Cytokinin response factors regulate PIN-FORMED auxin transporters. *Nature Communications* **6**, 8717.
- Skinner DJ, Baker SC, Meister RJ, Broadhvest J, Schneitz K, Gasser CS.** 2001. The *Arabidopsis* *HUELLENLOS* gene, which is essential for normal ovule development, encodes a mitochondrial ribosomal protein. *The Plant Cell* **13**, 2719–2730.
- Skinner DJ, Hill TA, Gasser CS.** 2004. Regulation of ovule development. *The Plant Cell* **16 Suppl**, S32–S45.
- Smyth DR, Bowman JL, Meyerowitz EM.** 1990. Early flower development in *Arabidopsis*. *The Plant Cell* **2**, 755–767.
- Snowdon RJ, Lühs W, Friedt W.** 2007. Genome mapping and molecular breeding in plants. *Oilseeds* **6**, 54–56.
- Sohlberg JJ, Myrenäs M, Kuusk S, Lagercrantz U, Kowalczyk M, Sandberg G, Sundberg E.** 2006. *STY1* regulates auxin homeostasis and affects apical-basal patterning of the *Arabidopsis* gynoecium. *The Plant Journal* **47**, 112–123.
- Stinchcombe JR, Weinig C, Ungerer M, Olsen KM, Mays C, Halldorsdottir SS, Purugganan MD, Schmitt J.** 2004. A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the flowering time gene *FRIGIDA*. *Proceedings of the National Academy of Sciences, USA* **101**, 4712–4717.
- Sun Y, Fan XY, Cao DM, et al.** 2010. Integration of brassinosteroid signal transduction with the transcription network for plant growth regulation in *Arabidopsis*. *Developmental Cell* **19**, 765–777.
- van Mourik H, van Dijk ADJ, Stortenbeker N, Angenent GC, Bemer M.** 2017. Divergent regulation of *Arabidopsis* *SAUR* genes: a focus on the *SAUR10*-clade. *BMC Plant Biology* **17**, 245.
- Vanstraelen M, Benková E.** 2012. Hormonal interactions in the regulation of plant development. *Annual Review of Cell and Developmental Biology* **28**, 463–487.
- Wetzstein HY, Yi W, Porter JA, Ravid N.** 2013. Flower position and size impact ovule number per flower, fruitset, and fruit size in pomegranate. *Journal of the American Society for Horticultural Science* **138**, 159–166.
- Willis KJ** 2017. *State of the world's plants 2017*. Kew: Royal Botanic Gardens.
- Wynn AN, Seaman AA, Jones AL, Franks RG.** 2014. Novel functional roles for *PERIANTHIA* and *SEUSS* during floral organ identity specification, floral meristem termination, and gynoecial development. *Frontiers in Plant Science* **5**, 130.
- Xu L, Zhao Z, Dong A, Soubigou-Taconnat L, Renou JP, Steinmetz A, Shen WH.** 2008. Di- and tri- but not monomethylation on histone H3 lysine 36 marks active transcription of genes involved in flowering time regulation and other processes in *Arabidopsis thaliana*. *Molecular and Cellular Biology* **28**, 1348–1360.
- Yang Y, Wang Y, Zhan J, Shi J, Wang X, Liu G, Wang H.** 2017. Genetic and cytological analyses of the natural variation of seed number per pod in rapeseed (*Brassica napus* L.). *Frontiers in Plant Science* **8**, 1890.
- Yuan J, Kessler SA.** 2019. A genome-wide association study reveals a novel regulator of ovule number and fertility in *Arabidopsis thaliana*. *PLoS Genetics* **15**, e1007934.
- Zuñiga-Mayo VM, Baños-Bayardo CR, Díaz-Ramírez D, Marsch-Martínez N, De Folter S.** 2018. Conserved and novel responses to cytokinin treatments during flower and fruit development in *Brassica napus* and *Arabidopsis thaliana*. *Scientific Reports* **8**, 6836.
- Zuñiga-Mayo VM, Gómez-Felipe A, Herrera-Ubaldo H, de Folter S.** 2019. Gynoecium development: networks in *Arabidopsis* and beyond. *Journal of Experimental Botany* **70**, 1447–1460.