



RESEARCH PAPER

# Duplication and functional divergence of a calcium sensor in the Brassicaceae

Shea M. Monihan, Courtney A. Magness, Choong-Hwan Ryu, Michelle M. McMahon, Mark A. Beilstein<sup>id</sup> and Karen S. Schumaker<sup>\*</sup>,<sup>id</sup>

School of Plant Sciences, University of Arizona, 1140 E. South Campus Drive, Tucson, AZ 85721, USA

\* Correspondence: [schumake@email.arizona.edu](mailto:schumake@email.arizona.edu)

Received 5 August 2019; Editorial decision 13 January 2020; Accepted 27 January 2020

Editor: Timothy Colmer, University of Western Australia, Australia

## Abstract

The presence of varied numbers of *CALCINEURIN B-LIKE10* (*CBL10*) calcium sensor genes in species across the Brassicaceae and the demonstrated role of *CBL10* in salt tolerance in *Arabidopsis thaliana* and *Eutrema salsugineum* provided a unique opportunity to determine if *CBL10* function is modified in different species and linked to salt tolerance. Salinity effects on species growth and cross-species complementation were used to determine the extent of conservation and divergence of *CBL10* function in four species representing major lineages within the core Brassicaceae (*A. thaliana*, *E. salsugineum*, *Schrenkiella parvula*, and *Sisymbrium irio*) as well as the first diverging lineage (*Aethionema arabicum*). Evolutionary and functional analyses indicate that *CBL10* duplicated within expanded lineage II of the Brassicaceae and that, while portions of *CBL10* function are conserved across the family, there are species-specific variations in *CBL10* function. Paralogous *CBL10* genes within a species diverged in expression and function probably contributing to the maintenance of the duplicated gene pairs. Orthologous *CBL10* genes diverged in function in a species-specific manner, suggesting that functions arose post-speciation. Multiple *CBL10* genes and their functional divergence may have expanded calcium-mediated signaling responses and contributed to the ability of certain members of the Brassicaceae to maintain growth in salt-affected soils.

**Keywords:** Brassicaceae, *CALCINEURIN B-LIKE10*, calcium sensor, functional divergence, gene duplication, salt tolerance

## Introduction

Calcium has emerged as an essential component of many signaling pathways in plants, underlying growth and development by linking perception of physiological and environmental cues to cellular responses. Specificity in signaling is achieved, in part, through an array of proteins that perceive changes in cytosolic calcium levels (calcium sensors). The potential for different calcium sensors to contribute to functional specificity is found in their diverse temporal and spatial expression patterns, different affinities for calcium, and range of target proteins.

In *Arabidopsis thaliana* (Arabidopsis), *CALCINEURIN B-LIKE10* (*AtCBL10*) encodes a calcium sensor that enables

plants to grow in salt-affected soils by preventing the toxic accumulation of sodium ions in the cytosol (Quan *et al.*, 2007). Upon perception of changes in cytosolic calcium levels, *AtCBL10* interacts with the *SALT-OVERLY-SENSITIVE2* (*AtSOS2*) protein kinase to activate the *AtSOS1* plasma membrane sodium/proton exchanger which transports sodium out of the cell using the energy stored in the proton gradient (Quan *et al.*, 2007; Lin *et al.*, 2009). Mutation of *AtCBL10* results in hypersensitivity when plants are grown in the presence of even low levels of salt (Quan *et al.*, 2007). Two *CBL10* genes have been found in a salt-tolerant relative of Arabidopsis, *Eutrema*

*salsugineum* (*EsCBL10a* and *EsCBL10b*; Monihan *et al.*, 2019). A comparative study of *CBL10* function in Arabidopsis and *E. salsugineum* provided a starting point for understanding how duplication of a calcium sensor increased calcium-mediated signaling capacity in *E. salsugineum* and contributed to its salt tolerance. Reduced expression of *EsCBL10a* and *EsCBL10b* singly and in combination revealed that both genes function in the response of *E. salsugineum* to salt and probably have different functions (Monihan *et al.*, 2019). Both *EsCBL10* genes complement the *Atcbl10* salt-sensitive phenotype, indicating that there is conservation of *CBL10* function in the two species (Monihan *et al.*, 2019). When the genes were expressed in a salt-sensitive strain of *Saccharomyces cerevisiae* (yeast) with *SOS2* and *SOS1* from Arabidopsis or *E. salsugineum*, cells expressing *EsCBL10b* had the greatest growth on media with salt, while cells expressing *EsCBL10a* had the weakest growth relative to cells expressing *AtCBL10* (Monihan *et al.*, 2019). This result suggests that *EsCBL10b* strongly activates the SOS pathway while *EsCBL10a* does so only weakly. The different expression patterns of the *EsCBL10* genes provided further evidence that these two genes diverged in function. *EsCBL10b*, like *AtCBL10*, is expressed primarily in leaves, with very low expression in roots, while *EsCBL10a* is expressed in both leaves and roots (Monihan *et al.*, 2019). *AtSOS3*, another calcium sensor in the CALCINEURIN B-LIKE family in Arabidopsis (also known as *AtCBL4*), is expressed in roots and also functions in plant responses to salt (Liu and Zhu, 1998; Halfter *et al.*, 2000). The *AtSOS3* and *AtCBL10* proteins have non-overlapping roles during growth in the presence of salt (Quan *et al.*, 2007). *EsCBL10a*, but not *AtCBL10* or *EsCBL10b*, was able to complement the salt-sensitive *Atsos3* mutant phenotype, suggesting that *EsCBL10a* has a function not present in *AtCBL10* or *EsCBL10b* (Monihan *et al.*, 2019). Together these results demonstrated that the duplication of *CBL10* in *E. salsugineum* resulted in two calcium sensors with both shared and distinct activities expanding the response of *E. salsugineum* to salt.

Arabidopsis and *E. salsugineum* belong to the Brassicaceae, a diverse family of plants containing both economically and scientifically important species (Beilstein *et al.*, 2006; Yang *et al.*, 2013). The number of sequenced Brassicaceae genomes coupled with its well-established phylogeny led us to analyze the extent of conservation and divergence of *CBL10* gene function in both evolutionary and genomic contexts. Specifically we determined if: (i) other species in the Brassicaceae have multiple *CBL10* genes and if they are descendants of the same duplication event that resulted in two genes in *E. salsugineum*; (ii) the functions of the orthologous genes have been retained; and (iii) paralogous genes within selected species diverged in function. These analyses in combination with assays of plant growth in the presence of salt were used to understand how changes in *CBL10* function may have contributed to Brassicaceae adaptation to soil salinity. A phylogenetic analysis of *CBL10* revealed that the duplication resulting in two genes in *E. salsugineum* probably occurred in expanded lineage II (to which *E. salsugineum* belongs) of the family and resulted in several species retaining orthologs of *EsCBL10a* and *EsCBL10b*. *CBL10* expression and function were examined

in four species representing major lineages within the core Brassicaceae (*A. thaliana*, *E. salsugineum*, *Schrenkiella parvula*, and *Sisymbrium irio*) as well as the first diverging lineage (*Aethionema arabicum*) (Beilstein *et al.*, 2006). These analyses indicate that, while portions of *CBL10* function are conserved across the Brassicaceae, there are also variations in *CBL10* function that are specific to each species.

## Materials and methods

### Phylogenetic tree

Nucleotide sequences were identified using Basic Local Alignment Search Tool (BLAST) with the coding sequence of *A. thaliana CBL10* (*AtCBL10*; At4G33000.2) and a threshold E-score of the order of  $1 \times 10^{-15}$ . The identified genes were used in a reciprocal BLAST against the Arabidopsis genome. Only those sequences that identified *AtCBL10* as the closest homolog were used to generate the phylogeny. Sequences for *Arabidopsis lyrata* (Hu *et al.*, 2011), *Boechera stricta* (Windsor *et al.*, 2006), *Brassica nigra*, *Brassica oleracea* (Liu *et al.*, 2014), *Brassica rapa* (Wang *et al.*, 2011), *Camelina sativa* (Kagale *et al.*, 2014), *Capsella grandiflora* (Slotte *et al.*, 2013), *Capsella rubella* (Slotte *et al.*, 2013), *Carica papaya* (Ming *et al.*, 2008), *E. salsugineum* Shandong (Yang *et al.*, 2013), and *Raphanus raphanistrum* (Moghe *et al.*, 2014) were retrieved from the Phytozome database ([www.phytozome.net](http://www.phytozome.net)). Sequences for *S. parvula* (Dassanayake *et al.*, 2011) were retrieved from the NCBI genome database, while sequences for *A. arabicum*, *Neslia paniculata*, *Leavenworthia alabamica*, and *S. irio* were provided by Dr Stephen Wright (University of Toronto; Haudry *et al.*, 2013; Slotte *et al.*, 2013), and a sequence for *Tarenaya hassleriana* (Cheng *et al.*, 2013) was retrieved from the Comparative Genomics (CoGe) platform (Lyons and Freeling, 2008; Lyons *et al.*, 2008). Sequences for *CBL10* from the *E. salsugineum* accessions Cracker Creek, and Yukon were obtained by PCR using primers designed to amplify *CBL10* from the Shandong accession. To strengthen the inference of the root for the gene tree within the Brassicaceae, outgroup sequences for *Populus euphratica* (Ma *et al.*, 2013), *Prunus mume* (Zhang *et al.*, 2012), *Durio zibethinus* (Teh *et al.*, 2017), *Theobroma cacao* (Argout *et al.*, 2011), and *Vitis vinifera* (Jaillon *et al.*, 2007) were obtained from the GenBank nucleotide database using the *CBL10* sequence from *C. papaya* as a query for a BLAST search (all matched with E-scores of the order of  $1 \times 10^{-140}$ ).

Exon boundaries for the Arabidopsis, *E. salsugineum*, *S. parvula*, *S. irio*, and *A. arabicum CBL10* genes were determined by comparing the genomic sequences with cDNAs generated by PCR [*AtCBL10* (Quan *et al.*, 2007); *EsCBL10a* and *EsCBL10b* (Monihan *et al.*, 2019); *S. parvula*, *S. irio*, and *A. arabicum CBL10* genes, see cloning strategy in the complementation assay section below]. For all other *CBL10* genes, exon boundaries were estimated after multiple sequence alignments to the experimentally determined annotations. Extensive length heterogeneity, indels, and low-complexity sequences within each of the eight introns resulted in poorly aligned sequences. The *CBL10* coding sequences from the nine exon regions were concatenated, aligned in MUSCLE 3.8.31 (Edgar, 2004) using translated amino acids, and analyzed using IQ-TREE 1.6.7 (Nguyen *et al.*, 2015) with standard model selection. Two sequences from *L. alabamica* failed the composition test, indicating statistically significant differences in composition of these two sequences relative to the rest of the alignment. The model selected by the Bayesian information criterion and the corrected Akaike information criterion was TIM3+F+I+G4, which was used in the subsequent likelihood tree search with support for branches estimated by 100 bootstrap replicates.

### Synteny analysis

The genomic regions of Arabidopsis containing *CBL10* and *E. salsugineum* containing *EsCBL10a* and *EsCBL10b* were compared with 15 genomes (*A. arabicum*, *A. lyrata*, *B. stricta*, *B. nigra*, *B. oleracea*, *B. rapa*, *C. sativa*, *C. grandiflora*, *C. rubella*, *C. papaya*, *L. alabamica*, *R. raphanistrum*, *S. parvula*, *S. irio*, and *T. hassleriana*) to identify collinear regions using the SynFind

feature on CoGe (Lyons and Freeling, 2008; Lyons *et al.*, 2008). The genes located on the two identified collinear regions of the Arabidopsis genome were used as a query for a BLAST search to detect putative homologs in the genomes of the other species (Altschul *et al.*, 1990). The species tree shown in the synteny analysis is based on Beilstein *et al.* (2010) and Yang *et al.* (2013).

#### Identification of transposable elements

RepeatMasker 4.0.6 was used to detect putative transposable elements in flanking regions of *CBL10/CBL10a* and *CBL10b* in Arabidopsis, *E. salsugineum*, *S. irio*, and *S. parvula* (A.F.A. Smit, R. Hubley, and P. Green; RepeatMasker at <http://repeatmasker.org>).

#### Plant growth

To determine the salt tolerance of each species, seeds of Arabidopsis, *E. salsugineum* Shangdong, *S. parvula*, *S. irio*, and *A. arabicum* were sown on SunGro Sunshine LC1 soil mix (SunGrow Horticulture) and stratified for 4 d at 4 °C in the dark to break dormancy. After cold treatment, plants were transferred to a growth chamber at 21 °C under a 16 h light/8 h dark photoperiod with light provided by Phillips F32T8/TL841 bulbs (135  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) and watered every 3 d with 0.25 $\times$  Hoagland's solution (Hoagland and Arnon, 1938) with cobalt chloride in place of cobalt nitrate. When true leaves developed, plants were treated with increasing salt (sodium chloride, NaCl) in 50 mM increments every 3 d until the indicated, final concentration was reached. Three weeks after the start of treatment, photographs were taken and the fresh weight of aerial tissue was recorded. Comparisons of growth were performed by analyzing the ratio of species growth in the absence and presence of salt at each NaCl concentration using Friedman's non-parametric, two-way ANOVA and Tukey's honestly significant difference (HSD) tests. Statistical significance was assigned at  $P \leq 0.05$ .

#### Complementation assays

*CBL10* gene function was analyzed by expressing each gene in the *Atcbl10* T-DNA insertion mutant (Arabidopsis Biological Resource Center, SALK\_056042; Monihan *et al.*, 2016) and the *Atsos3* ethyl methanesulfonate mutant (Dr Jian-Kang Zhu; Purdue University; Liu and Zhu, 1997). Full-length coding sequences without a stop codon were amplified from cDNA and cloned into pGEM-T Easy (Promega). All PCR amplifications were performed using Phusion High-Fidelity DNA Polymerase (ThermoFisher Scientific); primer sequences are provided in Supplementary Table S1 at JXB online. Because of the high degree of similarity between the *S. parvula* transcripts, forward primers were designed to anneal to the 5'-untranslated region (UTR) to enrich for a specific *CBL10* gene; primer sequences are provided in Supplementary Table S1. The *S. parvula* PCR products were then used as a template in a second reaction with primers that anneal to the coding sequence for cloning into pGEM-T Easy (Promega). All genes were digested with *XhoI* and *BamHI*, and subcloned into the corresponding site of the plant binary vector pEZT-NL (Drs Sean Cutler and David W. Ehrhardt; Carnegie Institution of Washington) downstream of the *Cauliflower mosaic virus* (CaMV) 35S promoter. *Agrobacterium tumefaciens* LBA4404 containing the binary vector was used to transform *Atcbl10* and *Atsos3* via the floral dip method (Clough and Bent, 1998). T<sub>1</sub> seed was germinated on soil for 1.5 weeks and then sprayed three times with 100 mg l<sup>-1</sup> Basta (Rely 200 Herbicide; Bayer Crop Science) at 3 d intervals. T<sub>1</sub> lines with antibiotic resistance were subsequently transferred to pots and grown to collect T<sub>2</sub> seed. Single insertion lines were identified by screening T<sub>2</sub> seed on 0.5 $\times$  MS medium [Murashige and Skoog medium; PhytoTechnology Laboratories containing 2.5 mM MES, 2% sucrose (w/v), and 1% agar (w/v) (A8678; Sigma), pH 5.7 (adjusted with potassium hydroxide)] and 7.5 mg l<sup>-1</sup> glufosinate ammonium (Santa Cruz Biotechnology). Lines with 75% resistance were selected. Homozygous lines were identified by screening T<sub>3</sub> seed on MS plates with glufosinate ammonium and selecting lines with 100% resistance. Gene-specific primers were used to confirm the identity and expression of all transgenes (Supplementary Fig. S1).

To monitor seedling growth, transgenic Arabidopsis seeds were sown on 0.5 $\times$  MS medium. Plates with seed were incubated at 4 °C in the dark for 2 d to break dormancy and transferred to a growth chamber at 21 °C under a 16 h light/8 h dark photoperiod with light provided by Phillips F32T8/TL841 bulbs (135  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ). For salt assays, seeds were germinated on medium without NaCl for 4 d, after which seedlings were transferred to medium without or with the indicated concentration of NaCl. NaCl concentrations were chosen for maximal differences in growth between the wild type and mutants. After 10 d of treatment, photographs were taken and seedling fresh weight was measured to quantify growth.

#### Transcript analysis

To determine expression patterns of the *CBL10* genes, leaves and roots of 11-day-old seedlings grown on 0.25 $\times$  MS medium were collected. RNA was isolated using the NucleoSpin RNA extraction kit (Macherey-Nagel) and used to synthesize cDNA (M-MLV Reverse Transcriptase; Promega). All PCRs were performed using recombinant Taq polymerase (Invitrogen) and were analyzed and compared in the linear range of the amplification (e.g. 23 cycles for *Actin* and 28 cycles for *CBL10*); primer sequences are provided in Supplementary Table S1.

#### Yeast salt screens

To monitor the ability of the *CBL10* proteins to activate the SOS pathway, *Saccharomyces cerevisiae* strain AXT3K (*ena1::HIS3::ena4*, *nha1::LEU2*, and *nhx1::KanMX4*; Quintero *et al.*, 2002) containing the pYPGE15 plasmid with *AtSOS1* (Jarvis *et al.*, 2014) and the pFL32T plasmid with *AtSOS2* and *AtSOS3* was modified to express the *CBL10* genes. Plasmids containing *AtCBL10*, *EsCBL10a*, *EsCBL10b*, or without a *CBL10* gene were generated as previously described (Monihan *et al.*, 2019). The full-length coding sequences of *SpCBL10a*, *SpCBL10b-1*, *SpCBL10b-2*, *SiCBL10a*, *SiCBL10b*, and *AaCBL10* were amplified from cDNA (Phusion High-Fidelity DNA polymerase; ThermoFisher Scientific; primer sequences are provided in Supplementary Table S1), cloned into pGEM-T Easy (Promega), digested with *XhoI* and *NotI*, and subcloned into the corresponding site of the pDR195 vector (Dr Alonso Rodriguez-Navarro; Rentsch *et al.*, 1995). Because of the high degree of similarity between the *S. parvula* transcripts, forward primers were designed to anneal to the 5'-UTR to enrich for a specific *CBL10* gene (Supplementary Table S1). The *S. parvula* PCR products were then used as a template in a second reaction with primers that anneal to the coding sequence for cloning into pGEM-T Easy (Promega). The pDR195 plasmids were digested with *AgeI* and *NotI*, and a fragment containing the *CBL10* gene was subcloned into the corresponding site of the pFL32T plasmid in place of *AtSOS3* to be expressed with *AtSOS2*. Transformed AXT3K cells were selected on synthetic dropout medium lacking both uracil and tryptophan (Clontech/TaKaRa) containing yeast nitrogen base without amino acids (VWR). Salt assays were carried out in alkali cation-free medium (AP; Rodriguez-Navarro and Ramos, 1984) containing 1 mM KCl with the designated concentrations of NaCl and cultured at 30 °C for 4 d.

#### Yeast two-hybrid assays

To determine if *EsCBL10a* and the *SpCBL10b* proteins might complement *Atsos3* by interaction with a similar CBL-interacting protein kinase (CIPK), yeast-two hybrid assays were performed between both *SpCBL10b* proteins and the four *EsCBL10a*-interacting *AtCIPK* proteins. Cloning of *AtCBL10*, *EsCBL10a*, *EsCBL10b*, and *AtCIPK* genes was described previously (Monihan *et al.*, 2019). *SpCBL10b-1* and *SpCBL10b-2* genes were PCR amplified using Phusion High-Fidelity DNA Polymerase (ThermoFisher Scientific); primer sequences are provided in Supplementary Table S1. The PCR products were digested with *EcoRI* and *BamHI*, and cloned into the corresponding site of the pGADT7 and pGBKT7 vectors (Clontech/TaKaRa) which allows for expression of the gene as a fusion protein with the GAL4 DNA activation domain (AD) or the GAL4 DNA-binding domain (BD), respectively. The pGADT7 clones were transformed into *S. cerevisiae* strain Y2HGOLD



(Clontech/TaKaRa), while the pGBKT7 clones were transformed into *S. cerevisiae* strain Y187 (Clontech/TaKaRa). Yeast were mated and grown on synthetic defined medium (SD) minus leucine and tryptophan (SD-LW) to select for diploid yeast expressing both constructs. To determine interaction, serial dilutions of yeast colonies were grown on SD-LW and without histidine (H), and incubated for 5 d. Interaction is shown in only one orientation because the SpCBL10b proteins fused to the GAL4 BD self-activated, causing strains expressing only these fusion proteins to grow on all selection media, masking any interaction with the CIPK proteins.

### Statistical analysis

To determine significant differences in growth, experiments were organized and analyzed as a randomized complete block design with genotypes and salt concentrations as treatments, and individual experiments as replicates. Treatment effects were assessed using a full-factorial mixed-model ANOVA in JMP, Version 11 (SAS Institute; 1989–2007). In these analyses, treatments were considered fixed effects and replicates random effects. The normality of the distributions of all dependent variables was analyzed by examining a plot of the residuals from a full-factorial ANOVA of untransformed data. A Shapiro–Wilk test (Shapiro and Wilk, 1965) was performed to assess normality, and Bartlett's (Bartlett, 1937) and Levene's (Levene, 1960) tests were performed to evaluate the homogeneity of variance. Based on the pattern of distribution and the results of these tests, a non-parametric approach was used to analyze the data throughout. Data were rank transformed using Microsoft Excel (function: RANK) followed by an ANOVA and Tukey's HSD test for multiple comparisons of means (Conover and Iman, 1981). The HSD values from rank-based ANOVA were then applied to the actual means for each measurement (i.e. not the ranks used in ANOVA). Statistical significance was assigned at  $P \leq 0.05$  throughout, and all tests of significance were two sided.

## Results


### *CBL10 duplicated within expanded lineage II of the Brassicaceae*

Previous studies have shown that there are two CALCINEURIN B-LIKE10 (CBL10) calcium sensors in *E. salsugineum*, a salt-tolerant relative of Arabidopsis. Reduced expression of the duplicated *E. salsugineum* CBL10 genes demonstrated that both genes function in the response of the plant to salt (Monihan et al., 2019). Cross-species complementation assays demonstrated that the two *E. salsugineum* CBL10 proteins have both shared and distinct activities (Monihan et al., 2019), suggesting that duplication of CBL10 increased calcium-mediated signaling capacity in *E. salsugineum*.

To understand CBL10's role in calcium-mediated signaling in an evolutionary context and determine when in the evolutionary history of the Brassicaceae the CBL10 duplication arose, CBL10 sequences were identified from species within and outside of the Brassicaceae and a phylogeny was inferred based on maximum likelihood. Of eight species examined within lineage I of the Brassicaceae, seven have a single CBL10 gene: *A. thaliana*, *A. lyrata*, *B. stricta*, *C. rubella*, *C. grandiflora*, *C. sativa*, and *N. paniculata* (Fig. 1). *Leavenworthia alabamica* was the only lineage I species examined, with multiple CBL10 genes probably due to a whole-genome triplication event; however, only two CBL10 genes were identified (Fig. 1; Haudry et al., 2013). CBL10 sequences from *B. rapa*, *B. nigra*, *B. oleracea*, *R. raphanistrum*, *S. irio*, and *S. parvula* (lineage II species) were compared with the CBL10 genes in *E. salsugineum* (which belongs to expanded lineage II; Beilstein et al., 2006;

Yang et al., 2013). *Sisymbrium irio*, *S. parvula*, and *E. salsugineum* have both CBL10a and CBL10b paralogs (Fig. 1), with an additional CBL10b gene in *S. parvula* due to a tandem duplication (Fig. 2; Dassanayake et al., 2011; Oh et al., 2014). *Raphanus raphanistrum* and all three Brassica species have three CBL10a genes, consistent with their whole-genome triplication events (Fig. 1; Lagercrantz and Lydiate, 1996; Moghe et al., 2014), but no CBL10b genes. *Aethionema arabicum*, which is a member of the first diverging lineage of the Brassicaceae, contains only a single CBL10 gene (Fig. 1). Outside of the Brassicaceae, *T. hassleriana* and *C. papaya*, members of the Cleomaceae and Caricaceae, respectively, both have a single CBL10 gene (Fig. 1). Sequences from *P. euphratica*, *P. mume*, *D. zibethinus*, *T. cacao*, and *V. vinifera* were included to provide additional data for the tree which was rooted on the *V. vinifera* sequence (data not shown). The tree is consistent with a single duplication in the ancestor of lineage II, leading to all known CBL10b genes which form a strongly supported clade (Fig. 1). The Brassica and *R. raphanistrum* CBL10 genes descended from a triplication in the ancestor of just those species, not from the duplication that led to CBL10b, indicating that the ancestor of the Brassica species and *R. raphanistrum* lost CBL10b. *Sisymbrium irio* CBL10a was placed with weak support among the Brassica species, suggesting that additional duplications (or other phenomena such as lineage sorting) must have occurred. While it is not possible to unequivocally determine the timing of the duplication due to low branch support, the presence of both CBL10a and CBL10b paralogs in multiple members of expanded lineage II and the absence of the paralogs in members of lineage I, *A. arabicum*, *T. hassleriana*, and *C. papaya*, suggest that the duplication of CBL10 occurred within expanded lineage II of the Brassicaceae.

The region surrounding the CBL10 sequence was compared to provide additional information about the origin of the genes. *EsCBL10a* is syntenic with CBL10 genes from Arabidopsis, *A. lyrata*, both *Capsella* species, *B. stricta*, *L. alabamica*, *C. sativa*, *T. hassleriana*, and *C. papaya*, and the CBL10a genes from all Brassica species, *S. parvula*, and *S. irio* (select species shown in Fig. 2A). Synteny in *N. paniculata* and *R. raphanistrum* was difficult to determine because the CBL10 sequences were found on short scaffolds. The fact that all CBL10 genes from the Brassica species and from *L. alabamica* share synteny with *EsCBL10a* (data not shown) further supports the conclusion that these CBL10 genes arose through independent whole-genome triplication events (Lagercrantz and Lydiate, 1996; Haudry et al., 2013). The CBL10 gene from *A. arabicum* was found on a short scaffold with insufficient sequence to assess synteny (scaffold 3407, Fig. 2A). A second scaffold in *A. arabicum* sharing synteny with Arabidopsis was detected, but no CBL10 gene was identified (scaffold 3487, Fig. 2A). The genomic region syntenic to *EsCBL10b* was identified in all species; however, a CBL10b gene was only detected in *E. salsugineum*, *S. parvula*, and *S. irio* (Fig. 2B). Numerous putative transposable elements were found surrounding the CBL10b genes, suggesting that the duplication of CBL10 might have been mediated by transposons (Supplementary Fig. S2). However, it was not possible to identify a single transposon associated with all three CBL10b




**Fig. 1.** The duplication of *CBL10* probably occurred within expanded lineage II in the Brassicaceae. Exons from *CBL10* nucleotide sequences were aligned and analyzed using maximum likelihood. Sequences from *P. euphratica*, *P. mume*, *D. zibethinus*, *T. cacao*, and *V. vinifera* were included to provide additional data for the tree which was rooted on the *V. vinifera* sequence. Circles above branches represent the percentage of 100 bootstrap replicates that support the topology; closed circles, 90–100; open circles, 70–89. *CBL10* genes from five species were chosen for further analysis and color coded. Green, *CBL10* genes from two species with a single gene; red, *CBL10a* genes from three species with multiple genes; blue, *CBL10b* genes from three species with multiple genes. Species within lineage I and expanded lineage II are indicated.

genes. Taken together, these results suggest that transposons may have mediated the duplication of *CBL10* and that the insertion of *CBL10b* into a different chromosomal position took place before the divergence of expanded lineage II species within the Brassicaceae.

#### Species in expanded lineage II are salt tolerant

As a first step in linking *CBL10* function to salt tolerance within the Brassicaceae, five species were selected for further analyses. In addition to *Arabidopsis* and *E. salsugineum*, *S. parvula* and *S. irio* were chosen because of their close relationship to *E. salsugineum* and the presence of multiple *CBL10* genes in their genomes (Beilstein et al., 2006; Fig. 3). Due to its position as a member of the first diverging group within the Brassicaceae and its more distant relationship with *E. salsugineum*, *A. arabicum* was also included in the analysis

(Beilstein et al., 2006; Fig. 3). The salt tolerance (FW of each species treated with increasing concentrations of salt) was measured. *Schrenkiella parvula*, *E. salsugineum*, and *S. irio* all maintained growth in concentrations of salt up to 300 mM, while growth of *Arabidopsis* and *A. arabicum* quickly decreased at concentrations as low as 100 mM (Fig. 3). Ratios of species growth in the absence and presence of salt at each salt concentration indicated that the salt tolerance of *S. parvula*, *E. salsugineum*, and *S. irio* was similar and distinct from the salt tolerance of *Arabidopsis* and *A. arabicum* which were similar (Supplementary Table S2). Because the increased salt tolerance of the expanded lineage II species analyzed might be due to an increase in the number of *CBL10* genes and/or a divergence in function of those genes, four assays (cross-species complementation of the *cb10* and *sos3* mutants in *Arabidopsis*, expression analysis, and SOS pathway activation) were used to compare the activities of the genes.




**Fig. 2.** The genomic positions of *AtCBL10* and the *CBL10b* genes differ. The organismal phylogeny is shown with the species name and the chromosome, contig, or scaffold on which *CBL10* was identified. Stars, short scaffolds; horizontal line, genomic region (not drawn to scale); pentagons, genes. In Arabidopsis, *CBL10* is black and the genes on either side were identified and assigned a color. In the other species, *CBL10* genes were identified and the surrounding genes were colored based on ontology with genes from Arabidopsis. Vertical lines, presence of genes not syntenic to those in Arabidopsis. Tandem duplicates of the flanking genes were collapsed to one gene/pentagon to simplify the figure. (a) Regions syntenic to *AtCBL10* and *EsCBL10a*. (b) Regions syntenic to *EsCBL10b*.

### *CBL10* function is conserved in species across the Brassicaceae

The *CBL10* phylogenetic tree suggests that all the genes identified are homologs of *AtCBL10* and may share activities. To determine if there is conservation of *CBL10* function throughout the Brassicaceae, the *CBL10* genes from *S. parvula*, *S. irio*, and *A. arabicum* were expressed in the Arabidopsis *cb110*

mutant (*Atcb110*). Our studies have shown that introns are necessary for full expression and function of *AtCBL10* when using the native promoter, but can lead to alternative splicing. To avoid alternative splicing of the *CBL10* genes under study and to ensure strong expression of their coding sequences, the CaMV 35S constitutive promoter was used for these cross-species complementation assays. The salt tolerance of four independently transformed, single insertion, homozygous






**Fig. 3.** Species in expanded lineage II in the Brassicaceae are salt tolerant. Seeds from *Arabidopsis*, *E. salsugineum*, *S. parvula*, *S. irio*, and *A. arabicum* were germinated and grown on soil for 1 week and then treated with increasing NaCl in 50 mM increments every 3 d until the indicated final concentration was reached. Three weeks after the start of treatment, aerial portions of the plants were harvested, photographed, and weighed. Scale bar=1 cm for all images. The average fresh weight was graphed and  $\pm$ SE is shown. One representative image of seven experiments is shown.

lines was assessed for each gene. All of the tested *CBL10* genes complemented the *Atcbl10* salt-sensitive phenotype, indicating that there is conservation of *CBL10* function throughout the Brassicaceae (Fig. 4).

#### *CBL10* function diverged in a species-specific manner in the Brassicaceae

It has previously been shown that the *CBL10* genes in *E. salsugineum* have different expression patterns. Like *AtCBL10*, *EsCBL10b* is expressed predominately in aerial tissue, whereas *EsCBL10a* is expressed throughout *E. salsugineum* (Monihan et al., 2019). To determine if *CBL10* expression correlates

with protein activity across the Brassicaceae, RNA was isolated from shoots and roots of each species and transcript accumulation was monitored. The exon-intron structure of the *CBL10* transcripts was very similar, and protein domains known to be important for *AtCBL10* function were present in all transcripts (Fig. 5). Expression patterns correlated with the phylogenetic relationship of the genes in lineage I and expanded lineage II species; *CBL10b* transcripts were high in leaves (similar to *AtCBL10*) while *CBL10a* transcripts accumulated in both leaves and roots (Fig. 5). The expression pattern of the *CBL10* gene from *A. arabicum* was opposite to what was seen for the *CBL10b* and *AtCBL10* genes; the transcript was high in roots (Fig. 5).




**Fig. 4.** *CBL10* genes from all Brassicaceae species complement the *Atcbl10* salt-sensitive phenotype. *CBL10* genes from *Arabidopsis* (*AtCBL10*, *At10*), *E. salsugineum* (*EsCBL10a*, *Es10a*; and *EsCBL10b*, *Es10b*), *S. parvula* (*SpCBL10a*, *Sp10a*; *SpCBL10b-1*, *Sp10b-1*; and *SpCBL10b-2*, *Sp10b-2*), *S. irio* (*SiCBL10a*, *Si10a*; and *SiCBL10b*, *Si10b*), and *A. arabicum* (*AaCBL10*, *Aa10*) were expressed in the *Atcbl10* mutant, and growth in the absence (control) and presence of salt (125 mM NaCl) was monitored. For salt assays, seeds were germinated on medium without NaCl for 4 d, after which seedlings were transferred to medium without or with the indicated concentration of NaCl. After 10 d of treatment, photographs were taken and total seedling fresh weight was measured. (A) Photographs of the wild type (WT), *Atcbl10*, and *Atcbl10* expressing each *CBL10* gene (*Atcbl10;CBL10*). The scale bar (1 cm, upper right panel) shows the magnification for all images. (B) Total seedling fresh weight was measured to quantify growth and is presented as a percentage of control. Data are means  $\pm$ SE of at least 24 seedlings per genotype grown in three independent experiments. \*Complementation of the *Atcbl10* salt-sensitive phenotype (Tukey–Kramer HSD,  $P < 0.05$ ).

*AtCBL10* has been shown to activate the SOS pathway which functions to prevent the toxic accumulation of sodium in the cytoplasm (Quan *et al.*, 2007; Lin *et al.*, 2009). In *E. salsugineum*, differences in *CBL10* activation of the pathway

were observed; *EsCBL10b* strongly activated the *Arabidopsis* and *E. salsugineum* SOS pathways, while *EsCBL10a* showed only weak activation (Monihan *et al.*, 2019). To determine if the *CBL10* proteins from *S. irio*, *S. parvula*, and *A. arabicum*





**Fig. 5.** Expression of the *CBL10a* and *CBL10b* genes differs. (a) Transcript structure of the *CBL10* genes from *Arabidopsis* (*AtCBL10*), *E. salisugineum* (*EsCBL10a* and *EsCBL10b*), *S. parvula* (*SpCBL10a*, *SpCBL10b-1*, and *SpCBL10b-2*), *S. irio* (*SiCBL10a* and *SiCBL10b*), and *A. arabicum* (*AaCBL10*). Black boxes, exons (size indicated below in bp); solid lines, introns (not drawn to scale); dotted lines, untranslated regions; gray boxes, primer annealing sites. Protein domains important for the function of *AtCBL10* are indicated above the *AtCBL10* transcript. HD, hydrophobic domain; Ca, calcium-binding domains; S, serine phosphorylation site. (b) *CBL10* transcript accumulation in leaves (L) and roots (R) of 11-day-old seedlings grown on 0.25× MS medium. *ACTIN2* (*ACT2*), loading control. One representative image of three replicates is shown.

function in the SOS pathway, each protein was expressed in a salt-sensitive strain of yeast (AXT3K) along with the *Arabidopsis* SOS2 protein kinase (*AtSOS2*) and SOS1  $\text{Na}^+/\text{H}^+$  exchanger (*AtSOS1*), and growth in the presence of salt was assessed as an indication of pathway activity. All *CBL10* proteins activated the SOS pathway, but differences in the level of activation were observed. *EsCBL10b* had the greatest activity followed by the *SpCBL10b* and *SpCBL10a* proteins which all had greater activity than *AtCBL10* (Fig. 6). The activities of *SiCBL10a* and *SiCBL10b* were similar to that of *AtCBL10*, while *EsCBL10a* and *AaCBL10* had the weakest activity (Fig. 6).

*EsCBL10a*, but not *AtCBL10* or *EsCBL10b*, can complement the salt-sensitive phenotype of the *Arabidopsis* *sos3* mutant (*Atsos3*), suggesting that *EsCBL10a* has a distinct function (Monihan et al., 2019). To determine when this function arose within the Brassicaceae, the *S. irio*, *S. parvula*, and *A. arabicum* *CBL10* genes were expressed in *Atsos3* downstream of the CaMV 35S promoter and the salt tolerance of five independently transformed, single insertion, homozygous lines was determined. Four of the nine genes examined complemented *Atsos3*: *EsCBL10a*, *SpCBL10b-1*, *SpCBL10b-2*, and *SiCBL10b* (Fig. 7). These results indicate that at least one gene from each expanded lineage II species can perform a function that complements the *Atsos3* salt-sensitive phenotype.


The ability of *EsCBL10a* to complement *Atsos3* and its weak ability to activate *Arabidopsis* and *E. salisugineum* SOS2 and

SOS1 in yeast suggested that it might perform its functions with a different kinase (Monihan et al., 2019). A yeast two-hybrid screen was performed with 26 kinases from the CIPK family to which *AtSOS2* belongs. Four CIPKs (*AtCIPK13*, *AtCIPK16*, *AtCIPK6*, and *AtCIPK18*) were identified as interacting specifically with *EsCBL10a* but not *AtCBL10* or *EsCBL10b* (Monihan et al., 2019). To determine if the *CBL10b* proteins from *S. parvula* might complement *Atsos3* through a mechanism similar to *EsCBL10a*, interaction was tested between the *SpCBL10b* proteins and the four *EsCBL10a*-interacting CIPKs. *SpCBL10b-1* interacted with two of the kinases, *AtCIPK6* and *AtCIPK18*, while *SpCBL10b-2* only interacted with *AtCIPK18* (Fig. 8). In the opposite orientation, the *S. parvula* proteins self-activated, masking any interaction with the CIPK proteins, so interaction is only shown for the CIPK proteins fused to the GAL4 BD and the *CBL10* proteins fused to the GAL4 AD.

## Discussion

### Cross-species analysis sheds light on the evolution of *CBL10* duplication in the Brassicaceae

All of the *CBL10* genes examined complemented the *Atcbl10* salt-sensitive phenotype, indicating that at least a portion of *CBL10* function is conserved throughout the Brassicaceae (Figs 4, 9). This is consistent with studies of *CBL10* genes in



**Fig. 6.** SOS pathway activation is greatest with EsCBL10b. A salt-sensitive strain of *S. cerevisiae* (AXT3K,  $\Delta ena1-4\Delta nha1\Delta nhx1$ ) was transformed with SOS1 and SOS2 from Arabidopsis in combination with *CBL10* from Arabidopsis (*AtCBL10*, *At10*), *E. salicaria* (*EsCBL10a*, *Es10a*; and *EsCBL10b*, *Es10b*), *S. parvula* (*SpCBL10a*, *Sp10a*; *SpCBL10b-1*, *Sp10b-1*; and *SpCBL10b-2*, *Sp10b-2*), *S. irio* (*SiCBL10a*, *Si10a*; and *SiCBL10b*, *Si10b*), and *A. arabicum* (*AaCBL10*, *Aa10*). Serial decimal dilutions of yeast cells were spotted onto control medium or medium containing 125 mM NaCl. Two independently transformed colonies were assayed in three biological replicates; one representative image is shown.

*Poplar trichocarpa* and *Solanum lycopersicum* which were also able to complement *Atcbl10*, indicating that some of *CBL10* function is conserved outside of the Brassicaceae (Tang *et al.*, 2014; Egea *et al.*, 2018). However, within the Brassicaceae, species-specific differences in *CBL10* expression and function were detected.

The absence of the *CBL10a* and *CBL10b* paralogs in members of lineage I, *A. arabicum*, *T. hassleriana*, and *C. papaya* in combination with the presence of *CBL10a* and *CBL10b* paralogs in multiple members of lineage II suggest that the duplication of *CBL10* occurred within expanded lineage II of the Brassicaceae (Fig. 1). Lack of variation in the C-termini of the *CBL10* nucleotide sequences (containing the four highly conserved EF-hand calcium-binding domains) reduced phylogenetic resolution, precluding assignment of the duplication to a specific branch of the tree.

At least one gene from each species within expanded lineage II has the ability to complement the *Atsos3* salt-sensitive phenotype (Figs 7, 9). Whether the ability to complement *Atsos3* evolved prior to or after subsequent speciation in the expanded lineage II species sampled remains less clear; however, there is support for the changes occurring after speciation. The ability of EsCBL10a to complement *Atsos3* resides in the hydrophobic domain (Monihan *et al.*, 2019). Conservation of amino acids in this domain in proteins that do not complement (*AtCBL10*, *EsCBL10b*, *SpCBL10a*, *SiCBL10a*, and *AaCBL10*) and variation in proteins that do (*EsCBL10a*, *SpCBL10b-1*, *SpCBL10b-2*, and *SiCBL10b*) suggests that sequences changed post-*CBL10* duplication and after speciation (Supplementary Fig. S3). Additional studies will be required to determine if the

other *CBL10b* proteins that complement *Atsos3* do so because of changes in this or another region of the protein.

With an increase in sequenced genomes and tools to analyze those genomes, questions have emerged regarding how genomic context influences which gene changes function after a duplication event. To determine if the change in function is equally likely in both genes or more likely in the gene inserted into a new genomic position, the ratio of non-synonymous (dN) to synonymous (dS) substitutions (dN/dS) is often calculated. Studies have generally concluded that the duplicated gene in the new genomic position often has the faster rate of change and, as a result, is the one most likely to change function (Han *et al.*, 2009; Dewey, 2011; Pegueroles *et al.*, 2013; Rosello and Kondrashov, 2014). In keeping with these findings, the *CBL10* genes in *S. parvula* and *S. irio* inserted into a new genomic position and acquired a function that complements *Atsos3*. However, *E. salicaria* appears to be an exception to this generalization. *EsCBL10a*, which remained in the original genomic position, acquired the ability to complement *Atsos3*, while *EsCBL10b*, which inserted into a new genomic position, never acquired the function or acquired the function and subsequently lost it. In addition to influencing protein function, duplication and insertion of a gene into a new genomic context can lead to regulation by different *cis*-acting elements, suggesting that the gene that changes genomic position is more likely to have a different expression pattern (Flagel and Wendel, 2009; Pegueroles *et al.*, 2013). Our current data do not support this model because the *CBL10b* genes inserted into a new genomic position yet they share a similar expression pattern with *AtCBL10*, while the *CBL10a* genes, which remained in the original genomic position, appear to have expanded expression into roots. Analysis of expression patterns from additional species will be needed to establish the major expression pattern.


#### Cross-species analysis sheds light on the molecular mechanism underlying *CBL10* duplication in the Brassicaceae

Transposable elements have been found to replicate and integrate into genomes, leading to duplication of genes (Xiao *et al.*, 2008; Flagel and Wendel, 2009; Panchy *et al.*, 2016). Because numerous transposable elements were found surrounding the *CBL10b* genes in the species studied (Supplementary Fig. S2), replicative transposition by transposable elements is a likely mechanism underlying the duplication of *CBL10*. *CBL10* duplication as a result of a polyploidization event is unlikely because no whole-genome duplication event is known to have occurred at the base of expanded lineage II and because, in the *CBL10*-containing regions of the genomes of species studied, *CBL10* is the only duplicated gene (Fig. 2). Because the *CBL10b* genes identified contain introns and are located on a different chromosome from the *CBL10a* genes, the duplication probably did not arise through a retroduplication or an unequal crossover event.


In Arabidopsis, the CBL calcium sensors (10 members) and the CIPK protein kinases (26 members) form signaling networks



A



B




**Fig. 7.** *EsCBL10a* and the *CBL10b* genes from *S. parvula* and *S. irio* complement the *Atsos3* salt-sensitive phenotype. *CBL10* genes from Arabidopsis (*AtCBL10*, *At10*), *E. salsguineum* (*EsCBL10a*, *Es10a*; and *EsCBL10b*, *Es10b*), *S. parvula* (*SpCBL10a*, *Sp10a*; *SpCBL10b-1*, *Sp10b-1*; and *SpCBL10b-2*, *Sp10b-2*), *S. irio* (*SiCBL10a*, *Si10a*; and *SiCBL10b*, *Si10b*), and *A. arabicum* (*AaCBL10*, *Aa10*) were expressed in the *Atsos3* mutant, and growth in the absence (control) and presence of salt (75 mM NaCl) was monitored. For salt assays, seeds were germinated on medium without NaCl for 4 d, after which seedlings were transferred to medium without or with the indicated concentration of NaCl. After 10 d of treatment, photographs were taken and total seedling fresh weight was measured. (A) Photographs of the wild type (WT), *Atsos3*, and *Atsos3* expressing each *CBL10* gene (*Atsos3*;*CBL10*). The scale bar (1 cm, upper right panel) shows the magnification for all images. (B) Total seedling fresh weight was measured to quantify growth and is presented as a percentage of the control. Data are means  $\pm$ SE of at least 24 seedlings per genotype grown in three independent experiments. \*Complementation of the *Atsos3* salt-sensitive phenotype (Tukey–Kramer HSD,  $P \leq 0.05$ ).

that link changes in cytosolic calcium levels to physiological responses (Luan, 2009). The weak ability of *EsCBL10a* to activate the SOS pathway suggested that it might function with a CIPK protein other than SOS2 (CIPK24) to complement

*Atsos3* (Fig. 6; Monihan et al., 2019). Using yeast two-hybrid assays, four CIPK proteins that interact with *EsCBL10a* but not *EsCBL10b* or *AtCBL10* were identified (Monihan et al., 2019). To begin to understand if *EsCBL10a* and the *SpCBL10b*





**Fig. 8.** The SpCBL10b proteins interact with two of the four EsCBL10a-interacting CIPKs. CBL10 proteins from Arabidopsis (AtCBL10, A10), *E. salsugineum* (EsCBL10a, Es10a; and EsCBL10b, Es10b), and *S. parvula* (SpCBL10b-1, Sp10b-1; and SpCBL10b-2, Sp10b-2) were fused to the GAL4 activation domain (AD) and interaction with the Arabidopsis CIPKs fused to the GAL4 binding domain (BD) was assessed using yeast two-hybrid assays. Serial decimal dilutions of diploid yeast harboring both constructs were spotted onto synthetic defined media (SD) minus leucine (L) and tryptophan (W), or minus LW and histidine (H). Two independently mated colonies were assayed in two biological replications; one representative image is shown.

|              | Genes           | Synteny | Expression<br>Leaf Root | <i>Atcbl10</i><br>comp | SOS pathway<br>activation | <i>Atsos3</i><br>comp |
|--------------|-----------------|---------|-------------------------|------------------------|---------------------------|-----------------------|
| Arabidopsis  | <i>CBL10</i>    |         |                         |                        |                           |                       |
| Eutrema      | <i>CBL10a</i>   |         |                         |                        |                           |                       |
| Schrenkiella | <i>CBL10a</i>   |         |                         |                        |                           |                       |
| Sisymbrium   | <i>CBL10a</i>   |         |                         |                        |                           |                       |
| Eutrema      | <i>CBL10b</i>   |         |                         |                        |                           |                       |
| Schrenkiella | <i>CBL10b-1</i> |         |                         |                        |                           |                       |
| Schrenkiella | <i>CBL10b-2</i> |         |                         |                        |                           |                       |
| Sisymbrium   | <i>CBL10b</i>   |         |                         |                        |                           |                       |
| Aethionema   | <i>CBL10</i>    |         |                         |                        |                           |                       |

**Fig. 9.** *CBL10* function diverged in a species-specific manner. Synteny, genes with the same color box are syntenic (white box indicates unknown synteny; Fig. 2). Expression, presence of *CBL10* transcript in leaves and roots (blue, present; white, absent; Fig. 5). *Atcbl10* comp, ability to complement the *Atcbl10* salt-sensitive phenotype (blue, complements; white, does not complement; Fig. 4). SOS pathway activation, ability to activate the Arabidopsis SOS pathway in yeast (strongest, strong, medium, and weak refer to the level of yeast growth correlating with strength of activation of the SOS pathway; Fig. 6). *Atsos3* comp, ability to complement the *Atsos3* salt-sensitive phenotype (blue, complements; white, does not complement; Fig. 7).

proteins complement *Atsos3* through a similar mechanism, yeast two-hybrid assays were used to examine interaction between the SpCBL10b proteins and the four EsCBL10a-interacting CIPK proteins. Only AtCIPK18 interacted with EsCBL10a and both SpCBL10b proteins; however, a role for *AtCIPK18* in salt tolerance has not been reported and the *Atcipk18* mutant does not have a salt-sensitive phenotype (Fig. 8; data not shown). SpCBL10b-1 but not SpCBL10b-2 interacted with AtCIPK6 which has been shown to interact with AtSOS3 to recruit the potassium transporter, AtAKT2, to the plasma membrane (Fig. 8; Held *et al.*, 2011). The duplication appears to have expanded the range of CIPK interactions. Whether there is convergence on a specific CIPK remains an open question.

*There are multiple ways in which CBL10 might contribute to salt tolerance in the Brassicaceae*

All three *Schrenkiella* genes complement *Atcbl10* (Figs 4, 9), but further examination of function revealed differences. In *E. salsugineum*, the *CBL10* genes have different functions; EsCBL10b strongly activates the SOS pathway while *EsCBL10a* has an unknown function that allows it to complement *Atsos3* (Monihan *et al.*, 2019). In *S. parvula*, a different pattern is observed; SpCBL10a, like EsCBL10b, strongly activates the SOS pathway although not as well (Figs 6, 9), and does not have a function that complements *Atsos3* (Figs 7, 9). SpCBL10b-1 and SpCBL10b-2 can both strongly activate the SOS pathway

(like EsCBL10b although also not as well, Figs 6, 9) and complement the *Atsos3* salt-sensitive phenotype (like EsCBL10a, Figs 7, 9). These results suggest that having three *CBL10* genes with strong and overlapping functions has contributed to the ability of *S. parvula* to maintain growth in the presence of salt.

*Eutrema salsugineum* and *S. irio* have similar levels of growth in the presence of salt, the same number of *CBL10* genes, similar *CBL10* expression patterns, all *CBL10* genes in the two species complement *Atcbl10*, and one of the two *CBL10* genes in each species can complement *Atsos3* (Figs 1–7). However, activation of the SOS pathway revealed differences in how *CBL10* functions within these species. In *E. salsugineum*, EsCBL10b strongly activates the pathway while EsCBL10a does so only weakly. In *S. irio*, SiCBL10a and SiCBL10b both moderately activate the SOS pathway, suggesting that there are multiple ways to acquire salt tolerance (Figs 6, 9). As was found in *E. salsugineum*, the *CBL10* genes in *S. irio* have diverged in function, probably increasing calcium-mediated signaling capacity and contributing to the ability of *S. irio* to grow in the presence of salt.

*Aethionema arabicum* is a member of the first diverging lineage of Brassicaceae and has only one *CBL10* gene (Fig. 1). *AaCBL10* may reside in a different genomic position; a scaffold syntenic to *AtCBL10* but lacking *AaCBL10* was identified in *A. arabicum* and *AaCBL10* was found on a short scaffold (Figs 2, 9). A different genomic location for *AaCBL10* might explain the altered expression of the transcript; while the *CBL10* genes in the other species studied are expressed in shoots and some also in roots, *AaCBL10* is the only gene expressed exclusively in roots (Figs 5, 9). Several results suggest that *AaCBL10* may not play a significant role in the ability of *A. arabicum* to grow in the presence of salt: (i) *A. arabicum* is sensitive to salt, suggesting that it has few active mechanisms to deal with the presence of salt in the soil (Fig. 3); (ii) while *AaCBL10* is functional in *Atcbl10*, it is probably complementing *AtCBL10*'s function in shoots (Figs 4, 9) and may not perform a similar function in *A. arabicum* roots; (iii) *AaCBL10* is unable to complement the salt-sensitive phenotype of *Atsos3* whose gene product functions in roots (Figs 7, 9; Quan et al., 2007); and (iv) *AaCBL10* only weakly activates the SOS pathway (Figs 6, 9). Because calcium is an important signaling molecule in many different plant processes, *AaCBL10* may function in other calcium-mediated responses in *A. arabicum*—a mutational analysis will be required to uncover the role of *CBL10* in this species.

Taken together, results from this study have demonstrated that: the paralogous *CBL10* genes within a species diverged in expression and function probably contributing to the maintenance of the duplicated gene pairs in their genomes; orthologous *CBL10* genes have diverged in function in a species-specific manner, suggesting that the function of the genes was not established immediately after the duplication but after speciation; and that species studied with multiple *CBL10* genes are better able to grow in the presence of salt.

## Supplementary data

Supplementary data are available at *JXB* online.

Fig. S1. Specificity of *CBL10* gene expression in transgenic *Arabidopsis*.

Fig. S2. Transposable elements may have mediated the *CBL10* duplication.

Fig. S3. Differences in *CBL10* function reside in the N-terminus.

Table S1. Primers.

Table S2. Ratio of species growth in the absence and presence of salt.

## Acknowledgements

We thank Brian M. Ortega and Katerina A. Oskolkoff for technical assistance, and Dr Steven E. Smith (School of Natural Resources and the Environment, University of Arizona) for statistical analyses. This work was supported by the United States National Science Foundation, Division of Integrative Organismal Systems Award 1552099 (to KSS).

## Author contributions

SMM and KSS designed the research; SMM, CAM, CHR, and MMM performed the experiments; SMM, KSS, MMM, and MAB analyzed the data; and SMM and KSS wrote the manuscript with input from all co-authors.

## References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.
- Argout X, Salse J, Aury JM, et al. 2011. The genome of *Theobroma cacao*. *Nature Genetics* **43**, 101–108.
- Bartlett MS. 1937. Properties of sufficiency and statistical tests. *Proceedings of the Royal Society A: Mathematical and Physical Sciences* **160**, 268–282.
- Beilstein MA, Al-Shehbaz IA, Kellogg EA. 2006. Brassicaceae phylogeny and trichome evolution. *American Journal of Botany* **93**, 607–619.
- Beilstein MA, Nagalingum NS, Clements MD, Manchester SR, Mathews S. 2010. Dated molecular phylogenies indicate a Miocene origin for *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **107**, 18724–18728.
- Cheng S, van den Bergh E, Zeng P, et al. 2013. The *Tarenaya hassleriana* genome provides insight into reproductive trait and genome evolution of crucifers. *The Plant Cell* **25**, 2813–2830.
- Clough SJ, Bent AF. 1998. Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The Plant Journal* **16**, 735–743.
- Conover WJ, Iman RL. 1981. Rank transformations as a bridge between parametric and nonparametric statistics. *American Statistician* **35**, 124–129.
- Dassanayake M, Oh DH, Haas JS, et al. 2011. The genome of the extremophile crucifer *Thellungiella parvula*. *Nature Genetics* **43**, 913–918.
- Dewey CN. 2011. Positional orthology: putting genomic evolutionary relationships into context. *Briefings in Bioinformatics* **12**, 401–412.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**, 1792–1797.
- Egea I, Pineda B, Ortíz-Atienza A, et al. 2018. The SiCBL10 calcineurin B-like protein ensures plant growth under salt stress by regulating Na<sup>+</sup> and Ca<sup>2+</sup> homeostasis. *Plant Physiology* **176**, 1676–1693.
- Flagel LE, Wendel JF. 2009. Gene duplication and evolutionary novelty in plants. *New Phytologist* **183**, 557–564.
- Halfter U, Ishitani M, Zhu JK. 2000. The *Arabidopsis* SOS2 protein kinase physically interacts with and is activated by the calcium-binding protein SOS3. *Proceedings of the National Academy of Sciences, USA* **97**, 3735–3740.
- Han MV, Demuth JP, McGrath CL, Casola C, Hahn MW. 2009. Adaptive evolution of young gene duplicates in mammals. *Genome Research* **19**, 859–867.

- Haudry A, Platts AE, Vello E, et al.** 2013. An atlas of over 90,000 conserved noncoding sequences provides insight into crucifer regulatory regions. *Nature Genetics* **45**, 891–898.
- Held K, Pascaud F, Eckert C, et al.** 2011. Calcium-dependent modulation and plasma membrane targeting of the AKT2 potassium channel by the CBL4/CIPK6 calcium sensor/protein kinase complex. *Cell Research* **21**, 1116–1130.
- Hoagland DR, Arnon DI.** 1938. The water-culture method for growing plants without soil. *California Agricultural Experiment Station Circular* **347**, 1–39.
- Hu TT, Pattyn P, Bakker EG, et al.** 2011. The *Arabidopsis lyrata* genome sequence and the basis of rapid genome size change. *Nature Genetics* **43**, 476–481.
- Jaillon O, Aury JM, Noel B, et al.** 2007. The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* **449**, 463–467.
- Jarvis DE, Ryu CH, Beilstein MA, Schumaker KS.** 2014. Distinct roles for SOS1 in the convergent evolution of salt tolerance in *Eutrema salsugineum* and *Schrenkiella parvula*. *Molecular Biology and Evolution* **31**, 2094–2107.
- Kagale S, Koh C, Nixon J, et al.** 2014. The emerging biofuel crop *Camelina sativa* retains a highly undifferentiated hexaploid genome structure. *Nature Communications* **5**, 3706.
- Lagercrantz U, Lydiate DJ.** 1996. Comparative genome mapping in Brassica. *Genetics* **144**, 1903–1910.
- Levene H.** 1960. Robust tests for equality of variances. In: Olkin I, ed. *Contributions to probability and statistics*. Stanford, CA: Stanford University Press, 278–292.
- Lin H, Yang Y, Quan R, Mendoza I, Wu Y, Du W, Zhao S, Schumaker KS, Pardo JM, Guo Y.** 2009. Phosphorylation of SOS3-LIKE CALCIUM BINDING PROTEIN8 by SOS2 protein kinase stabilizes their protein complex and regulates salt tolerance in *Arabidopsis*. *The Plant Cell* **21**, 1607–1619.
- Liu J, Zhu JK.** 1997. An *Arabidopsis* mutant that requires increased calcium for potassium nutrition and salt tolerance. *Proceedings of the National Academy of Sciences, USA* **94**, 14960–14964.
- Liu J, Zhu JK.** 1998. A calcium sensor homolog required for plant salt tolerance. *Science* **280**, 1943–1945.
- Liu S, Liu Y, Yang X, et al.** 2014. The *Brassica oleracea* genome reveals the asymmetrical evolution of polyploid genomes. *Nature Communications* **5**, 3930.
- Luan S.** 2009. The CBL–CIPK network in plant calcium signaling. *Trends in Plant Science* **14**, 37–42.
- Lyons E, Freeling M.** 2008. How to usefully compare homologous plant genes and chromosomes as DNA sequences. *The Plant Journal* **53**, 661–673.
- Lyons E, Pedersen B, Kane J, et al.** 2008. Finding and comparing syntenic regions among *Arabidopsis* and the outgroups papaya, poplar, and grape: CoGe with rosids. *Plant Physiology* **148**, 1772–1781.
- Ma T, Wang J, Zhou G, et al.** 2013. Genomic insights into salt adaptation in a desert poplar. *Nature Communications* **4**, 2797.
- Ming R, Hou S, Feng Y, et al.** 2008. The draft genome of the transgenic tropical fruit tree papaya (*Carica papaya* Linnaeus). *Nature* **452**, 991–996.
- Moghe GD, Hufnagel DE, Tang H, Xiao Y, Dworkin I, Town CD, Conner JK, Shiu SH.** 2014. Consequences of whole-genome triplication as revealed by comparative genomic analyses of the wild radish *Raphanus raphanistrum* and three other Brassicaceae species. *The Plant Cell* **26**, 1925–1937.
- Monihan SM, Magness CA, Yadegari R, Smith SE, Schumaker KS.** 2016. *Arabidopsis* CALCINEURIN B-LIKE10 functions independently of the SOS pathway during reproductive development in saline conditions. *Plant Physiology* **171**, 369–379.
- Monihan SM, Ryu CH, Magness CA, Schumaker KS.** 2019. Linking duplication of a calcium sensor to salt tolerance in *Eutrema salsugineum*. *Plant Physiology* **179**, 1176–1192.
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ.** 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**, 268–274.
- Oh DH, Hong H, Lee SY, Yun DJ, Bohnert HJ, Dassanayake M.** 2014. Genome structures and transcriptomes signify niche adaptation for the multiple-ion-tolerant extremophyte *Schrenkiella parvula*. *Plant Physiology* **164**, 2123–2138.
- Panchy N, Lehti-Shiu M, Shiu SH.** 2016. Evolution of gene duplication in plants. *Plant Physiology* **171**, 2294–2316.
- Pegueroles C, Laurie S, Albà MM.** 2013. Accelerated evolution after gene duplication: a time-dependent process affecting just one copy. *Molecular Biology and Evolution* **30**, 1830–1842.
- Quan R, Lin H, Mendoza I, Zhang Y, Cao W, Yang Y, Shang M, Chen S, Pardo JM, Guo Y.** 2007. SCABP8/CBL10, a putative calcium sensor, interacts with the protein kinase SOS2 to protect *Arabidopsis* shoots from salt stress. *The Plant Cell* **19**, 1415–1431.
- Quintero FJ, Ohta M, Shi H, Zhu JK, Pardo JM.** 2002. Reconstitution in yeast of the *Arabidopsis* SOS signaling pathway for Na<sup>+</sup> homeostasis. *Proceedings of the National Academy of Sciences, USA* **99**, 9061–9066.
- Rentsch D, Laloi M, Rouhara I, Schmelzer E, Delrot S, Frommer WB.** 1995. NTR1 encodes a high affinity oligopeptide transporter in *Arabidopsis*. *FEBS Letters* **370**, 264–268.
- Rodríguez-Navarro A, Ramos J.** 1984. Dual system for potassium transport in *Saccharomyces cerevisiae*. *Journal of Bacteriology* **159**, 940–945.
- Roselló OPI, Kondrashov FA.** 2014. Long-term asymmetrical acceleration of protein evolution after gene duplication. *Genome Biology and Evolution* **6**, 1949–1955.
- Shapiro SS, Wilk MB.** 1965. An analysis of variance test for normality (Complete Samples). *Biometrika* **52**, 591.
- Slotte T, Hazzouri KM, Ågren JA, et al.** 2013. The *Capsella rubella* genome and the genomic consequences of rapid mating system evolution. *Nature Genetics* **45**, 831–835.
- Tang RJ, Yang Y, Yang L, Liu H, Wang CT, Yu MM, Gao XS, Zhang HX.** 2014. Poplar calcineurin B-like proteins PtCBL10A and PtCBL10B regulate shoot salt tolerance through interaction with PtSOS2 in the vacuolar membrane. *Plant, Cell & Environment* **37**, 573–588.
- Teh BT, Lim K, Yong CH, et al.** 2017. The draft genome of tropical fruit durian (*Durio zibethinus*). *Nature Genetics* **49**, 1633–1641.
- Wang X, Wang H, Wang J, et al.** 2011. The genome of the mesopolyploid crop species *Brassica rapa*. *Nature Genetics* **43**, 1035–1039.
- Windsor AJ, Schranz ME, Formanová N, Gebauer-Jung S, Bishop JG, Schnabelrauch D, Kroymann J, Mitchell-Olds T.** 2006. Partial shotgun sequencing of the *Boechera stricta* genome reveals extensive microsynteny and promoter conservation with *Arabidopsis*. *Plant Physiology* **140**, 1169–1182.
- Xiao H, Jiang N, Schaffner E, Stockinger EJ, van der Knaap E.** 2008. A retrotransposon-mediated gene duplication underlies morphological variation of tomato fruit. *Science* **319**, 1527–1530.
- Yang R, Jarvis DE, Chen H, et al.** 2013. The reference genome of the halophytic plant *Eutrema salsugineum*. *Frontiers in Plant Science* **4**, 46.
- Zhang Q, Chen W, Sun L, et al.** 2012. The genome of *Prunus mume*. *Nature Communications* **3**, 1318.