

Bioimage informatics

OrgDyn: feature- and model-based characterization of spatial and temporal organoid dynamics

Zaki Hasnain¹, Andrew K. Fraser², Dan Georgess², Alex Choi^{2,3}, Paul Macklin³, Joel S. Bader⁴, Shelly R. Peyton⁵, Andrew J. Ewald^{2,4} and Paul K. Newton^{1,6,*}

¹Department of Aerospace & Mechanical Engineering, University of Southern California, Los Angeles, CA 90089, USA, ²Department of Cell Biology and Center for Cell Dynamics, Johns Hopkins University, Baltimore, MD 21218, USA, ³Intelligent Systems Engineering, Indiana University, Bloomington, IN 47408, USA, ⁴Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 21218, USA, ⁵Department of Chemical Engineering, University of Massachusetts Amherst, Amherst, MA 01003, USA and ⁶Department of Mathematics, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA 90089, USA

*To whom correspondence should be addressed

Associate Editor: Alfonso Valencia

Received on August 26, 2019; revised on December 23, 2019; editorial decision on February 2, 2020; accepted on February 16, 2020

Abstract

Summary: Organoid model systems recapitulate key features of mammalian tissues and enable high throughput experiments. However, the impact of these experiments may be limited by manual, non-standardized, static or qualitative phenotypic analysis. OrgDyn is an open-source and modular pipeline to quantify organoid shape dynamics using a combination of feature- and model-based approaches on time series of 2D organoid contour images. Our pipeline consists of (i) geometrical and signal processing feature extraction, (ii) dimensionality reduction to differentiate dynamical paths, (iii) time series clustering to identify coherent groups of organoids and (iv) dynamical modeling using point distribution models to explain temporal shape variation. OrgDyn can characterize, cluster and model differences among unique dynamical paths that define diverse final shapes, thus enabling quantitative analysis of the molecular basis of tissue development and disease.

Availability and Implementation: <https://github.com/zakih/organoidDynamics> (BSD 3-Clause License).

Contact: newton@usc.edu

Supplementary information: [Supplementary data](#) are available at *Bioinformatics* online.

1 Introduction

Our understanding of cell, developmental and cancer biology relies upon the analysis of isolated cells cultured at low density on flat, rigid substrates. Recently, 3D cell culture systems have become more physiologically relevant models for study cell behavior (Cheung *et al.*, 2013; Gencoglu *et al.*, 2017; Nguyen-Ngoc *et al.*, 2012; Nguyen-Ngoc *et al.*, 2015; Padmanaban *et al.*, 2019; Schwartz *et al.*, 2017; Shamir and Ewald, 2014). These systems include 3D cellular spheroids or tumor organoids and 3D matrices in which to grow and visualize cells akin to the way they grow in the human body. To fully realize the potential of these 3D culture models, we urgently need better visualization and analysis methods to handle the large amount of data that results from time-dependent analysis of 3D culture models and subsequent 2D projections.

Conventional approaches to quantify the morphologies of diverse 3D cell samples have either (i) often been limited to qualitative scoring systems, such as binned percentages or categorical scales, (ii)

focused on a few spatial descriptors such as distance from centroid (Maeda *et al.*, 2008), Fourier modes (Sánchez-Corrales *et al.*, 2018), membrane extension and retraction (Satulovsky *et al.*, 2008), roundness and hollowness (Åkerfelt *et al.*, 2015) or (iii) developed comprehensive spatial descriptors for non-temporal datasets (Borten *et al.*, 2018).

OrgDyn introduces techniques for quantifying the dynamical evolution of organoid morphology by (i) extracting spatio-temporal shape descriptors from time series of 2D organoid contours, (ii) capturing phenotypic spatio-temporal heterogeneity in clusters and (iii) modeling and quantifying dynamical variations (Fig. 1A). At the core of the pipeline are geometrical- and signal processing-based descriptive features of organoid shape, as these are often the focus of shape assessment studies (Borten *et al.*, 2018; Kriegel *et al.*, 2018; Meijering *et al.*, 2012; Meijering, 2012; Pincus and Theriot, 2007; Sánchez-Corrales *et al.*, 2018; Zimmer *et al.*, 2002). We introduce local curvature shape descriptors, and leverage all these features concurrently to increase robustness of spatio-temporal phenotypic




Fig. 1. OrgDyn tools and example. (A) OrgDyn pipeline. (B) Sample FGf2 treated organoid DIC images and contours using OrgDyn script for ImageJ. (C) Area and polar moment of area time series for FGf2 and basal organoids. (D) PDM metrics for 39 organoids grouped by (E) hierarchical clustering of 39 organoids' time series in principal component space

analyses. A standardized and intuitive pipeline of organoid morphometric analysis can benefit the experimental community by providing a common path to quantification and form a baseline for future advancements in analytical techniques.

2 Materials and methods

The input for OrgDyn is a time series of 2D organoid contours (step 1 in Fig. 1A). Point distribution models (PDM) of each organoid are generated, and metrics of dynamical complexity are calculated (step 2 in Fig. 1A). Ten features are extracted per contour: area A , perimeter P , form factor a_f , solidity a_b , polar moment of area J_{zz} , fraction of convex f_{vex} and concave f_{cav} points, and the number of modes N_{90} , the mean mode amplitude \bar{A}_{90} and standard deviation σ_{90} of mode amplitudes in the 90% discrete Fourier transform of a contour (step 3 in Fig. 1A). Together, these features form multivariate time series for each organoid, and these are cast to principal component (PC) space and then clustered (steps 4 and 5 Fig. 1A). Detailed methods are presented in Supplementary Section 2.

2.1 Numerical implementation

The preprocessing, feature extraction and PDM algorithms are implemented using MATLAB 2017b. Dimensionality reduction and clustering algorithms are implemented in R version 3.4.3.

3 Example

We demonstrate OrgDyn on a 3D culture model of normal mouse mammary development (Ewald *et al.*, 2008; Nguyen-Ngoc *et al.*, 2015) consisting of basal and FGf2-treated organoid groups. The organoids were imaged every 30 min using differential interference contrast (DIC) microscopy for 130 h, creating a time series of contours of each organoid's boundary (Fig. 1B). Among the 10 features (Fig. 1A, step 3), area and the polar moment of area showed a dramatic difference in the growth trajectories (Fig. 1C). Each contour of an organoid's time series is cast in reduced PC space in Figure 1E, and connecting together these points creates a reduced dimension time series of each organoid which encompasses the majority of variance in the original features. Clustering the organoids' PC space time series

reveals groups of organoids which have similar dynamical histories. The hierarchical clustering splits cluster-1 from the other clusters first, forming the division between basal and FGF2 types. However, each cluster represents unique phenotypes, and clusters 2–5 are all FGF2 subtypes ranging from the most to least complex morphologies in clusters-2 and cluster-3, respectively (Fig. 1E). Figure 1D shows the PDM metrics grouped by the clusters in Fig. 1E, where the low variance captured by the first mode and AUC of the mode-variance curve confirm that cluster-2 organoids require the greatest number of dynamical modes to capture their evolution. Detailed example is presented in [Supplementary Section 3](#).

Acknowledgements

We gratefully acknowledge partial support from the Breast Cancer Research Foundation (BCRF) and the Jayne Koskinas & Ted Giovanis Foundation (JKTG) for Health and Policy.

Funding

This work was supported by the Jayne Koskinas Ted Giovanis (JKTG) Foundation for Health and Policy and the Breast Cancer Research Foundation, private foundations committed to critical funding of cancer research. The opinions, findings, conclusions or recommendations expressed in this material are those of the author(s) and not necessarily those of the JKTG foundation. Dan Georgess was supported by a Postdoctoral Fellowship Grant from the Susan G. Komen Foundation [PDF15332336]. National Institutes of Health/National Cancer Institute [U01CA217846 to A.J.E. and J.S.B.; U54CA2101732 to A.J.E.]. A.J.E. was supported by NIH/National Institute of General Medical Sciences(NIGMS) T32GM007309.

Conflict of Interest: none declared.

References

- Åkerfelt, M. *et al.* (2015) Automated tracking of tumor-stroma morphology in microtissues identifies functional targets within the tumor microenvironment for therapeutic intervention. *Oncotarget*, **6**, 30035.
- Borten, M.A. *et al.* (2018) Automated brightfield morphometry of 3D organoid populations by OrganoSeg. *Sci. Rep.*, **8**, 5319.
- Cheung, K.J. *et al.* (2013) Collective invasion in breast cancer requires a conserved basal epithelial program. *Cell*, **155**, 1639–1651.
- Ewald, A.J. *et al.* (2008) Collective epithelial migration and cell rearrangements drive mammary branching morphogenesis. *Dev. Cell*, **14**, 570–581.
- Gencoglu, M.F. *et al.* (2017) Comparative study of multicellular tumor spheroid formation methods and implications for drug screening. *ACS Biomater. Sci. Eng.*, **4**, 410–420.
- Kriegel, F.L. *et al.* (2018) Cell shape characterization and classification with discrete fourier transforms and self-organizing maps. *Cytometry Part A*, **93**, 323–333.
- Maeda, Y.T. *et al.* (2008) Ordered patterns of cell shape and orientational correlation during spontaneous cell migration. *PLoS One*, **3**, e3734.
- Meijering, E. (2012) Cell segmentation: 50 years down the road. *IEEE Signal Process. Mag.*, **29**, 140–145.
- Meijering, E. *et al.* (2012) Methods for cell and particle tracking. In: *Methods in Enzymology*. Vol. 504. Elsevier, pp. 183–200.
- Nguyen-Ngoc, K.-V. *et al.* (2012) ECM microenvironment regulates collective migration and local dissemination in normal and malignant mammary epithelium. *Proc. Natl. Acad. Sci. USA*, **109**, E2595–E2604.
- Nguyen-Ngoc, K.-V. *et al.* (2015) 3D culture assays of murine mammary branching morphogenesis and epithelial invasion. In: *Tissue Morphogenesis*. Humana Press, New York, NY, pp. 135–162.
- Padmanaban, V. *et al.* (2019) E-cadherin is required for metastasis in multiple models of breast cancer. *Nature*, **573**, 439–444.
- Pincus, Z. and Theriot, J. (2007) Comparison of quantitative methods for cell-shape analysis. *J. Microscopy*, **227**, 140–156.
- Sánchez-Corrales, Y.E. *et al.* (2018) Morphometrics of complex cell shapes: lobe contribution elliptic Fourier analysis (LOCO-EFA). *Development*, **145**, 1–13. doi: 10.1242/dev.156778.
- Satulovsky, J. *et al.* (2008) Exploring the control circuit of cell migration by mathematical modeling. *Biophys. J.*, **94**, 3671–3683.
- Schwartz, A.D. *et al.* (2017) A biomaterial screening approach reveals micro-environmental mechanisms of drug resistance. *Integr. Biol.*, **9**, 912–924.
- Shamir, E.R. and Ewald, A.J. (2014) Three-dimensional organotypic culture: experimental models of mammalian biology and disease. *Nat. Rev. Mol. Cell Biol.*, **15**, 647–664.
- Zimmer, C. *et al.* (2002) Segmentation and tracking of migrating cells in video-microscopy with parametric active contours: a tool for cell based drug testing. *IEEE Trans. Med. Imaging*, **21**, 1212–1221.