

RESEARCH

Open Access



Engineering central pathways for industrial-level (3R)-acetoin biosynthesis in *Corynebacterium glutamicum*

Lingxue Lu^{1†}, Yufeng Mao^{2†}, Mengyun Kou¹, Zhenzhen Cui¹, Biao Jin¹, Zhishuai Chang¹, Zhiwen Wang¹, Hongwu Ma² and Tao Chen^{1*}

Abstract

Background: Acetoin, especially the optically pure (3S)- or (3R)-enantiomer, is a high-value-added bio-based platform chemical and important potential pharmaceutical intermediate. Over the past decades, intense efforts have been devoted to the production of acetoin through green biotechniques. However, efficient and economical methods for the production of optically pure acetoin enantiomers are rarely reported. Previously, we systematically engineered the GRAS microorganism *Corynebacterium glutamicum* to efficiently produce (3R)-acetoin from glucose. Nevertheless, its yield and average productivity were still unsatisfactory for industrial bioprocesses.

Results: In this study, cellular carbon fluxes in the acetoin producer CGR6 were further redirected toward acetoin synthesis using several metabolic engineering strategies, including blocking anaplerotic pathways, attenuating key genes of the TCA cycle and integrating additional copies of the *alsSD* operon into the genome. Among them, the combination of attenuation of citrate synthase and inactivation of phosphoenolpyruvate carboxylase showed a significant synergistic effect on acetoin production. Finally, the optimal engineered strain CGS11 produced a titer of 102.45 g/L acetoin with a yield of 0.419 g/g glucose at a rate of 1.86 g/L/h in a 5 L fermenter. The optical purity of the resulting (3R)-acetoin surpassed 95%.

Conclusion: To the best of our knowledge, this is the highest titer of highly enantiomerically enriched (3R)-acetoin, together with a competitive product yield and productivity, achieved in a simple, green processes without expensive additives or substrates. This process therefore opens the possibility to achieve easy, efficient, economical and environmentally-friendly production of (3R)-acetoin via microbial fermentation in the near future.

Keywords: (3R)-Acetoin, *Corynebacterium glutamicum*, Metabolic engineering, Microbial fermentation, Citrate synthase, Green chemistry

Background

Acetoin (3-hydroxy-2-butanone) is a popular food

additive with a pleasant butter-like flavor [1]. Since it can be obtained from biomass and possesses reactive carbonyl and hydroxyl moieties, the United States Department of Energy designated acetoin as one of 30 promising platform chemicals that were given priority for development and utilization in 2004 [2]. Acetoin is chiral, and its enantiomers (3S)- and (3R)-acetoin are more valuable than the racemate as they are potential pharmaceutical intermediates [3]. Moreover, they can be used

*Correspondence: chentao@tju.edu.cn

[†]Lingxue Lu and Yufeng Mao contributed equally to this work

¹Frontier Science Center for Synthetic Biology and Key Laboratory of Systems Bioengineering of Ministry of Education, SynBio Research Platform, Collaborative Innovation Center of Chemical Science and Engineering, School of Chemical Engineering and Technology, Tianjin University, Tianjin 300072, China

Full list of author information is available at the end of the article



© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

to synthesize liquid crystal materials and novel optically active α -hydroxyketone derivatives [4].

At present, the commercial production of acetoin is mostly based on chemical methods with many disadvantages, such as high cost, high pollution, and low yield [5]. Moreover, the use of chemosynthetic, non-natural acetoin in food and cosmetics is restricted due to safety concerns [6, 7]. The biotechnological production of safe and natural acetoin could be more ecological and sustainable than their chemical counterparts [8, 9]. These methods, including microbial fermentation [8, 10, 11], whole-cell biocatalysis [12, 13] and enzymatic biocatalysis [14, 15], have consequently gained great attention over the past decades. *Saccharomyces cerevisiae* JHY617-SDN was able to efficiently accumulate 100.2 g/L acetoin from glucose during fed-batch fermentation [16]. *Gluconobacter oxydans* NL71 could produce 165.9 g/L acetoin from 2,3-butanediol in a whole-cell catalysis process [17]. This is the highest acetoin titer ever reported. However, the enantiomeric excess of the produced acetoin was not reported, and efficient methods for the production of optically pure (3*R*)-acetoin were rarely reported. Nevertheless, some representative publications with good results do exist. Using a two-enzyme coupling system, 12.2 g/L of optically pure (3*S*)-acetoin was efficiently produced from diacetyl with a high productivity of 9.76 g/L/h [18]. By constructing an efficient *E. coli* whole-cell biocatalyst, Guo et al. [19] obtained a high (3*R*)-acetoin titer of 86.7 g/L from optically pure (*R,R*)-2,3-butanediol. However, the expensive chiral substrate, and the complicated, costly processing steps, such as protein purification through Ni-NTA affinity chromatography or centrifugation to concentrate the catalyst cells, made it economically unfeasible in industrial applications.

Acetoin production from petrochemicals such as the expensive chiral 2,3-butanediol or the noxious diacetyl is costly and unsustainable. Consequently, (3*R*)-acetoin production from renewable substrates via microbial fermentation is highly favored in recent studies. Many microorganisms can naturally synthesize optically pure (3*R*)-acetoin, including *Klebsiella pneumoniae* [20], *Serratia marcescens* [21, 22], *Lactococcus lactis* [23] and *Bacillus* sp. [24]. When the acetoin degradation pathway was disrupted, the *Klebsiella pneumoniae* strain $\Delta budC\Delta aco$ was able to accumulate 62.3 g/L (3*R*)-acetoin in 57 h [20]. However, this pathogenic microorganism can be hazardous to humans, which entails unacceptable safety risks in industrial-scale production. Dai et al. [25] isolated the marine strain *Bacillus subtilis* CGMCC 13141 capable of producing 83.7 g/L of (3*R*)-acetoin with a high yield of 0.447 g/g glucose


in fed-batch fermentation. However, the titer was still insufficient for industrial demands.

In our previous study [26], we systematically engineered *Corynebacterium glutamicum*, the famous amino acids industrial workhorse with GRAS (generally recognized as safe) status, to develop a safe and efficient industrial (3*R*)-acetoin producer. The best strain CGR7 was able to produce 96.2 g/L (3*R*)-acetoin with an optical purity of more than 95% in fed-batch mode with a yield of 0.360 g/g glucose and productivity of 1.30 g/L/h [9], highlighting *C. glutamicum* as a competitive producer of enantiopure acetoin with fantastic potential for industrial-level production. While this was still the highest reported titer at the time of writing, the corresponding yield (73.5% of the theoretical value) and average productivity were still unsatisfactory for economical production. In this study, the previously engineered strain CGR6 (ATCC13032 $\Delta pta\Delta ackA\Delta ldh\Delta butA\Delta nagD$; $\Delta ackA::P_{tuf}-alsSD$) [9] was chosen as a chassis for further optimization. In this strain, the biosynthesis pathways of the major by-products lactate, glycerin, and acetate, as well as the downstream product 2,3-butanediol were disrupted to increase the pool of the precursor pyruvate and prevent acetoin from being reduced to downstream 2,3-butanediol. Moreover, a copy of the acetoin synthesis operon *alsS-alsD* under the control of the strong constitutive promoter P_{tuf} was inserted into the locus of the deleted gene *ackA* to enhance acetoin production. Here, several strategies were successively applied to redirect more carbon flux toward acetoin synthesis in strain CGR6, including interdicting anaplerotic pathways, weakening key genes of the TCA cycle, and inserting additional copies of the *alsSD* operon into the genome of the host. The optimal engineered strain CGS11 (Fig. 1) achieved a titer of 102.45 g/L acetoin with a yield of 0.419 g/g glucose at a rate of 1.86 g/L/h in fed-batch fermentation.

Materials and methods

Reagents, strains and media

Primers were synthesized by GENEWIZ (Suzhou, China). Plasmids were extracted using the Axyprep™ Plasmid Miniprep Kit (Axygen, USA) and the SanPrep Column Plasmid Mini-Prep Kit (Sangon Biotech, Shanghai, China). BHI Broth was purchased from Hopebio (Qingdao, China). Yeast extract was purchased from Angel (Hubei, China). Acetoin, creatine, 1-naphthol, ethyl 2-acetoxy-2-methylacetoacetate, oxaloacetate and 5,5'-dithiobis(2-nitrobenzoic) acid were purchased from Sigma (Merck, USA). The 2,3,5,6-tetramethylpyrazine standard was purchased from Tokyo Chemical Industry (Tokyo, Japan). Other reagents were purchased from Sangon Biotech (Shanghai, China).



The strains and plasmids used in this study were listed in Table 1. *Escherichia coli* DH5 α was used for plasmid construction and was grown in lysogeny broth (LB) medium containing (per liter) 10 g tryptone, 5 g yeast extract, and 10 g NaCl. Brain heart infusion (BHI) medium was used for the transformation and tube culture of *C. glutamicum*. CGIII medium composed of (per liter) 10 g tryptone, 10 g yeast extract, 2.5 g NaCl and 20 g glucose was used for pre-cultures of *C. glutamicum*. Batch fermentation of acetoin was conducted in CGX-IIP medium containing (per liter): 10 g yeast extract, 5 g (NH₄)₂SO₄, 5 g urea, 1 g KH₂PO₄, 1 g K₂HPO₄, 0.25 g MgSO₄·7H₂O, 0.01 g CaCl₂, 0.01 g FeSO₄·7H₂O, 0.1 mg MnSO₄·H₂O, 1 mg ZnSO₄·7H₂O, 0.2 mg CuSO₄·5H₂O, 0.02 mg NiCl₂·6H₂O, 0.4 mg biotin and 38 g glucose, pH 7.0. Fed-batch fermentation was conducted in LBRC medium containing (per liter): 10 g yeast extract,

50 g corn steep liquor, 1 g urea, 0.5 g K₂HPO₄, 0.5 g MgSO₄·7H₂O and 2 g sodium acetate, supplemented with the indicated amount of glucose, pH 7.0. The stock comprising 1000 g/L glucose used for fed-batch fermentation, was prepared by putting 100 g glucose and 36 mL ddH₂O in a Schott-Duran bottle and sterilizing at 110 °C for 10 min, then stored at 60 °C. Antibiotics were added where appropriate as follows: for *C. glutamicum*, kanamycin 25 mg/L, for *E. coli*, kanamycin 40 mg/L.

Construction of plasmids and strains

All the primers used in this study are listed in supplementary Additional file 1: Table S1. All DNA manipulations, including restriction enzyme digestion and vector isolation were carried out using standard protocols [27]. The suicide plasmid pD-*sacB* was used for genome editing in

Table 1 Strains and plasmids used in this study

Strain/plasmid	Relevant characteristics	References
<i>E. coli</i> DH5a	Host for plasmid cloning	Invitrogen
ATCC 13032	<i>C. glutamicum</i> wild type Biotin auxotrophic	ATCC ^a
CGR6	ATCC13032 Δ pta Δ ackA Δ ldh Δ butA Δ nagD, Δ ackA::P _{tuf} -alsSD	[9]
CGS1	CGR6 Δ ppc	This study
CGS2	CGR6 Δ pyc	This study
CGS3	CGR6 Δ ppc Δ pyc	This study
CGS4	CGS3 ICD ^{A94D}	This study
CGS5	CGS3 ICD ^{G407S}	This study
CGS6	CGS3 ICD ^{R453S}	This study
CGS7	CGR6 P ₁ -gltA	This study
CGS8	CGS1 P ₁ -gltA	This study
CGS9	CGS8 Δ butA::P _{tuf} -alsSD, Δ nagD::P _{tuf} -alsSD	This study
CGS11	CGS9 pEC-XK99E-alsSD- Δ laclq	This study
Plasmids		
pD-sacB	Kan ^R ; vector for in-frame deletion (<i>sacB</i> _{B,sub} ; <i>lacZa</i> ; OriV _{E.c.})	[56]
pD-sacB-ppc	pD-sacB carrying the flanking sequences of the <i>ppc</i> gene	This study
pD-sacB-pyc	pD-sacB carrying the flanking sequences of the <i>pyc</i> gene	This study
pD-sacB-butA	pD-sacB carrying the flanking sequences of the <i>butA</i> gene	[41]
pD-sacB-nagD	pD-sacB carrying the flanking sequences of the <i>nagD</i> gene	[9]
pD-sacB-icd-mut-1	Kan ^R , containing the nucleotide sequence for ICD amino acid exchange A94D	This study
pD-sacB-icd-mut-2	Kan ^R , containing the nucleotide sequence for ICD amino acid exchange G407S	This study
pD-sacB-icd-mut-3	Kan ^R , containing the nucleotide sequence for ICD amino acid exchange R453S	This study
pD-sacB-P ₁ -gltA	Kan ^R , containing P ₁ promoter and <i>gltA</i> flanks	Unpublished work
pD-sacB-butA-alsSD	Kan ^R , containing P _{tuf} -alsSD flanks	This study
pD-sacB-nagD-alsSD	Kan ^R , containing P _{tuf} -alsSD flanks	This study
pEC-XK99E	Kan ^R ; <i>C. glutamicum</i> / <i>E. coli</i> shuttle vector (P _{trc} , <i>laclq</i> ; pGA1, OriV _{C.g.} , OriV _{E.c.})	[57]
pEC-XK99E-alsSD- Δ laclq	Derived from pEC-XK99E, for the overexpression of <i>alsS</i> and <i>alsD</i> under promoter P _{trc}	[9]

^a ATCC, American Type Culture Collection

C. glutamicum via two-step homologous recombination [28].

To delete the *pyc* gene, the vector pD-sacB-*pyc* was constructed as follows: the upstream sequence of *pyc* and a *pyc*-specific fragment were amplified from *C. glutamicum* genomic DNA by PCR using the primer pairs *pyc*-1/*pyc*-2 and *pyc*-3/*pyc*-4, respectively, and fused by PCR. The resulting product was digested with *Bam*HI and *Sal*II and ligated between the corresponding sites of pD-sacB. The plasmid pD-sacB-*ppc* was constructed analogously, using *Bam*HI and *Hind*III.

To introduce mutations into the endogenous gene *icd*, the plasmid pD-sacB-*icd*-mut-1 was constructed as follows: the flanking regions of the *icd* gene with relevant modifications were amplified from genomic DNA of *C. glutamicum* using the primer pairs ICD^{A94D}-1/ICD^{A94D}-2 and ICD^{A94D}-3/ICD^{A94D}-4. The corresponding flanking fragments were fused using ICD^{A94D}-1/ICD^{A94D}-4. The fused product was digested with *Bam*HI and *Sal*II and

ligated between the corresponding sites of pD-sacB to construct pD-sacB-*icd*-mut-1. The plasmids pD-sacB-*icd*-mut-2 and pD-sacB-*icd*-mut-3 were constructed analogously, using *Sma*I and *Sal*II.

To integrate additional copies of the acetoin operon into the chromosome, the plasmid pD-sacB-*butA*-alsSD was constructed as follows: the acetoin operon (*alsS*-*alsD*) was amplified from the genome of CGR6 using the primer pair BalsSD-1/BalsSD-2. The resulting fragment was digested with *Xho*I/*Sal*II and ligated between the corresponding sites of pD-sacB-*butA*. The plasmid pD-sacB-*nagD*-alsSD was constructed analogously.

Fermentation conditions

To prepare *C. glutamicum* pre-cultures, single colonies were used to inoculate 5 mL of BHI medium and grown at 30 °C and 220 rpm overnight, after which the entire resulting culture was used to inoculate 50 mL of CGIII medium, and grown to an OD₆₀₀ of 10. For batch fermentation, the

seed culture was used to inoculate a 250-mL shake flask containing 50 mL of CGXIIP medium to an initial OD_{600} of 1 and grown at 30 °C and 220 rpm on a rotary shaker.

For fed-batch fermentation, 100 mL of LBRC seed culture was used to inoculate a 5-L fermenter (Bailun, Shanghai, China) containing 1.8 L of LBRC medium. All cultivations were carried out at 30 °C with an aeration rate of 1 vvm. The agitation speed was maintained at 600 rpm. The initial pH of the medium was 6.5. During the fermentation process, the pH value was not controlled. The initial glucose concentration was 50 g/L, and an appropriate amount of 1000 g/L glucose stock was added to maintain its concentration between 10 and 50 g/L.

Analytical methods

The biomass concentration was calculated from OD_{600} values using an experimentally determined correlation 1 OD_{600} unit is equal to 0.25 g/L cell dry weight (CDW) [29]. Glucose was measured using an SBA Bio-analyzer (Shandong Academy of Sciences, China) after appropriate dilution. Metabolite concentrations were determined by HPLC, using an HPX-87H (300 mm × 7.8 mm) organic acid and sugar analysis column (Bio-Rad, China), kept at 60 °C, and a refractive index detector. The injection volume was 10 μ L, and the mobile phase consisted of 5 mM H_2SO_4 at a flow rate of 0.4 mL/min. The concentration of 2,3,5,6-tetramethylpyrazine was analyzed by GC-FID (PERSEE, Beijing, China) equipped with an HP-5 (19091 J-413, 30 m × 0.32 mm) capillary column (Agilent, USA) as described previously [30]. The optical purity of acetoin was determined by GC as described previously [9]. The fermentation broth was quenched by 40% pre-cooled (−40 °C) methanol and centrifuged at 5000 × g for 1 min immediately to harvest cells, then resuspended cells with extracting solution provided by the Pyruvate Assay Kit (Solarbio, Beijing, China) and disrupted cells by ultrasonication. The reaction, including the construction of pyruvate standard curves, was carried out using the Pyruvate Assay Kit.

Enzyme activity assays

Cells of the engineered strains were grown to the middle exponential phase (12 h) and late exponential phase (22 h), after which the fermentation broth was centrifuged at 5000 × g for 10 min to harvest cells, which were washed twice with 1 mL 200 mM phosphate buffered saline (pH 7.0) and then resuspended in 0.3 mL of the same buffer. The cells were disrupted by ultrasonication and cell debris was removed by centrifugation at 5000 × g and 4 °C for 10 min. Enzyme activity was assayed in the resulting supernatant. Total protein concentrations were determined according to the Bradford method [31]. The

formed amount of acetoin generated by acetolactate was measured to determine the acetolactate synthase (ALS) activity as described previously [32]. The acetolactate decarboxylase (ALDC) activity was assayed by measuring the production of acetoin as described previously [33]. The citrate synthase activity was determined by measuring the amount of CoA formed as described previously [34].

Results and discussion

Improvement of acetoin production by deleting *ppc/pyc* to reduce succinate accumulation

To analyze the carbon distribution in strain CGR6, batch fermentation in flasks was carried out. As shown in Fig. 2a, CGR6 was able to produce 11.30 g/L acetoin when glucose was almost depleted at 24 h, corresponding to a yield of 0.302 g/g glucose. As expected, no 2,3-butandiol or lactate was detected, since their synthesis pathways were blocked. Although 0.52 g/L α -ketoglutarate was detected when fermenting this strain in CGXIIP medium [9], it was not detected in CGXIIP medium in this study. The major by-product was succinate, with a titer of 2.63 g/L, followed by acetate (1.12 g/L) and glycerin (0.23 g/L). In our previous work, two key genes of succinate synthesis, *ppc* encoding phosphoenolpyruvate (PEP) carboxylase and *pyc* encoding pyruvate carboxylase, were respectively deleted via one-step homologous recombination (single crossover) with a tetracycline resistance marker [9]. The *ppc/pyc*-deficient strain showed an improved acetoin production in batch fermentation in flasks. However, the use of the tetracycline resistance marker impeded further genetic manipulation and was not acceptable for an industrial producer. In this study, we deleted the *ppc* and *pyc* genes in CGR6 using a markerless two-step recombination method (double crossover) [35], resulting in the strains CGS1 and CGS2, respectively.

As shown in Fig. 2b, succinate production was almost completely abolished in the *ppc* knockout strain CGS1, with a titer of only 0.03 g/L. The glucose consumption rate of CGS1 was decreased compared with that of CGR6, and 11.96 g/L of acetoin was obtained when glucose was exhausted at 29 h. This corresponds to a yield of 0.328 g/g glucose, which was 9.3% higher than that of CGR6. By contrast, the succinate production in the *pyc* knockout strain CGS2 was almost unchanged (Fig. 2c), but its acetoin production was enhanced with a titer of 11.75 g/L when glucose was depleted at 29 h, and a yield of 0.323 g/g glucose. The glucose consumption rate of CGS2 was also decreased compared with that of CGR6, but it was somewhat higher than that of CGS1 at 24 h.

Mutants deficient in *ppc* rather than *pyc* could efficiently reduce succinate accumulation, indicating that




Fig. 2 Time profiles of the biomass (g/L), glucose, organic acid and acetoin concentrations of strains CGR6 (a), CGS1 (b) and CGS2 (c) cultured with 38 g/L glucose. Error bars indicate the standard deviations from three independent cultures

Table 2 Fermentation characteristics of *C. glutamicum* strains cultivated in CGXIIP medium supplemented with initial 38 g/L glucose measured at 24 h

Strain	Biomass (g/L)	Consumed glucose (g/L)	Acetoin yield (g/g glucose)	Acetoin productivity (g/L/h)	Titer (g/L)			
					Acetoin	Acetate	Glycerin	Succinate
CGR6	10.29±0.27	37.42±0.06	0.302±0.009	0.47±0.01	11.30±0.26	1.12±0.14	0.23±0.02	2.63±0.15
CGS1	10.19±0.03	32.54±0.59	0.309±0.005	0.42±0.01	10.06±0.26	0.06±0.01	0.29±0.01	0.03±0.01
CGS2	10.18±0.11	33.50±1.39	0.326±0.020	0.45±0.04	10.91±0.86	0.44±0.04	0.27±0.03	2.67±0.22
CGS5	8.67±0.34	24.55±1.08	0.272±0.009	0.28±0.02	6.68±0.54	0.72±0.07	0.32±0.01	<0.01
CGS7	9.58±0.21	35.84±1.43	0.337±0.011	0.50±0.01	12.07±0.38	0.58±0.03	0.20±0.03	2.02±0.13
CGS8	9.34±0.12	30.17±0.98	0.415±0.005	0.52±0.01	12.53±0.03	0.15±0.01	0.17±0.03	0.03±0.02
CGS9	7.88±0.45	24.45±1.52	0.484±0.015	0.49±0.01	11.83±0.35	0.29±0.03	0.08±0.01	0.03±0.02
CGS11	8.06±0.44	24.54±0.46	0.497±0.005	0.51±0.01	12.22±0.20	0.14±0.01	0.05±0.01	0.03±0.01

Error bars indicate the standard deviations from three independent cultures

PEP rather than pyruvate is the key precursor of succinate under aerobic conditions. In agreement with previous reports [36, 37], the growth of CGS1 and CGS2 was only mildly decreased, but their acetate titers (respectively 0.06 and 0.44 g/L) were unexpectedly significantly decreased compared with that of CGR6 (1.12 g/L) (Table 2). We suspected that the reduced acetate accumulation in CGS1 and CGS2 was resulted from the decreased glucose consumption rates (Table 2), which can reduce the flux in overflow metabolism. The only another detected by-product was glycerin, with a titer of about 0.30 g/L in both CGS1 and CGS2. The two strains both showed improved acetoin yields, which was consistent with our previous results [9]. However, strain CGS3 with *ppc* and *pyc*-deficient was unable to grow on glucose in mineral medium and still grew poorly with 10 g/L yeast extract addition (data not shown), which was consistent with previous reports [37, 38]. After considering the acetoin titer, yield and by-products accumulation, strain CGS1 was chosen for further manipulation.

Improvement of acetoin production by blocking anaplerotic pathways and introducing isocitrate dehydrogenase mutants

Most studies on improving precursor pyruvate (Fig. 1) availability focused on complete inactivation or attenuation of the pyruvate dehydrogenase complex (PDHC) by deleting the *aceE* gene or reducing its promoter activity [39, 40]. In our previous work, the *aceE* gene was also deleted to conserve pyruvate and improve acetoin production. However, additional acetate was required for cell growth and the best strain CGL3 (*C. glutamicum* $\Delta aceE\Delta ldh\Delta butA$; pEC-XK99E-*alsSD*) could only accumulate 8.33 g/L acetoin from about 33 g/L glucose and 10 g/L acetate under optimal conditions [41]. In cases of attenuating PDHC, although the growth of the resulting

strain was independent of acetate addition, the cell growth and glucose consumption rate were dramatically reduced [39]. Therefore, deletion or attenuation of PDHC might not be optimal for manufacturing optically pure (3R)-acetoin.

A double deletion of *ppc* and *pyc* is generally lethal for *C. glutamicum* due to a lack of oxaloacetate [37]. However, a novel strategy for improving the biosynthesis of pyruvate-derived metabolites by introducing newly identified isocitrate dehydrogenase mutants (A94D, G407S or R453S) was recently proposed [38]. These *icd* mutations both lowered the ICD activity and activated the glyoxylate shunt, and were consequently able to recover oxaloacetate re-supply and cell growth of *C. glutamicum* $\Delta ppc\Delta pyc$ [38]. Thus, three ICD mutants A94D, G407S and R453S were introduced into CGS3 to test their effects on acetoin production, resulting in strains CGS4, CGS5 and CGS6, respectively. As shown in Additional file 1: Fig. S1, strains CGS4 and CGS6 showed almost the same growth inhibition as strain CGS3. This is probably because strain CGS3 is an acetoin producing host, in which the major carbon fluxes have already been redirected toward acetoin synthesis, and are therefore insufficient to support an activated glyoxylate shunt to re-supply oxaloacetate. Nevertheless, CGS5 with the mutant ICD^{G407S} successfully recovered cell growth to some extent. However, acetoin production in CGS5 was dramatically decreased to a titer of only 11.37 g/L when glucose was exhausted at 34 h, and a yield of 0.280 g/g glucose, which was 14.6% lower than that of CGS1 (Additional file 1: Fig. S2). This indicated that the carbon fluxes from acetoin synthesis rather than its competing pathways were redirected to regain biomass synthesis, but the reason for this is still unknown.

Improvement of acetoin production by reducing citrate synthase activity

Since directly blocking acetyl-CoA or oxaloacetate synthesis severely affected the growth performance and failed to improve acetoin production, the focus of engineering was moved to citrate synthase (CS, encode by *gltA*), which condenses acetyl-CoA and oxaloacetate to citrate and is one of the most important sinks for the flux of these two precursors. Therefore, reduction of CS activity was viewed as a promising strategy to lower the synthesis of acetyl-CoA and oxaloacetate, and thereby conserve pyruvate for improved acetoin production. *C. glutamicum* mutants devoid of CS were unable to grow on glucose [34]. Therefore, a reduction rather than inactivation of CS activity was preferable. Moreover, regulation of the *gltA* gene, which is responsible for 95% of CS activity in *C. glutamicum* [42], was widely applied to improve the biosynthesis of pyruvate-derived products [34, 43–45]. However, the impact of reducing CS activity on acetoin production is still unclear. Therefore, we used a weak promoter P_I with approximately 1% activity of the P_{tac} [46] to replace the native promoter of *gltA* gene in CGR6 and CGS1, yielding strains CGS7 and CGS8 respectively.

The strains' CS activity was measured to confirm its successful downregulation. As shown in Additional file 1: Fig. S3A, the CS activity of CGS7 at 12 h in the middle exponential phase was 75.8% lower than that of CGR6, and 60.0% lower at 22 h in the late exponential phase. As shown in Table 2, the succinate yield of CGS7 at 24 h was 20.0% lower than that of CGR6 (0.056 versus 0.070 g succinate/g glucose). Moreover, the cell growth and glucose consumption were also decreased (Fig. 3a), suggesting that the flux into the TCA cycle was indeed reduced. Unexpectedly, the acetate yield decreased by 46.7% (0.016 versus 0.030 g acetate/g glucose). We speculated that the decreased glucose consumption rate in CGS7 would reduce the fluxes to acetate synthesis caused by overflow metabolism, and the newly available carbon fluxes flowed to acetoin effectively, leading to an increase of the acetoin yield at 24 h in contrast to CGR6 (0.337 versus 0.302 g acetoin/g glucose). Then, the yield further increased to 0.350 g/g glucose with a titer of 13.61 g/L when glucose was depleted at 29 h (Fig. 3a). However, the results indicated that neither weakening *gltA* nor the deletion of *ppc* could effectively improve the intracellular pyruvate pool, but a combination of both eventually led to a 31.1% increase of intracellular pyruvate (Additional file 1: Fig. S3B). At the same time, the acetoin production was significantly enhanced to 14.56 g/L, with a yield of 0.389 g/g glucose when glucose was exhausted at 29 h, which was 18.6% higher than that of CGS1 (Fig. 3b). While deletion of *ppc* obviously inhibited glucose consumption of CGS8 compared to CGS7 (Table 2), however, the acetoin

productivity was 0.52 g/L/h at 24 h, even higher than that of CGS7 (0.50 g/L/h). Moreover, the by-products acetate (0.15 g/L), glycerin (0.17 g/L) and succinate (0.03 g/L) remained at low concentrations. After considering acetoin production and by-product accumulation, CGS8 was selected for further engineering.

Furthermore, reduction of CS exhibited a fantastic synergistic effect on acetoin production with inactivation of PEP carboxylase, which is responsible for 90% of total oxaloacetate synthesis in *C. glutamicum* [47, 48]. As shown in Table 2, *ppc* knockout in CGS1 almost had no effect on acetoin yield (0.309 versus 0.302 g/g glucose), and *gltA* attenuation in CGS7 only led to a 11.6% increase in acetoin yield (0.337 versus 0.302 g/g glucose). However, the combination of the two gene modifications resulted in that the acetoin yield was significantly improved by 37.4% (0.415 versus 0.302 g/g glucose). The variations of intracellular pyruvate further confirmed the synergistic effect of CS and PEP carboxylase on acetoin production (Additional file 1: Fig. S3B), which would also be potential for improving other pyruvate-acetolactate-derived metabolites production.

Effect of enhancing the acetoin synthesis pathway on acetoin production

To further pull the carbon flux from pyruvate toward product accumulation, two additional *alsSD* operon (under the control of the constitutive promoter P_{tuj}) copies were inserted into the chromosome of CGS8 at the $\Delta butA$ and $\Delta nagD$ sites to generate the strain CGS9.

The activities of ALS and ALDC were listed in Table 3. The ALS activity of CGS9 was 2.33-fold higher than that of CGR6 at 12 h in the middle log phase, and subsequently increased by 11.6% at 22 h in late log phase. However, the ALDC activity of CGS9 was increased by only 47.0% compared with that of CGR6 at 12 h. Then, it decreased by 15.1% at 22 h, but was still 37.3% higher than that of CGR6. With the increase of *alsSD* copies, the intracellular pyruvate concentration was slightly decreased by 8.2%. However, when glucose was exhausted at 34 h, 15.70 g/L acetoin was accumulated with a yield of 0.408 g/g glucose (Fig. 4a), which was merely 4.9% higher than that of CGS8. Therefore, the increased ALS and ALDC activity appeared to still be insufficient to increase acetoin production.

To further pull pyruvate into acetoin synthesis pathway, the previously constructed *alsSD* over-expression plasmid pEC-XK99E-*alsSD*- $\Delta lacIq$ [9], was introduced into CGS9 to yield CGS11. Both the activities of ALS and ALDC were significantly improved compared with CGS9. Moreover, the intracellular pyruvate pool was further decreased by 6.7% compared with that of CGS9 (Additional file 1: Fig. S3B), indicating that more flux




Fig. 3 Time profiles of the biomass (g/L), glucose, organic acid and acetoin concentrations of strains CGS7 (a) and CGS8 (b) cultured with 38 g/L glucose. Error bars indicate the standard deviations from three independent cultures

might have been re-directed toward acetoin synthesis. However, the acetoin production was still not obviously enhanced. When glucose was depleted at 34 h, the final acetoin titer was 16.10 g/L with a yield of 0.419 g/g glucose (Fig. 4b), which was only 7.7% higher

than that of CGS8. Although the intracellular pyruvate pool of CGS11 was still 12.1% higher than that of CGR6, the carbon fluxes from pyruvate to competing pathways were successfully controlled as the titers of the main by-products, acetate (0.29 versus 1.12 g/L),

Table 3 Enzyme activities of different *C. glutamicum* strains

Strains	Middle log phase (12 h)		Late log phase (22 h)	
	ALS activity (U/mg)	ALDC activity (U/mg)	ALS activity (U/mg)	ALDC activity (U/mg)
CGR6	0.511 ± 0.087	0.608 ± 0.067	0.521 ± 0.060	0.553 ± 0.045
CGS9	1.192 ± 0.094	0.894 ± 0.044	1.330 ± 0.122	0.759 ± 0.054
CGS11	8.242 ± 0.362	2.864 ± 0.127	6.383 ± 0.778	1.074 ± 0.035

Error bars indicate the standard deviations from three independent cultures

glycerin (0.06 versus 0.23 g/L) and succinate (0.12 versus 2.63 g/L), were much lower than those of CGR6. No 2,3-butandiol or lactate was detected. Moreover, despite the decreases of cell growth and glucose consumption, the acetoin productivity was still comparable to that of strain CGS8. Therefore, strain CGS11 was a promising candidate for further scale-up fermentation.

Notably, both strain CGS9 and CGS11 showed much higher acetoin yields at 24 h (Table 2). The acetoin yield of strain CGS11 was even beyond the maximum theoretical yield of 0.489 g/g glucose and reached 0.498 g/g glucose at 24 h, which was mainly caused by the rich nutrients from the yeast extract in the CGXIIP medium. Moreover, all the detected by-products remained at low concentrations and biomass was relatively lower in CGS11 at 24 h (Table 2). Unexpectedly, the acetoin yield was then noticeably decreased to 0.419 g/g glucose at 34 h. The decreases of ALS and ALDC activities over time (Table 3) were initially suspected to be responsible for the decreased acetoin yield. However, the yields of by-products or biomass were not accordingly increased. Another possibility is that acetoin was re-used as an alternative carbon source when glucose was almost exhausted. Acetoin utilization is mainly catalyzed by the acetoin dehydrogenase (AoDH) complex [20, 49]. However, no candidate gene encoding a putative AoDH was identified in *C. glutamicum* through homologous sequence alignment (data not shown). Furthermore, acetoin degradation was more clearly observable when glucose was exhausted in fermentations of strains CGR6, CGS1 and CGS2 (Fig. 2a–c), but their biomass instead decreased during acetoin degradation. Thus, it did not appear that acetoin was consumed as a reserve carbon source, which would be expected to further support cell growth.

It is worth noting that two molecules of acetoin ($C_4H_8O_2$) can be chemically converted to one molecule of 2,3,5,6-tetramethylpyrazine (TMP, $C_8H_{12}N_2$, also called ligustrazine) in the presence of inorganic ammonium salts such as $(NH_4)_2SO_4$ or diammonium phosphate [6, 30]. As shown in Additional file 1: Fig. S4, no TMP was accumulated in the fermentation broth of strain CGS11 at 24 h, but a titer of 1.27 g/L TMP was indeed detected at 34 h, which corresponded to a consumption of 1.64 g/L

acetoin. Taking this portion of the generated acetoin into calculation, the yield could even reach a high level of 0.462 g/g glucose, which indicated that 94.4% of carbon fluxes had been directed toward acetoin synthesis.

Fed-batch fermentation of CGS11 for efficient (3R)-acetoin production

After considering the promising shake-flask results, strain CGS11 was chosen for fed-batch fermentation in order to evaluate its potential for further industrial application. Before scale-up production, CGS11 was evaluated in the LBRC medium, which was optimized for acetoin production in *C. glutamicum* in our previous work [9]. As shown in Fig. 5, the acetoin production was significantly enhanced with a titer of 23.53 g/L (3R)-acetoin and a yield of 0.553 g/g glucose when glucose was exhausted at 29 h. The acetoin production in CGS11 was also much higher than that in our previous optimal strain CGR7 (17.10 g/L (3R)-acetoin with a yield of 0.428 g/g glucose from LBRC medium) [9]. The rich nutrients such as lactate (initial 2.07 g/L) and amino acids from corn steep liquor (CSL) in LBRC medium should be the major reason for the improved acetoin production. And the remarkable effect on acetoin production with CSL addition was consistent with previous report in *B. subtilis* [50, 51]. Moreover, when using organic nitrogen source to replace inorganic nitrogen source $(NH_4)_2SO_4$, no TMP was detected (Additional file 1: Fig. S4), which might be another reason for the improved acetoin production. Given that CSL is cheap and abundant, the cost caused by addition of CSL could be easily made up by the significant enhancement of acetoin production. The LBRC medium was therefore adopted for further fed-batch fermentation.

During the fed-batch fermentation process, the cell growth, residual glucose, product concentrations and relative dissolved oxygen (DO) were determined. As shown in Fig. 6, a titer of 102.45 g/L acetoin was obtained at 55 h, corresponding to an average productivity of 1.86 g/L/h. The final fermentation volume was 2.45 L and a total of 625.85 g glucose was added in 7 batches, while the remaining glucose amount at 55 h was 11 g/L. Therefore, the acetoin yield was 0.419 g/g glucose, reaching 85.7% of the theoretical yield. The




Fig. 4 Time profiles of the biomass (g/L), glucose, organic acid and acetoin concentrations of strains CGS9 (a) and CGS11 (b) cultured with 38 g/L glucose. Error bars indicate the standard deviations from three independent cultures

optical purity of the produced (3*R*)-acetoin surpassed 95% (Additional file 1: Fig. S5), which was consistent with our previous results [9]. No TMP was detected during the entire fed-batch fermentation (Additional file 1: Fig. S4). The final concentrations of acetate,

succinate, glycerin and lactate were 1.94, 0.60, 0.41 and 0.33 g/L, respectively. However, α -ketoglutarate, which was undetectable during batch fermentation in shake flask, started to accumulate after 33 h and reached a final concentration of 1.71 g/L at 55 h. Nevertheless,




Fig. 5 Time profiles of the biomass (g/L), glucose, organic acid and acetoin concentrations of strain CGS11 in LBRC medium in shake fermentation




Fig. 6 (3R)-acetoin production from glucose using CG11 in fed-batch fermentation. The strain was cultured in LBRC medium with initial 50 g/L glucose at 30 °C and 600 rpm in a 5-L fermenter under aeration of 1 vvm. A stock solution comprising 1000 g/L glucose was added when the glucose concentration dropped below 15 g/L to keep the glucose concentration between 10 and 50 g/L

all the by-products remained at acceptably low concentrations.

We noticed that the glucose consumption and acetoin production were prone to stop when acetoin titer surpassed 100 g/L (from 55 to 62 h, as shown in Fig. 6). With an initial acetoin concentration of 80 g/L in shake fermentation, the specific growth rate of *B. amyloliquefaciens* was decreased by 99% [52], while wild-type *B. subtilis* 168 was even unable to grow on plates containing 50 g/L acetoin [53]. Therefore, the toxicity of the very high acetoin concentrations will no doubt prevent glucose assimilation and acetoin production of CGS11, and even activate its underlying acetoin catabolism. Since acetoin metabolism is complex and still not clearly

elucidated, several successful strategies, such as physical/chemical mutagenesis [54], adaptive evolution [52, 53] or omics focusing on global transcriptional/metabolism level responses to acetoin stress [55], could be adopted to deeply understand and improve the acetoin tolerance of *C. glutamicum* in future studies.

To our best knowledge, the value of 102.45 g/L is the highest titer of highly enantiomerically enriched (3*R*)-acetoin reported to date (Table 4), as well as the best result obtained for acetoin production via microbial fermentation. Moreover, the yield and productivity were also good enough to merit further industrial application. Furthermore, the strains engineered in this work could also be used as platform strains to be further developed

Table 4 Comparison of the production of acetoin via microbial fermentation process in the literature and in this study

Strains/enzymes	Substrate	Titer (g/L)	Enantiomer	Yield (g/g)	Productivity (g/L/h)	Refs.
Production of optically pure acetoin						
<i>C. glutamicum</i> CGS11	Glucose	102.45	(3 <i>R</i>)-	0.419	1.86	This study
<i>C. glutamicum</i> CGR7	Glucose	96.20	(3 <i>R</i>)-	0.360	1.30	[9]
<i>Bacillus subtilis</i> CGMCC 13141	Glucose	83.70	(3 <i>R</i>)-	0.448	1.02	[25]
<i>Klebsiella pneumoniae</i>	Glucose	62.30	(3 <i>R</i>)-	0.140	1.09	[20]
<i>E. coli</i> (pAC-NOX)	Glucose	60.30	(3 <i>R</i>)-	0.422	1.55	[58]
<i>Bacillus subtilis</i> DL01	Sugarcane molasses	58.20	(3 <i>R</i>)-	0.291	0.77	[59]
<i>Lactococcus lactis</i> CS4701m	Glucose	5.80	(3 <i>S</i>)-	0.347	0.19	[23]
Production of racemic acetoin (or optical purity not stated)						
<i>S. cerevisiae</i> JHY617-SDN	Glucose	100.10	Ng	0.440	1.82	[16]
<i>Gluconobacter oxydans</i> DSM 2003	2,3-Butanediol	89.20	Ng	0.891	1.24	[60]
<i>B. subtilis</i> HS019	Glucose	82.50	Ng	0.460	1.33	[53]
<i>Bacillus licheniformis</i> WX-02 Δ <i>budC</i> Δ <i>acoR</i>	Glucose	78.80	Ng	0.310	0.58	[7]
<i>S. marcescens</i> H32- <i>nox</i>	Sucrose	75.20	Ng	0.360	1.88	[61]
<i>Enterobacter aerogenes</i> EJW-03	Glucose	71.70	Ng	0.320	2.87	[62]
<i>B. amyloliquefaciens</i> E-11	Glucose	71.50	Ng	0.413	1.63	[52]
<i>B. licheniformis</i> MW3 (Δ <i>budC</i> Δ <i>gdh</i>)	Glucose	64.20	Ng	0.412	2.38	[63]
<i>B. subtilis</i> ZB02	Glucose, xylose, and arabinose	62.20	Ng	0.288	0.86	[64]
<i>S. marcescens</i> H32	Sucrose	60.50	Ng	Ng	1.44	[65]
<i>B. subtilis</i> F126-2	Glycerol	60.48	Ng	Ng	1.26	[66]
<i>Lactococcus lactis</i> CML B4	Glucose	59.00	Ng	0.360	2.11	[67]
<i>Paenibacillus polymyxa</i> CJX518	Glucose	57.20	Ng	0.400	0.30	[68]
<i>B. subtilis</i> TH-49	Glucose	56.90	Ng	Ng	0.89	[69]
<i>B. subtilis</i> BMN	Glucose	56.70	Ng	0.378	0.68	[70]
<i>P. polymyxa</i> CS107	Glucose	55.30	Ng	0.370	1.32	[71]
<i>B. subtilis</i> JNA-UD-6	Glucose	53.90	Ng	0.359	0.37	[72]
<i>Enterobacter cloacae</i> SDM 45	Glucose	55.20	Ng	0.373	2.69	[73]
	Lignocellulosic hydrolysate	45.60	Ng	NG	1.52	
<i>B. amyloliquefaciens</i> FMME044	Glucose	51.20	Ng	0.430	1.42	[74]
<i>Bacillus</i> sp. H15-1.	Degermed maize flour hydrolysate	50.80	Ng	0.285	NG	[75]
<i>C. glutamicum</i> CGT2	Glucose	40.51	Ng	0.240	0.51	[76]

Ng indicates that the related information is Not Given

as microbial cell factories for production of (3S)-acetoin and chiral 2,3-butanediol.

Conclusions

In summary, this is the first report on reducing CS activity to improve acetoin production. Moreover, combination of attenuation of CS and inactivation of PEP showed a significant synergistic effect on acetoin production. The highly engineered *C. glutamicum* strain CGS11 produced a titer of 102.45 g/L acetoin with a yield of 0.419 g/g glucose at a rate of 1.86 g/L/h in a 5 L fermenter, which is the highest titer of highly enantiomerically enriched (3R)-acetoin together with a competitive product yield and productivity, which opens the possibility of finally realizing the promises of industrial (3R)-acetoin production via green chemical process in the near future.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12934-020-01363-8>.

Additional file 1: Table S1 Primers used in this study. **Fig. S1.** Growth curves of strains CGS3, CGS4, CGS5, CGS6 and CGR6 in CGXIIY medium (per liter) containing 1 g yeast extract, 5 g (NH₄)₂SO₄, 5 g urea, 1 g KH₂PO₄, 1 g K₂HPO₄, 0.25 g MgSO₄·7H₂O, 0.01 g CaCl₂, 0.01 g FeSO₄·7H₂O, 0.1 mg MnSO₄·H₂O, 1 mg ZnSO₄·7H₂O, 0.2 mg CuSO₄·5H₂O, 0.02 mg NiCl₂·6H₂O and 0.4 mg biotin and 40 g glucose, pH 7.0. **Fig. S2.** Time profiles of the biomass (g/L), glucose, organic acid and acetoin concentrations of strain CGS5 in CGXIIIP medium. **Fig. S3.** (A) The activity of citrate synthase in CGR6 and CGS7. (B) The intracellular pyruvate concentrations of different strains at 12 h. Error bars indicate the standard deviations from three independent cultures. **Fig. S4.** Measurement of 2,3,5,6-tetramethylpyrazine concentrations by GC-FID. A: The 2,3,5,6-tetramethylpyrazine standard had a retention time of 10.598 min; B: Batch fermentation products of CGS11 in CGXIIIP medium at 24 h; C: Batch fermentation products of CGS11 in CGXIIIP medium at 34 h; D: Batch fermentation products of CGS11 in LBRC medium at 29 h; E: Fed-batch fermentation products of CGS11 in LBRC medium at 55 h. **Fig. S5.** Identification of acetoin enantiomers by GC-FID. A: The optically pure standards of (3R)- and (3S)- had retention times of 9.215 and 9.598 min, respectively; B: Fermentation products of CGS11 in CGXIIIP medium; C: Fed-batch fermentation products of CGS11 in LBRC medium at 55 h.

Acknowledgements

Not applicable.

Authors' contributions

LL, YM, and TC conceived and designed the experiments. LL, MK, ZC, BJ and ZC performed the experiments. YM, LL, TC, ZW and HM analyzed the data. YM, LL and TC wrote and revised this manuscript. TC supervised the work. All authors contributed to the discussion of the research. All authors read and approved the final manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (Grants NSFC-21576191, NSFC-21621004 and NSFC-21776208).

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Frontier Science Center for Synthetic Biology and Key Laboratory of Systems Bioengineering of Ministry of Education, SynBio Research Platform, Collaborative Innovation Center of Chemical Science and Engineering, School of Chemical Engineering and Technology, Tianjin University, Tianjin 300072, China. ² Key Laboratory of Systems Microbial Biotechnology, Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences, Tianjin 300308, China.

Received: 23 February 2020 Accepted: 5 May 2020

Published online: 12 May 2020

References

- Xiao Z, Lu JR. Generation of acetoin and its derivatives in foods. *J Agric Food Chem*. 2014;62:6487–97.
- Werpy T, Petersen GE. Top value added chemicals from biomass. Washington, DC: Department of Energy; 2004.
- Xiao Z, Xu P. Acetoin metabolism in bacteria. *Crit Rev Microbiol*. 2007;33:127–40.
- Saito S IH, Ohno K: Alpha-hydroxyketone derivatives, liquid crystal compositions containing said derivatives, and liquid crystal devices using said compositions. US Patent, 5164112. 1992.
- Gu L, Lu T, Li X, Zhang Y. A highly efficient thiazolydene catalyzed acetoin formation: reaction, tolerance and catalyst recycling. *Chem Commun*. 2014;50:12308–10.
- Xu Y, Jiang Y, Li X, Sun B, Teng C, Yang R, Xiong K, Fan G, Wang W. Systematic characterization of the metabolism of acetoin and its derivative ligustrazine in *Bacillus subtilis* under micro-oxygen conditions. *J Agric Food Chem*. 2018;66:3179–87.
- Li L, Wei X, Yu W, Wen Z, Chen S. Enhancement of acetoin production from *Bacillus licheniformis* by 2,3-butanediol conversion strategy: metabolic engineering and fermentation control. *Process Biochem*. 2017;57:35–42.
- Yang T, Rao Z, Zhang X, Xu M, Xu Z, Yang ST. Metabolic engineering strategies for acetoin and 2,3-butanediol production: advances and prospects. *Crit Rev Biotechnol*. 2017;37:990–1005.
- Mao YF, Fu J, Tao R, Huang C, Wang ZW, Tang YJ, Chen T, Zhao XM. Systematic metabolic engineering of *Corynebacterium glutamicum* for the industrial-level production of optically pure D-(–)-acetoin. *Green Chem*. 2017;19:5691–702.
- Drejfer EB, Chan DTC, Haupka C, Wendisch VF, Brautaset T, Irla M. Methanol-based acetoin production by genetically engineered *Bacillus methanolicus*. *Green Chem*. 2020;22:788–802.
- Windhorst C, Gescher J. Efficient biochemical production of acetoin from carbon dioxide using *Cupriavidus necator* H16. *Biotechnol Biofuels*. 2019;12:163.
- Xiao Z, Lv C, Gao C, Qin J, Ma C, Liu Z, Liu P, Li L, Xu P. A novel whole-cell biocatalyst with NAD⁺ regeneration for production of chiral chemicals. *PLoS ONE*. 2010;5:e8860.
- Samuel N, Bao T, Zhang X, Yang T, Xu M, Li X, Komera I, Philibert T, Rao Z. Optimized whole cell biocatalyst from acetoin to 2,3-butanediol through coexpression of acetoin reductase with NADH regeneration systems in engineered *Bacillus subtilis*. *J Chem Technol Biotechnol*. 2017;92:2477–87.
- Cui Z, Mao Y, Zhao Y, Chen C, Tang Y-J, Chen T, Ma H, Wang Z. Concomitant cell-free biosynthesis of optically pure D-(–)-acetoin and xylitol via a novel NAD⁺ regeneration in two-enzyme cascade. *J Chem Technol Biotechnol*. 2018;93:3444–51.
- Cui Z, Zhao Y, Mao Y, Shi T, Lu L, Ma H, Wang Z, Chen T. *In vitro* biosynthesis of optically pure D-(–)-acetoin from meso-2,3-butanediol using 2,3-butanediol dehydrogenase and NADH oxidase. *J Chem Technol Biotechnol*. 2019;94:2547–54.
- Bae SJ, Kim S, Hahn JS. Efficient production of acetoin in *Saccharomyces cerevisiae* by disruption of 2,3-butanediol dehydrogenase and expression of NADH oxidase. *Sci Rep*. 2016;6:27667.
- Zhou X, Zhou X, Zhang H, Cao R, Xu Y. Improving the performance of cell biocatalysis and the productivity of acetoin from 2,3-butanediol using a compressed oxygen supply. *Process Biochem*. 2018;64:46–50.

18. Gao C, Zhang L, Xie Y, Hu C, Zhang Y, Li L, Wang Y, Ma C, Xu P. Production of (3S)-acetoin from diacetyl by using stereoselective NADPH-dependent carbonyl reductase and glucose dehydrogenase. *Biores Technol*. 2013;137:111–5.
19. Guo Z, Zhao X, He Y, Yang T, Gao H, Li G, Chen F, Sun M, Lee JK, Zhang L. Efficient (3R)-acetoin production from meso-2,3-butanediol using a new whole-cell biocatalyst with co-expression of meso-2,3-butanediol dehydrogenase, NADH oxidase, and *Vitreoscilla* hemoglobin. *J Microbiol Biotechnol*. 2017;27:92–100.
20. Wang D, Zhou J, Chen C, Wei D, Shi J, Jiang B, Liu P, Hao J. R-acetoin accumulation and dissimilation in *Klebsiella pneumoniae*. *J Ind Microbiol Biotechnol*. 2015;42:1105–15.
21. Lv X, Dai L, Bai F, Wang Z, Zhang L, Shen Y. Metabolic engineering of *Serratia marcescens* MG1 for enhanced production of (3R)-acetoin. *Bioresour Bioprocess*. 2016;3:52.
22. Bai F, Dai L, Fan J, Truong N, Rao B, Zhang L, Shen Y. Engineered *Serratia marcescens* for efficient (3R)-acetoin and (2R,3R)-2,3-butanediol production. *J Ind Microbiol Biotechnol*. 2015;42:779–86.
23. Liu X, Solem C, Jensen PR. Integrating biocompatible chemistry and manipulating cofactor partitioning in metabolically engineered *Lactococcus lactis* for fermentative production of (3S)-acetoin. *Biotechnol Bioeng*. 2016;113:2744–8.
24. Zhong H, Wang L, Zhao J-Y, Xiao Z. Fermentative production of chiral acetoin by wild-type *Bacillus* strains. *Prep Biochem Biotechnol*. 2019;50:116–22.
25. Dai J, Wang Z, Xiu Z-L. High production of optically pure (3R)-acetoin by a newly isolated marine strain of *Bacillus subtilis* CGMCC 13141. *Bioprocess Biosyst Eng*. 2019;42:475–83.
26. Becker J, Rohles CM, Wittmann C. Metabolically engineered *Corynebacterium glutamicum* for bio-based production of chemicals, fuels, materials, and healthcare products. *Metab Eng*. 2018;50:122–41.
27. Deininger P. Molecular cloning: a laboratory manual, 2nd ed. In: Sambrook J, Fritsch EF, Maniatis T. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1989. Analytical Biochemistry. 1990, 186:182–183.
28. Eggeling L, Bott M. Handbook of *Corynebacterium glutamicum*. Boca Raton: CRC Press; 2010.
29. Meiswinkel TM, Rittmann D, Lindner SN, Wendisch VF. Crude glycerol-based production of amino acids and putrescine by *Corynebacterium glutamicum*. *Biores Technol*. 2013;145:254–8.
30. Xiao Z, Hou X, Lyu X, Xi L, Zhao J-Y. Accelerated green process of tetramethylpyrazine production from glucose and diammonium phosphate. *Biotechnol Biofuels*. 2014;7:106.
31. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*. 1976;72:248–54.
32. Holtzclaw WD, Chapman LF. Degradative acetolactate synthase of *Bacillus subtilis*: purification and properties. *J Bacteriol*. 1975;121:917–22.
33. Yang Y-T, Peredelchuk M, Bennett GN, San K-Y. Effect of variation of *Klebsiella pneumoniae* acetolactate synthase expression on metabolic flux redistribution in *Escherichia coli*. *Biotechnol Bioeng*. 2000;69:150–9.
34. van Ooyen J, Noack S, Bott M, Reth A, Eggeling L. Improved L-lysine production with *Corynebacterium glutamicum* and systemic insight into citrate synthase flux and activity. *Biotechnol Bioeng*. 2012;109:2070–81.
35. Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene*. 1994;145:69–73.
36. Blombach B, Buchholz J, Busche T, Kalinowski J, Takors R. Impact of different CO₂/HCO₃⁻ levels on metabolism and regulation in *Corynebacterium glutamicum*. *J Biotechnol*. 2013;168:331–40.
37. Peters-Wendisch PG, Kreutzer C, Kalinowski J, Patek M, Sahn H, Eikmanns BJ. Pyruvate carboxylase from *Corynebacterium glutamicum*: characterization, expression and inactivation of the *pyc* gene. *Microbiology*. 1998;144(Pt 4):915–27.
38. Schwentner A, Feith A, Münch E, Busche T, Rückert C, Kalinowski J, Takors R, Blombach B. Metabolic engineering to guide evolution—creating a novel mode for L-valine production with *Corynebacterium glutamicum*. *Metab Eng*. 2018;47:31–41.
39. Buchholz J, Schwentner A, Brunnenkan B, Grabris C, Grimm S, Gerstmeir R, Takors R, Eikmanns BJ, Blombach B. Platform engineering of *Corynebacterium glutamicum* with reduced pyruvate dehydrogenase complex activity for improved production of L-lysine, L-valine, and 2-ketoisovalerate. *Appl Environ Microbiol*. 2013;79:5566.
40. Blombach B, Schreiner ME, Holatko J, Bartek T, Oldiges M, Eikmanns BJ. L-Valine production with pyruvate dehydrogenase complex-deficient *Corynebacterium glutamicum*. *Appl Environ Microbiol*. 2007;73:2079–84.
41. Ma H, Liu H, Zhu N, Chen T. Metabolic engineering of *Corynebacterium glutamicum* for acetoin production. *Tianjin Daxue Xuebao (Ziran Kexue yu Gongcheng Jishu Ban) J Tianjin Univ Sci Technol*. 2014;47:967–72.
42. Radmacher E, Eggeling L. The three tricarboxylate synthase activities of *Corynebacterium glutamicum* and increase of L-lysine synthesis. *Appl Microbiol Biotechnol*. 2007;76:587.
43. Milke L, Ferreira P, Kallscheuer N, Braga A, Vogt M, Kappelmann J, Oliveira J, Silva AR, Rocha I, Bott M, et al. Modulation of the central carbon metabolism of *Corynebacterium glutamicum* improves malonyl-CoA availability and increases plant polyphenol synthesis. *Biotechnol Bioeng*. 2019;116:1380–91.
44. Vogt M, Haas S, Klaffl S, Polen T, Eggeling L, van Ooyen J, Bott M. Pushing product formation to its limit: metabolic engineering of *Corynebacterium glutamicum* for L-leucine overproduction. *Metab Eng*. 2014;22:40–52.
45. Ma Y, Cui Y, Du L, Liu X, Xie X, Chen N. Identification and application of a growth-regulated promoter for improving L-valine production in *Corynebacterium glutamicum*. *Microb Cell Fact*. 2018;17:185.
46. Zhang S, Liu D, Mao Z, Mao Y, Ma H, Chen T, Zhao X, Wang Z. Model-based reconstruction of synthetic promoter library in *Corynebacterium glutamicum*. *Biotech Lett*. 2018;40:819–27.
47. Petersen S, de Graaf AA, Eggeling L, Mollney M, Wiechert W, Sahn H. In vivo quantification of parallel and bidirectional fluxes in the anaplerosis of *Corynebacterium glutamicum*. *J Biol Chem*. 2000;275:35932–41.
48. Park SM, Shaw-Reid C, Sinskey AJ, Stephanopoulos G. Elucidation of anaplerotic pathways in *Corynebacterium glutamicum* via ¹³C-NMR spectroscopy and GC-MS. *Appl Microbiol Biotechnol*. 1997;47:430–40.
49. Wang M, Fu J, Zhang X, Chen T. Metabolic engineering of *Bacillus subtilis* for enhanced production of acetoin. *Biotech Lett*. 2012;34:1–9.
50. Tian Y, Fan Y, Liu J, Zhao X, Chen W. Effect of nitrogen, carbon sources and agitation speed on acetoin production of *Bacillus subtilis* SF4-3. *Electron J Biotechnol*. 2016;19:41–9.
51. Yang TW, Rao ZM, Zhang X, Xu MJ, Xu ZH, Yang ST. Effects of corn steep liquor on production of 2,3-butanediol and acetoin by *Bacillus subtilis*. *Process Biochem*. 2013;48:1610–7.
52. Luo Q, Wu J, Wu M. Enhanced acetoin production by *Bacillus amyloliquefaciens* through improved acetoin tolerance. *Process Biochem*. 2014;49:1223–30.
53. Wang S, Hou Y, Chen X, Liu L. Kick-starting evolution efficiency with an autonomous evolution mutation system. *Metab Eng*. 2019;54:127–36.
54. Jia X, Peng X, Liu Y, Han Y. Conversion of cellulose and hemicellulose of biomass simultaneously to acetoin by thermophilic simultaneous saccharification and fermentation. *Biotechnol Biofuels*. 2017;10:232.
55. Yuan H, Xu Y, Chen Y, Zhan Y, Wei X, Li L, Wang D, He P, Li S, Chen S. Metabolomics analysis reveals global acetoin stress response of *Bacillus licheniformis*. *Metabolomics*. 2019;15:25.
56. Zhu N, Xia H, Wang Z, Zhao X, Chen T. Engineering of acetate recycling and citrate synthase to improve aerobic succinate production in *Corynebacterium glutamicum*. *PLoS ONE*. 2013;8:e60659.
57. Kirchner O, Tauch A. Tools for genetic engineering in the amino acid-producing bacterium *Corynebacterium glutamicum*. *J Biotechnol*. 2003;104:287–99.
58. Xu Q, Xie L, Li Y, Lin H, Sun S, Guan X, Hu K, Shen Y, Zhang L. Metabolic engineering of *Escherichia coli* for efficient production of (3R)-acetoin. *J Chem Technol Biotechnol*. 2015;90:93–100.
59. Dai J-Y, Cheng L, He Q-F, Xiu Z-L. High acetoin production by a newly isolated marine *Bacillus subtilis* strain with low requirement of oxygen supply. *Process Biochem*. 2015;50:1730–4.
60. Wang X, Lv M, Zhang L, Li K, Gao C, Ma C, Xu P. Efficient bioconversion of 2, 3-butanediol into acetoin using gluconobacter oxydans DSM 2003. *Biotechnol Biofuels*. 2013;6(1):92.
61. Sun J-A, Zhang L-Y, Rao B, Shen Y-L, Wei D-Z. Enhanced acetoin production by *Serratia marcescens* H32 with expression of a water-forming NADH oxidase. *Biores Technol*. 2012;119:94–8.

62. Jang JW, Jung HM, Im DK, Jung MY, Oh MK. Pathway engineering of *Enterobacter aerogenes* to improve acetoin production by reducing by-products formation. *Enzyme Microbial Technol.* 2017;106:114–8.
63. Lü C, Ge Y, Cao M, Guo X, Liu P, Gao C, Xu P, Ma C. Metabolic engineering of *Bacillus licheniformis* for production of acetoin. *Front Bioeng Biotechnol.* 2020;8:125.
64. Zhang B, Li X-L, Fu J, Li N, Wang Z, Tang Y-J, Chen T. Production of acetoin through simultaneous utilization of glucose, xylose, and arabinose by engineered *Bacillus subtilis*. *PLoS ONE.* 2016;11:e0159298.
65. Sun J, Zhang L, Rao B, Han Y, Chu J, Zhu J, Shen Y, Wei D. Enhanced acetoin production by *Serratia marcescens* H32 using statistical optimization and a two-stage agitation speed control strategy. *Biotechnol Bioprocess Eng.* 2012;17:598–605.
66. Fan X, Wu H, Jia Z, Li G, Li Q, Chen N, Xie X. Metabolic engineering of *Bacillus subtilis* for the co-production of uridine and acetoin. *Appl Microbiol Biotechnol.* 2018;102:8753–62.
67. Roncal T, Caballero S. Díaz de Guereñu MDM, Rincón I, Prieto-Fernández S, Ochoa-Gómez JR. Efficient production of acetoin by fermentation using the newly isolated mutant strain *Lactococcus lactis* subsp. *lactis* CML B4. *Process Biochem.* 2017;58:35–41.
68. Liu L, Xu QM, Chen T, Cheng JS, Yuan YJ. Artificial consortium that produces riboflavin regulates distribution of acetoin and 2,3-butanediol by *Paenibacillus polymyxa* CJX518. *Eng Life Sci.* 2017;17:1039–49.
69. Xu H, Jia S, Liu J. Production of acetoin by *Bacillus subtilis* TH-49. In 2011 International Conference on Consumer Electronics, Communications and Networks (CECNet); 16–18 April 2011. 2011, pp 1524–1527.
70. Zhang X, Zhang R, Bao T, Rao Z, Yang T, Xu M, Xu Z, Li H, Yang S. The rebalanced pathway significantly enhances acetoin production by disruption of acetoin reductase gene and moderate-expression of a new water-forming NADH oxidase in *Bacillus subtilis*. *Metab Eng.* 2014;23:34–41.
71. Zhang L, Chen S, Xie H, Tian Y, Hu K. Efficient acetoin production by optimization of medium components and oxygen supply control using a newly isolated *Paenibacillus polymyxa* CS107. *J Chem Technol Biotechnol.* 2012;87:1551–7.
72. Zhang X, Zhang R, Yang T, Zhang J, Xu M, Li H, Xu Z, Rao Z. Mutation breeding of acetoin high producing *Bacillus subtilis* blocked in 2,3-butanediol dehydrogenase. *World J Microbiol Biotechnol.* 2013;29:1783–9.
73. Zhang L, Liu Q, Ge Y, Li L, Gao C, Xu P, Ma C. Biotechnological production of acetoin, a bio-based platform chemical, from a lignocellulosic resource by metabolically engineered *Enterobacter cloacae*. *Green Chem.* 2016;18:1560–70.
74. Zhang Y, Li S, Liu L, Wu J. Acetoin production enhanced by manipulating carbon flux in a newly isolated *Bacillus amyloliquefaciens*. *Biores Technol.* 2013;130:256–60.
75. Xiao Z, Gu R, Hou X, Zhao JY, Zhu H, Lu JR. Non-sterilized fermentative production of acetoin with 2,3-butanediol as a main byproduct from maize hydrolysate by a newly isolated thermophilic *Bacillus* strain. *J Chem Technol Biotechnol.* 2017;92:2845–52.
76. Tao R, Mao YF, Jing FU, Huang C, Wang ZW, Chen T. Effects of inactivation of acetate synthetic pathway genes and overexpression of NADH oxidase on acetoin production in *Corynebacterium glutamicum*. *Microbiol China.* 2017;44:2530–8.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

