

Clinical efficacy and safety of stem cell therapy for knee osteoarthritis

A meta-analysis

Rui Huang, MD^a, Wei Li, MD^b, Ying Zhao, MD^c, Fan Yang, MD^d, Meng Xu, PhD^{e,*}

Abstract

Background: We performed a meta-analysis of the efficacy and safety of stem cell therapy as a clinical treatment of knee osteoarthritis. This meta-analysis is expected to provide evidence of the efficacy of stem cell therapy, which is currently controversial, as a conservative treatment for knee osteoarthritis.

Methods: An online search for relevant articles was conducted in the PubMed, EMBASE, and Cochrane Library databases. The search terms were “stem cells” and “osteoarthritis.” We conducted a quality assessment of the included articles and extracted the following indicators: Visual Analogue Scale (VAS) score, Subjective International Knee Documentation Committee (IKDC) score, Western Ontario and McMaster Universities (WOMAC) subscales, and adverse events. The RevMan5.3 software was used for determining effect sizes.

Results: Nine randomized controlled trials involving 339 patients were included. VAS score and IKDC score from baseline to 24 months were improved in the stem cell therapy group compared to those in the control group. However, no significant difference was observed between the 2 groups in IKDC score changes from baseline to 6 and 12 months, as well as in WOMAC-Pain, WOMAC-Stiffness, and WOMAC-Physical Function score changes at each visit point.

Conclusion: Stem cell therapy is certainly superior to traditional treatments in the conservative treatment of KOA; it considerably reduces pain with no obvious additional side effects.

Abbreviations: AD-MSCs = adipose-derived mesenchymal stem cells, BMAC = bone marrow aspirate concentrate, BM-MSCs = bone marrow mesenchymal stromal cells, CI = confidence interval, HA = hyaluronic acid, HTO = high tibial osteotomy, IKDC = International Knee Documentation Committee, IL-1RA = IL-1 receptor antagonist, MSCs = mesenchymal stem cells, PBSC = peripheral blood stem cells, PRG = progenza, PRISMA = Preferred Reporting Items for Systematic Reviews and Meta-Analysis, PRP = platelet-poor plasma, RCTs = randomized controlled trials, SMD = standardized mean difference, VAS = Visual Analogue Scale, WOMAC = Western Ontario and McMaster Universities.

Keywords: knee osteoarthritis, meta-analysis, stem cell therapy

1. Introduction

Knee osteoarthritis is a chronic degenerative bone metabolic disease that commonly occurs in middle-aged and older adults; it affects patients' daily activities and even causes disability.^[1,2] Its

clinical features mainly include cartilage degenerative lesions, with clinical manifestations such as joint swelling, pain, and deformity. Thus, the main therapeutic purposes of knee osteoarthritis are to reduce or eliminate pain, correct joint deformities, and improve joint function through cartilage repair.^[3]

In recent years, replacement of damaged articular cartilage by chondrocytes or cartilage tissue has been considered a potential approach for treating knee osteoarthritis. Studies have shown that it is feasible to induce human pluripotent stem cells to differentiate into chondrocytes; therefore, stem cell therapy has become a new method for local treatment of knee osteoarthritis. For example, mesenchymal stem cells (MSCs) have multi-directional differentiation potential and can be differentiated into osteoblasts and chondrocytes under specific induction conditions in vitro and in vivo, thereby repairing bone and articular cartilage.^[4,5] However, there is still a dispute on the clinical effects of stem cells,^[6–8] for which a multitude of clinical trials and meta-analyses have been conducted.^[9,10]

We herein present a meta-analysis of the controversial efficacy and safety of stem cell therapy as a clinical treatment of knee osteoarthritis. This study is markedly distinguished from previous meta-analyses^[9,10] because it focused on bone marrow MSCs, peripheral blood stem cells, and amniotic fluid free stem cells. In addition, we used updated data from several latest high-

Editor: Eric Bush.

The authors have no conflicts of interests to disclose.

^a Department of neurosurgery, The Second Hospital of Jilin University,

^b Department of pediatrics, The First Hospital of Jilin University, ^c Department of hand surgery, The Second Hospital of Jilin University, ^d College of Mathematics, Jilin University, ^e Orthopaedic Medical Center, The Second Hospital of Jilin University, Changchun, China.

* Correspondence: Meng Xu, The Second Hospital of Jilin University, Changchun, Jilin China (e-mail: xumeng@jlu.edu.cn).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Huang R, Li W, Zhao Y, Yang F, Xu M. Clinical efficacy and safety of stem cell therapy for knee osteoarthritis: A meta-analysis. *Medicine* 2020;99:11(e19434).

Received: 17 September 2019 / Received in final form: 11 December 2019 / Accepted: 5 February 2020

<http://dx.doi.org/10.1097/MD.00000000000019434>

level randomized controlled trials (RCTs).^[11,12] This meta-analysis is expected to provide an evidence of the efficacy of stem cell therapy as a conservative treatment of knee osteoarthritis.

2. Methods

All analyses were based on previous published studies; thus, no ethical approval and patient consent are required.

2.1. Study selection

In accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement,^[13] 2 researchers independently screened the literature, as well as extracted and cross-checked the relevant data. If disagreements occurred, a decision regarding data extraction was made by a third researcher.

2.2. Search strategy

We conducted the search in PubMed (1970-May 2019), Embase (1970-May 2019), and The Cochrane Library (1970-May 2019) databases for relevant articles, with “stem cells” and “osteoarthritis” as search terms. We also manually screened relevant Chinese and English language journals and reference lists to include potential studies. The search strategy for PubMed is detailed herein as an example: (((“Stem Cells”[Mesh]) OR (((((((((((Cell, Stem[Title/Abstract]) OR Stem Cell[Title/Abstract]) OR Progenitor Cells[Title/Abstract]) OR Cell, Progenitor[Title/Abstract]) OR Cells, Progenitor[Title/Abstract]) OR Progenitor Cell[Title/Abstract]) OR Mother Cells[Title/Abstract]) OR Cell, Mother[Title/Abstract]) OR Cells, Mother[Title/Abstract]) OR Mother Cell[Title/Abstract]) OR Colony-Forming Unit[Title/Abstract]) OR Colony Forming Unit[Title/Abstract]) OR Colony-Forming Units[Title/Abstract]) OR Colony Forming Units[Title/Abstract])) AND (((((((((((Osteoarthritis[Title/Abstract]) OR Osteoarthritis[Title/Abstract]) OR Osteoarthroses[Title/Abstract]) OR Arthritis, Degenerative[Title/Abstract]) OR Arthritides, Degenerative[Title/Abstract]) OR Degenerative Arthritides[Title/Abstract]) OR Degenerative Arthritis[Title/Abstract]) OR Osteoarthritis Deformans[Title/Abstract]) OR Polyarthritides[Title/Abstract]) OR Arthritides[Title/Abstract]) OR Polyarthritides[Title/Abstract]) OR Arthritis[Title/Abstract])) OR “Osteoarthritis”[Mesh]).

2.3. Eligibility criteria

The study inclusion criteria included:

- (1) studies involving patients with knee osteoarthritis;
- (2) studies including stem cell therapy as the test group, as well as placebo, hyaluronic acid, and steroid treatments as the control groups;
- (3) RCTs;
- (4) studies that used at least one of the following indicators: Visual Analogue Scale (VAS) score, Western Ontario and McMaster Universities (WOMAC) subscale, International Knee Documentation Committee (IKDC) score, and incidence of adverse events.

Studies were ineligible if they met any of the following conditions:

- (1) studies that used animals or cadavers as research objects;
- (2) studies that were unable to extract or convert valid data;

- (3) retrospective studies, literature reviews, or conference papers with no full text.

2.4. Data extraction

Data were extracted independently by 2 researchers using a predesigned data sheet. Valid data were converted as per the Cochrane Handbook for Systematic Reviews of Interventions,^[14] in the case where standard deviation could not be acquired. If disagreements occurred, the decision regarding data extraction was done by the third reviewer. Each RCT was concurrently assessed with risk of bias.

2.5. Outcome measures

- VAS is a scoring scale that intuitively quantifies the intensity of pain in the knee. A lower score indicates milder pain.
- The WOMAC subscale is a rating scale that assesses the structure, stiffness, and function of the knee in pain. A lower score indicates better knee condition.
- IKDC is a subjective scale for assessing the knee joint. A higher score indicates better symptoms, functions, and physical activities of the knee joint.
- Adverse events refer to treatment-related adverse reactions, including joint effusion, stiffness, and pain.

2.6. Statistical analysis

Statistical analysis was conducted using the RevMan 5.3 software (Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014). The chi-square test was used to assess inter-study heterogeneity. $I^2 > 50\%$ indicated heterogeneity. A random effects model was used; otherwise, a fixed effects model was used. Relative risk and standardized mean difference were used for assessing binary variables and continuous variables, respectively. The 95% confidence interval estimates and hypothesis testing results for each variable were listed in a forest plot. For each endpoint with high heterogeneity, a sensitivity analysis, in which the included studies were removed one at a time, was conducted to screen the source of heterogeneity. A publication bias assessment using a funnel plot was performed if there were no less than 10 studies included.

3. Results

3.1. Literature search

We retrieved 7054 relevant articles, and ultimately included 9 RCTs^[11,12,15–21] involving 399 patients (Fig. 1). In the studies by Kuah,^[12] Lamo-Espinosa,^[16] and Thomas Vangsnæs et al,^[19] there were 2 parallel test groups, namely the high- and low-dose groups, in comparison with the control group. Therefore, for each study mentioned above, we conducted statistical analyses in 2 RCTs: high-dose vs control and low-dose vs control.

3.2. Study characteristics

There were 203 patients in the stem cell therapy group and 196 patients in the control group. The specific features and Jadad scores of the patients^[22,23] are listed in Table 1. The Jadad scale is a 7-point scale that includes random sequence generation, randomized hiding, blind method, withdrawal, and dropout.




Figure 1. Flowchart of literature retrieval.

3.3. Clinical outcomes

3.3.1. VAS. From baseline to 3 months, 4 studies^[12,15,16,20] were included, involving 6 RCTs with 87 patients in the stem cell group and 79 patients in the control group Fig. 2. There was no heterogeneity ($I^2=0\%$) between the studies; thus, the fixed

effects model was used for the analysis. According to Figure 2, SMD (standardized mean difference) = -0.36 , 95% CI (confidence interval) $[-0.67, -0.05]$, and $P=.02$. The VAS score in the stem cell group was significantly lower than that in the control group.

Table 1

Main characteristics of all the eligible studies included in the analysis.

Author	Year	Intervention		Numbers		Follow up (months)		Jadad score
		SCs (dose)	Control	SCs	Control	SCs	Control	
Lee et al ^[11]	2019	AD-MSCs (100 million)	Saline	12	12	6	6	6
Kuah et al ^[12]	2018	PRG (6.7 million/3.9 million)	Placebo	8	4	12	12	7
Shapiro et al ^[15]	2016	BMAC & PRP	Saline	25	25	6	6	6
Lamo-Espinosa et al ^[16]	2016	BM-MSCs & HA (10 million/100 million)	HA	10	10	12	12	7
Vega et al ^[21]	2015	MSCs (40 million)	HA	15	15	12	12	3
Vangness Jr et al ^[19]	2014	MSCs (50 million/150 million)	HA	18	19	24	24	7
Wong et al ^[17]	2013	MSCs & HTO	HTO	28	28	24.8	24.5	6
Saw et al ^[18]	2012	PBSC & HA	HA	25	24	24	24	6
Bhattacharya et al ^[20]	2011	Amniotic fluid	Triamcinolone acetonide	26	26	6	6	5

AD-MSCs = adipose-derived mesenchymal stem cells, BMAC = bone marrow aspirate concentrate, BM-MSCs = bone marrow mesenchymal stromal cells, HA = hyaluronic acid, HTO = high tibial osteotomy, MSCs = mesenchymal stem cells, PBSC = peripheral blood stem cells, PRG = progenza, PRP = platelet-poor plasma.




Figure 2. Forest plot of the change of VAS score. VAS=visual analogue scale score.

From baseline to 6 months, 4 studies^[12,15,16,20] were included, involving 6 RCTs with 87 patients in the stem cell group and 79 patients in the control group. Because there was a high heterogeneity (I²=86%) between the studies, the study by Bhattacharya et al^[20] was removed from the sensitivity analysis, and the I² value was reduced to 0%. The fixed effects model was used. According to Figure 2, SMD=-0.86, 95% CI [-1.21, -0.52], and P<.00001. The VAS score in the stem cell group was significantly lower than that in the control group.

From baseline to 12 months, 3 studies^[12,16,21] were included, involving 5 RCTs with 51 patients in the stem cell group and 43 patients in the control group. There was a low heterogeneity (I²=8%) between the studies, and thus the fixed effects model was used. According to Figure 2, SMD=-0.86, 95% CI [-1.30, -0.43], and P=0.0001. The VAS score in the stem cell group was significantly lower than that in the control group.

3.3.2. WOMAC-Pain. From baseline to 3 months, 2 studies^[12,16] were included, involving four RCTs with 36 patients in the stem cell group and 28 patients in the control group Fig. 3. There was a low heterogeneity (I²=40%) between the studies, and thus the fixed effects model was used. According to Figure 3, SMD=-0.22, 95% CI [-0.73, 0.30], and P=.41. There was no significant difference in WOMAC-Pain score between the groups.

From baseline to 6 months, 2 studies^[12,16] were included, involving four RCTs with 36 patients in the stem cell group and 28 patients in the control group. There was a low heterogeneity (I²=48%) between the studies, and thus the fixed effects model was used. According to Figure 3, SMD=-0.08, 95% CI [-0.59,

0.44], and P=.77. There was no significant difference in WOMAC-Pain score between the groups.

From baseline to 12 months, 3 studies^[12,16] were included, involving 4 RCTs with 43 patients in the stem cell group and 39 patients in the control group. There was no heterogeneity (I²=0%) between the studies, and thus the fixed effects model was used. According to Figure 3, SMD=-0.09, 95% CI [-0.53, 0.36], and P=.70. There was no significant difference in WOMAC-Pain score between the groups.

3.3.3. WOMAC-Stiffness. From baseline to 3 months, 2 studies^[12,16] were included, involving 4 RCTs with 36 patients in the stem cell group and 28 patients in the control group Fig. 4. There was no heterogeneity (I²=0%) between the studies, and the fixed effects model was used. According to Figure 4, SMD=-0.51, 95% CI [-1.02, 0.01], and P=.05. There was no significant difference in WOMAC-Stiffness score between the groups.

From baseline to 6 months, 2 studies^[12,16] were included, involving four RCTs with 36 patients in the stem cell group and 28 patients in the control group. There was a low heterogeneity (I²=36%) between the studies, and the fixed effects model was used. According to Figure 4, SMD=-0.25, 95% CI [-0.76, 0.27], and P=.35. There was no significant difference in WOMAC-Stiffness score between the groups.

From baseline to 12 months, 2 studies^[12,16] were included, involving 4 RCTs with 36 patients in the stem cell group and 28 patients in the control group. There was a low heterogeneity (I²=9%) between the studies, and the fixed effects model was used.




Figure 3. Forest plot of the change of WOMAC-Pain score. WOMAC=Western Ontario and McMaster Universities subscore.

According to Figure 4, SMD = -0.46, 95% CI [-0.98, 0.05], and P = .08. There was no significant difference in WOMAC-Stiffness score between the groups.

3.3.4. WOMAC-Function. From baseline to 3 months, 2 studies^[12,16] were included, involving 4 RCTs with 36 patients

in the stem cell group and 28 patients in the control group Fig. 5. There was no heterogeneity (I² = 0%) between the studies, and the fixed effects model was used. According to Figure 5, SMD = 0.15, 95% CI [-0.35, 0.66], and P = .55. There was no significant difference in WOMAC-Function score between the groups.




Figure 4. Forest plot of the change of WOMAC-Stiffness score. WOMAC=Western Ontario and McMaster Universities subscore.




Figure 5. Forest plot of the change of WOMAC-Function. WOMAC=Western Ontario and McMaster Universities subscore.

From baseline to 6 months, 2 studies^[12,16] were included, involving four RCTs with 36 patients in the stem cell group and 28 patients in the control group. There was a low heterogeneity (I²=29%) between the studies, and the fixed effects model was used. According to Figure 5, SMD=0.43, 95% CI [-0.09, 0.95], and P=.1. There was no significant difference in WOMAC-Function score between the groups.

From baseline to 12 months, 2 studies^[12,16] were included, involving 4 RCTs with 36 patients in the stem cell group and 28

patients in the control group. There was no heterogeneity (I²=0%) between the studies, and the fixed effects model was used. According to Figure 5, SMD=0.18, 95% CI [-0.33, 0.68], and P=.49. There was no significant difference in WOMAC-Function score between the groups.

3.3.5. IKDC. From baseline to 6 months, 2 studies^[17,18] were included, involving 2 RCTs with 53 patients in the stem cell group and 52 patients in the control group Fig. 6. There was a




Figure 6. Forest plot of the change of IKDC score. IKDC=International Knee Documentation Committee.




Figure 7. Forest plot of the change of adverse events.

high heterogeneity ($I^2=51\%$) between the studies, and the random effects model was used. According to Figure 6, SMD = 0.16, 95% CI [-0.39, 0.72], and $P = .56$. There was no significant difference in IKDC score between the groups.

From baseline to 12 months, 2 studies^[17,18] were included, involving 2 RCTs with 53 patients in the stem cell group and 52 patients in the control group. There was a high heterogeneity ($I^2=83\%$) between the studies, and the random effects model was used. According to Figure 6, SMD = 0.36, 95% CI [-0.58, 1.31], and $P = .45$. There was no significant difference in IKDC score between the groups.

From baseline to 24 months, 2 studies^[17,18] were included, involving 2 RCTs with 53 patients in the stem cell group and 52 patients in the control group. There was a high heterogeneity ($I^2=75\%$) between the studies, and the random effects model was used. According to Figure 6, SMD = 0.53, 95% CI [-0.25, 1.32], and $P = .18$. There was no significant difference in IKDC score between the groups.

3.3.6. Adverse events. There were 2 included studies,^[11,12] involving three RCTs with 28 patients in the stem cell group and 20 patients in the control group Fig. 7. There was a high heterogeneity ($I^2=73\%$) between the studies, and thus the random effects model was used. According to Figure 7, SMD = 1.54, 95% CI [0.57, 4.19], and $P = .40$. There was no significant difference in incidence of adverse events between the groups.

4. Discussion

4.1. Key findings

Changes in VAS and IKDC scores from baseline to 24 months were superior in the stem cell group than in the control group, whereas there were no statistical differences in the changing trend of other indicators between the 2 groups, including the changes in IKDC scores at 6 months, IKDC score at 12 months, WOMAC-Pain score, WOMAC-Stiffness score, WOMAC-Function score, WOMAC-Pain score, and incidence of adverse events.

Pain relief is key to treating knee osteoarthritis, with VAS scores as an important endpoint for pain assessment. Kuah et al^[12] found that relative to placebo, stem cell therapy considerably relieved pain at 3, 6, and 12 months after treatment. This conclusion has been confirmed in our meta-analysis. We found that the VAS scores in the stem cell group were significantly reduced at each visit point. Inflammatory response is known as one of the causes of pain. MSCs can release anti-inflammatory factors, thereby relieving pain. Lamo-Espinosa et al^[16] believed that stem cells have a paracrine function and their anti-inflammatory properties contribute to pain relief. In addition, studies have found that in an acute renal failure model, MSCs can

promote recovery of renal function by releasing anti-inflammatory factors and inhibiting production of pro-inflammatory cytokines, such as interleukin-1 β , tumor necrosis factor, and interferon- γ .^[24] Similar findings were observed in a pulmonary fibrosis model, in which MSCs release IL-1 receptor antagonist (IL-1RA) to inhibit interleukin-1 α -producing T cells and TNF-producing macrophages, indicating that MSCs have anti-inflammatory properties.^[25] WOMAC-Pain scores showed no statistical difference between the 2 groups at each visit point, but these data regarding WOMAC-Pain score were obtained only from 3 studies;^[12,16,21] therefore, further studies with larger sample sizes are warranted to verify these findings.

Functional improvement of the knee joint is one of the ultimate purposes of knee osteoarthritis treatment. In this study, we used WOMAC-Stiffness, WOMAC-Function, and IKDC scores to comprehensively assess knee joint function. Statistical analysis results showed that there was no significant difference between the 2 groups in IKDC scores at 6 and 12 months, as well as in WOMAC-Stiffness and WOMAC-Function scores at each visit point. Studies have found that mesenchymal stem cell implantation achieves better outcomes in patients with grade 3 knee osteoarthritis than those in patients with grade 4 knee osteoarthritis.^[26] We thus concluded that treatment with MSCs are effective in preventing or limiting the progression of knee osteoarthritis at the early stage. In the studies by Kuah^[12] and Lamo-Espinosa et al,^[16] patients with grade 3 osteoarthritis or higher accounted for 75% and over 80% of the total patients, respectively. Most patients developed knee osteoarthritis at the middle and late stages, for whom treatment with MSCs had no significant efficacy and was not conducive to functional recovery. Moreover, in most tissue engineering methods, MSCs are combined with cell scaffolds containing chondrogenic growth factors to form fully functional hyaline cartilage. Such a regimen is commonly used in small-animal models of surgically induced cartilage or osteochondral defects, but cannot be used for repairing large-area cartilage defects associated with knee osteoarthritis.^[27] In addition, Centeno,^[28] Emadedin,^[29] and Vangsnest et al^[19] pointed out that treatment with approximately 2×10^7 stem cells can afford good clinical results. Kuah^[12] and Lamo-Espinosa et al^[16] reported that a stem cell dose of lower or higher than 2×10^7 may also impact the therapeutic efficacy of stem cells. In addition, the change in IKDC score at 24 months was higher in the stem cell group than in the control group; however, these data were extracted from only 2 studies. Therefore, further investigation with a larger sample size is warranted.

For adverse events, we sent an e-mail to the authors of the relevant research^[18,19,21] to obtain data on the number of patients who experienced treatment-related adverse events,

including arthralgia, joint effusion, and joint stiffness, in both the stem cell and control groups. Because of the lack of response from the other studies, only 2 studies^[11,12] were included, involving 3 RCTs. There were no statistical differences in adverse events between the 2 groups, indicating that stem cell treatment has no obvious side effects. A study addressing 87 patients with lupus erythematosus^[30] showed no adverse events associated with transplantation after 4 years of follow-up. Similarly, no graft-related adverse events occurred in many patients undergoing stem cell therapy for other diseases.^[31–35] These findings indicated that the human body has good tolerance to MSCs, and that stem cell treatment has no significant side effects.

4.2. Limitations

Differences in the original RCT protocols led to insufficient representation of some outcome indicators. Thus, high-quality, large-scale RCTs are required to verify our findings. In addition, there was no uniform standard in the preparation and use of stem cells, which may cause certain heterogeneity.

5. Conclusion

Compared to traditional methods, stem cell treatment has a certain superiority as a conservative treatment of knee osteoarthritis, in terms of markedly reducing pain without inducing side effects.

Author contributions

Conceptualization: Rui Huang.

Data curation: Rui Huang, Wei Li, Ying Zhao.

Formal analysis: Rui Huang, Wei Li, Ying Zhao, Fan Yang.

Investigation: Rui Huang, Wei Li, Ying Zhao.

Methodology: Rui Huang, Wei Li, Fan Yang.

Project administration: Meng Xu.

Software: Rui Huang, Wei Li, Ying Zhao, Fan Yang.

Supervision: Rui Huang, Wei Li, Ying Zhao, Meng Xu.

Validation: Ying Zhao, Fan Yang, Meng Xu.

Visualization: Rui Huang, Wei Li, Fan Yang.

Writing – original draft: Rui Huang, Wei Li.

Writing – review & editing: Ying Zhao, Meng Xu.

References

- [1] Krasnokutsky S, Samuels J, Abramson SB. Osteoarthritis in 2007. *Bulletin of the Nyu Hospital for Joint Diseases* 2007;65:222–8.
- [2] Clouet J, Vinatier C, Merceron C, et al. From osteoarthritis treatments to future regenerative therapies for cartilage. *Drug Discovery Today* 2009;14:913–25.
- [3] Abramson SB, Attur M. Developments in the scientific understanding of osteoarthritis. *Arthritis Res Ther* 2009;11:227–327.
- [4] Wakitani S, Imoto K, Yamamoto T, et al. Human autologous culture expanded bone marrow mesenchymal cell transplantation for repair of cartilage defects in osteoarthritic knees. *Osteoarthritis Cartilage* 2002;10:199–206.
- [5] Davatchi F, Abdollahi BS, Mohyeddin M, et al. Mesenchymal stem cell therapy for knee osteoarthritis. Preliminary report of four patients. *Int J Rheum Dis* 2016;14:211–5.
- [6] Sotiropoulou PA, Perez SA, Maria S, et al. Characterization of the optimal culture conditions for clinical scale production of human mesenchymal stem cells. *Stem Cells* 2010;24:462–71.
- [7] J Mary M, Kenneth D, Stephen B, et al. Reduced chondrogenic and adipogenic activity of mesenchymal stem cells from patients with advanced osteoarthritis. *Arthritis Rheum* 2010;46:704–13.
- [8] Banfi A, Bianchi G, Notaro R, et al. Replicative aging and gene expression in long-term cultures of human bone marrow stromal cells. *Tissue Eng* 2002;8:901–10.
- [9] Yubo M, Yanyan L, Li L, et al. Clinical efficacy and safety of mesenchymal stem cell transplantation for osteoarthritis treatment: a meta-analysis. *Plos One* 2017;12:e0175449.
- [10] Xia P, Wang X, Lin Q, et al. Efficacy of mesenchymal stem cells injection for the management of knee osteoarthritis: a systematic review and meta-analysis. *Int Orthop* 2015;39:2363–72.
- [11] Lee WS, Kim HJ, Kim KI, et al. Intra-articular injection of autologous adipose tissue-derived mesenchymal stem cells for the treatment of knee osteoarthritis: a phase IIb, randomized, placebo-controlled clinical trial. *Stem Cells Transl Med*. 2019;8:504–11.
- [12] Kuah D, Sivell S, Longworth T, et al. Safety, tolerability and efficacy of intra-articular Progenza in knee osteoarthritis: a randomized double-blind placebo-controlled single ascending dose study. *J Transl Med* 2018;16:49.
- [13] Panic N, Leoncini E, De Belvis G, et al. Evaluation of the endorsement of the preferred reporting items for systematic reviews and meta-analysis (PRISMA) statement on the quality of published systematic review and meta-analyses. *Stefania Boccia. European Journal of Public Health* 2013;23(suppl_1):e83138.
- [14] Jüni P, Altman DG, Egger M. Assessing the quality of randomized controlled trials. "Systematic reviews in health care: meta-analysis in context. 2001;87–108.
- [15] Shapiro SA, Kazmerchak SE, Heckman MG, et al. A prospective, single-blind, placebo-controlled trial of bone marrow aspirate concentrate for knee osteoarthritis. *Am J Sports Med* 2017;45:82–90.
- [16] Lamo-Espinosa JM, Mora G, Blanco JF, et al. Intra-articular injection of two different doses of autologous bone marrow mesenchymal stem cells versus hyaluronic acid in the treatment of knee osteoarthritis: multicenter randomized controlled clinical trial (phase I/II). *J Transl Med* 2016;14:246.
- [17] Wong KL, Lee KB, Tai BC, et al. Injectable cultured bone marrow-derived mesenchymal stem cells in varus knees with cartilage defects undergoing high tibial osteotomy: a prospective, randomized controlled clinical trial with 2 years' follow-up. *Arthroscopy* 2013;29:2020–8.
- [18] Khay-Yong S, Adam A, Caroline SYJ, et al. Articular cartilage regeneration with autologous peripheral blood stem cells versus hyaluronic acid: a randomized controlled trial. *Arthroscopy* 2013;29: 684–94.
- [19] C Thomas V, Jack F, Joel B, et al. Adult human mesenchymal stem cells delivered via intra-articular injection to the knee following partial medial meniscectomy: a randomized, double-blind, controlled study. *J Bone Joint Surg Am* 2014;96:90–8.
- [20] Bhattacharya N. Clinical use of amniotic fluid in osteoarthritis: a source of cell therapy. *Transplantation* 2011;90:395–403.
- [21] Aurelio V, Miguel Angel MF, Francisco DC, et al. Treatment of knee osteoarthritis with allogeneic bone marrow mesenchymal stem cells: a randomized controlled trial. *Transplantation* 2015;99:1681–90.
- [22] Moher D, Jadad AR, Nichol G, et al. Assessing the quality of randomized controlled trials: an annotated bibliography of scales and checklists. *Control Clin Trials* 1995;16:62–73.
- [23] Jadad AR, Rennie D. The randomized controlled trial gets a middle-aged checkup. *J Am Med Assoc* 1998;279:319–20.
- [24] Tögel F, Hu Z, Weiss K, et al. Administered mesenchymal stem cells protect against ischemic acute renal failure through differentiation-independent mechanisms. *Am J Physiol Renal Physiol* 2005;289:F31–42.
- [25] Ortiz LA, Maria D, Cheryl F, et al. Interleukin 1 receptor antagonist mediates the antiinflammatory and antifibrotic effect of mesenchymal stem cells during lung injury. *Proc Natl Acad Sci U S A* 2007;104: 11002–7.
- [26] Koh YG, Choi YJ. Infrapatellar fat pad-derived mesenchymal stem cell therapy for knee osteoarthritis. *Knee* 2012;19:902–7.
- [27] Grässel S, Lorenz J. Tissue-engineering strategies to repair chondral and osteochondral tissue in osteoarthritis: use of mesenchymal stem cells. *Curr Rheumatol Rep* 2014;16:452.
- [28] Centeno CJ, Busse D, Kisdaj J, et al. Increased knee cartilage volume in degenerative joint disease using percutaneously implanted, autologous mesenchymal stem cells. *Pain Phys* 2008;11:343–53.
- [29] Emadedin M, Aghdami N, Taghiyar L, et al. Intra-articular injection of autologous mesenchymal stem cells in six patients with knee osteoarthritis. *Arch Iranian Med* 2012;15:422.
- [30] Wang D, Zhang H, Liang J, et al. Allogeneic mesenchymal stem cell transplantation in severe and refractory systemic lupus erythematosus: 4 years of experience. *Cell Transplant* 2013;22:2267–77.

- [31] Koç ON, Gerson SL, Cooper BW, et al. Rapid hematopoietic recovery after confusion of autologous-blood stem cells and culture-expanded marrow mesenchymal stem cells in advanced breast cancer patients receiving high-dose chemotherapy. *J Clin Oncol* 2000;18:307.
- [32] Lazarus HM, Koc ON, Devine SM, et al. Cotransplantation of HLA-identical sibling culture-expanded mesenchymal stem cells and hematopoietic stem cells in hematologic malignancy patients. *Biol Blood Marrow Transplant* 2005;11:389–98.
- [33] Hare JM, Fishman JE, Gerstenblith G, et al. Comparison of allogeneic vs autologous bone marrow–derived mesenchymal stem cells delivered by transcatheter injection in patients with ischemic cardiomyopathy: the POSEIDON randomized trial. *JAMA* 2012;308:2369–79.
- [34] Bernardo ME, Pagliara D, Locatelli F. Mesenchymal stromal cell therapy: a revolution in regenerative medicine? *Bone Marrow Transplant* 2012;47:164–71.
- [35] Wang P, Li Y, Huang L, et al. Effects and safety of allogeneic mesenchymal stem cell intravenous infusion in active ankylosing spondylitis patients who failed NSAIDs: a 20-week clinical trial. *Cell Transplant* 2014;23:1293–303.