



# What Makes Nutrition Research So Difficult to Conduct and Interpret?

Mara Z. Vitolins<sup>1</sup> and Talsi L. Case<sup>2</sup>

<sup>1</sup>Department of Epidemiology and Prevention, Wake Forest School of Medicine, Winston-Salem, NC; <sup>2</sup>Nutrition and Foods Bachelor of Science Dietetics Program [student], Appalachian State University, Boone, NC

Conducting and interpreting nutrition research involves consideration of the research question, study design, wide variability of nutrients in foods and dietary patterns, food environment, approaches used to collect and analyze dietary data, and manner in which results are reported. This article reviews all of these considerations with regard to diabetes-related nutrition research.

Nutrition researchers are trained to examine the complex interplay between foods eaten and health and disease in individuals or populations. Given the huge potential impact of diet on a person's health and the fact that everyone eats, it is no surprise that studies in human nutrition are crucially important. However, nutrition research has its challenges. To fully appreciate the many challenges surrounding nutrition research, it is important to understand some of the key elements involved, including research designs, the complexity of the food environment, and approaches to collecting and analyzing dietary data.

Nutrition research conducted to examine the role of diet in managing diabetes is particularly challenging because it involves assessing and intervening in patients' lifestyle and habits, which are influenced by human nature; a food environment that is ever-changing; and data that are largely self-reported. Another challenge is how nutrition research results are described; unfortunately, nutrition research results are often simplified and misinterpreted. Entire books have described in great detail the fundamental approaches to and challenges of conducting nutrition research (1,2). Such a detailed discussion is beyond the scope of this article, which rather aims to create awareness of a handful of these issues.

## Research Study Designs Are Important

When designing a nutrition research study, it is crucial to determine the best study design to answer the research question (1,2). Several research designs are commonly used to assess the impact of a dietary intervention on health and disease. Some of these include tightly controlled

feeding studies, randomized clinical trials, and observational studies. Each design has advantages and disadvantages.

A feeding trial involves providing study participants with all meals, snacks, and beverages to control what they consume. Such trials provide the most control over participants' dietary intake. Provided meals are carefully prepared, ingredients are weighed and measured, and participants are instructed to eat all the food they are provided (or to bring back what they did not eat). As an example, Bell et al. (3,4) used single-meal feeding trials to examine insulin dosing for protein and fat intake in participants with type 1 diabetes.

Feeding trials are expensive, and special facilities are needed to prepare and provide food for the participants. Feeding trials are usually short-term, not only because of the costs involved, but also because participants have to greatly modify their lives to adhere to eating only the foods and beverages provided by the trial. These trials usually enroll people who are quite different from the general public. For example, if a nutrition researcher wanted to study the impact of different amounts of dietary fiber on A1C levels, and wanted to include women with prediabetes who are 30–40 years of age, recruitment could prove to be difficult, and retaining participants could be even more challenging. The participants would need to be in the trial for at least 3 months to get meaningful changes in A1C; however, women in this age-group are likely to be married, have children, and be employed. To participate in such a trial, they would have to be willing to avoid eating any of the food prepared for the rest of their family, they would not be able share their

Corresponding author: Mara Z. Vitolins, [mvtolin@wakehealth.edu](mailto:mvtolin@wakehealth.edu)

<https://doi.org/10.2337/ds19-0077>

©2020 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

trial-provided food with anyone else, and they would need to visit the research unit nearly every day to pick up meals.

The burden on participants enrolled in feeding trials is great, and the risk for nonadherence increased as the trial goes on. If participants do not strictly adhere to eating the foods they are provided, the findings of the trial can be seriously affected. Yet, it is difficult to precisely determine participants' degree of adherence because they might eat foods other than those they were provided, share their foods with others, or not follow the protocol requiring them to bring back any foods they do not eat.

Feeding trials are not appropriate for answering questions about longer-term dietary exposures. Many diseases develop over time, so fully examining whether a particular dietary pattern affects health requires a long follow-up period and a large sample size. Randomized controlled intervention trials have been conducted to examine the effects of long-term dietary modification (5–7). Although these trials are not feeding trials, they still require a large budget and come with their own complexities. Research teams who have been able to conduct long-term dietary modification trials have relied on many different research methods and tools. Having close, ongoing contact with participants, carefully monitoring their adherence, and asking them to self-monitor what they eat are just a few of the key components of such trials.

These trials are labor-intensive for both researchers and participants, and in most cases, adherence has been shown to decrease after several months. Participant adherence to the intervention (and the control condition) is essential, but not always realized. Although nonadherence to protocol-specified behaviors is not a problem specific to nutrition research, the act of selecting, preparing, and eating foods is complex and repeated multiple times each day, providing many opportunities for nonadherence to the diet modification specified by the trial.

The Diabetes Prevention Program (DPP) trial is a good example of a large, randomized controlled intervention trial (8). Registered dietitians instructed participants in the intervention group to follow a healthful eating pattern, but meals were not provided. Participants had to learn approaches to reducing their fat and calorie intake to achieve a weight loss goal of 7% of baseline body weight and then to maintain their weight loss (8). They were required to self-monitor their behaviors, and dietitians reviewed this documentation regularly and provided constructive feedback. Participants attended weekly sessions during the first 24 weeks of the study. However, during the maintenance phase, contacts with the study team decreased to about twice per month. DPP participants had to remain adherent over a long period

of time, so when contact frequency decreased, the research team had to work even more diligently to assist participants in meeting their study goals. If they had not done so, it would have been impossible to test the study hypothesis that lifestyle intervention would delay or prevent the development of type 2 diabetes in people with impaired glucose tolerance who were at high risk for developing the disease.

Observational studies have been used to track dietary intake in large numbers of participants and can be used to track such data over many years. Participants do not have to change their dietary behaviors to participate in an observational study. These studies are not as expensive to conduct, and the burden on participants and investigators is much lower.

Observational studies are not carefully controlled like clinical trials, so their results may be less reliable. Observational studies compare individuals who self-select to consume either a healthful or unhealthful diet, and the two groups may differ in other characteristics that could influence health (9,10). Nonetheless, such studies are vitally important for identifying possible connections between diet and health that can be further tested in randomized clinical trials.

## **Food Is Complicated**

Nutrition researchers must be aware of and account for numerous food-related issues when collecting and analyzing dietary data. A few such issues are described below.

Foods and food products are available in different varieties, brands, and flavors. There are many options of the same type of food, yet the ingredients in each option may differ in ways that matter greatly to nutrition researchers. One brand of microwavable popcorn may contain butter, whereas another may contain buttery flavoring (butter-flavored coconut oil).

How food is cooked can change its nutrient profile. If vegetables are eaten fresh, many of their nutrients are preserved; however, if vegetables are boiled, many water-soluble nutrients can be lost. Many nutrients, including vitamin C, thiamin, and folic acid, are sensitive to temperature (11).

Where a food is prepared can make a substantial difference in the nutrient content because the ingredients used in their preparation can vary widely. Home-cooked foods versus restaurant-prepared foods can greatly differ in their nutrient composition.

Seasonal variations can make a difference in the nutrient content of fruits and vegetables. Also, those purchased directly from local farms have different levels of nutrients than those picked before they have ripened and been transported many miles to grocery stores.

The food environment is ever-changing, and food ingredients will continue to change, although consumers may not notice these changes at all. For example, until relatively recently, synthetic sources of *trans* fats were ubiquitous in bakery products as an emulsifier to improve creaming and reduce the rancidity of frying oil (12). Now, artificial *trans* fats are no longer contained in bakery products, or they have been greatly reduced (13). If a researcher wanted to examine the intake of cookies over time on health outcomes, using data overlapping this important change in the *trans* fat content of the cookies would greatly complicate this evaluation.

### People Are Also Complicated

People vary in many ways, including by sex, race/ethnicity, BMI, economic status, metabolic rate, food preferences, exercise patterns, and fitness levels, among others. All of these differences could affect what study participants eat, how they metabolize what they eat, and how much they remember about what they eat.

The ability to assess dietary intake to explore diet-disease connections is essential to nutrition research but remains problematic for a number of reasons. Dietary data are typically obtained from participants who self-report what they eat (1,14). Common approaches used to collect dietary data include food records, food frequency questionnaires, and 24-hour recalls (1,14). To complete food records, participants are asked to carefully describe everything they eat or drink over a period of a few days, typically 3–4 days. Food records are burdensome to complete because participants must weigh and measure all of their foods and beverages and write all of this information down. To complete food frequency questionnaires, participants are asked to reflect on what they have typically eaten during the past 6 or 12 months, and it is known that participant recall is fallible. The 24-hour recall method collects dietary intake over a much shorter period of time, but participants can misreport what they eat.

Such misreporting can vary by participants' personal characteristics. For example, a number of studies have shown that women and individuals who are heavier tend to underreport foods consumed (15–18). Underreporting of food intake by participants may reflect social desirability (i.e., not wanting to be judged by what they eat) (16,17). Underreporting may also reflect an inability to accurately estimate how much they actually consume. Extensive research has been conducted to identify approaches to mathematically adjust for misreported intake. Unfortunately, these corrections are not a perfect solution to fully address the problem of misreported food intake. Biomarkers have been studied to assist in the assessment of foods consumed. However, they carry with them a number of different, but still complicating, issues, including individual differences in nutrient absorption (19).

### Food and Nutrition Databases Have Limitations

Food and nutrition databases are used to convert a consumed food to the specific nutrients contained in the food (20). These databases contain information about foods' sources, profiles, nutrients, and dietary components.

It is not hard to imagine the complexity and difficulty involved in attempting to capture all of the different types and varieties of foods available to the public in one database. Each nutrient contained in each food is quantified. Significant effort goes into annually updating the content of food and nutrient databases as new foods are introduced to the marketplace almost daily. The databases are enormous because the intention is usually to include as many foods (and their variations) as possible. For example, there are more than 40 chicken soup options from which to choose in the U.S. Department of Agriculture/Agricultural Research Service Food and Nutrient Database for Dietary Studies (21).

Food and nutrient databases cannot include every component of every food. For example, nutrients are not the only components contained in foods, and the nonnutritive components in foods are usually not included in these databases. Vegetables and fruits contain many nonnutritive compounds, including dithiolthiones, monoterpenes, indoles sterols, and sulfhydryls. Although these components are not nutritional, they are in the foods research participants eat, and they may be important to health.

It is also known that food packaging may have an impact on health. Bisphenol A (BPA) is a good example of a non-nutritive but important exposure that was linked to food packaging. BPA is considered an endocrine disruptor, and the primary source of exposure to BPA is through the diet (22). BPA has been known to leach into food from internal epoxy resin coatings on cans used for canning foods such as soups and vegetables. BPA is not included in food and nutrition databases.

### Reporting of Nutrition Research Findings Can Be Confusing

Once a research study is published, its results are likely to be picked up by the lay press. Such reporting is where a good deal of nutrition confusion is created. Nutrition is a topic of great interest to people, particularly those who have been diagnosed with diabetes, who often inquire about which diet is best to help them achieve target glucose levels or to lose or maintain body weight. Where they get answers to such questions matters; unfortunately, most people learn about nutrition research findings in the popular press or rely on the Internet for nutrition information. Although a wealth of nutrition information is available, some of it may be unreliable.

The online newsletter "Obesity and Energetics Offerings," from the Indiana School of Public Health in Bloomington and

the University of Alabama–Birmingham, provides some excellent examples of headlines that have failed to accurately describe the results of nutrition studies. The newsletter’s “Headline vs. Study” section on 11 October 2019 (23), for example, provided a link to an article in *Women’s Health* magazine titled, “Snacking on Nuts Found to Help Prevent Weight Gain” (24), followed by a link to the actual research report on which the article was based. The study was not a randomized controlled trial, the nuts were not provided to participants, and the participants self-reported both their nut intake and their weight. Although consumers reading the magazine headline might have thought that snacking on nuts would help them control their weight, all the research really found was an association—not necessarily causal—between self-reported nut intake and self-reported weight (25).

Similarly, data from the Women’s Health Initiative study cohort were used to explore links between chocolate consumption and the risk of diabetes (26). The authors cautiously presented their finding of an inverse relationship between chocolate intake and incident diabetes at moderate levels of chocolate consumption. However, press reports about the study simply suggested that eating chocolate would stave off diabetes (27).

Because patients may not be able to interpret the results of nutrition-related studies they see reported in the news, health care providers should be prepared to discuss these issues when patients inquire. Most Americans likely do not understand the nuances of such research or how links between foods or diets and chronic disease outcomes are stronger when supported by consistent evidence from basic science research, feeding studies, observational cohort studies, and randomized controlled clinical trials. People do not always realize that findings from one study cannot stand alone and that results from multiple studies are usually needed to create the robust evidence necessary for nutrition recommendations. Health care providers can help patients better understand nutrition research findings reported in the popular media by asking them to think about how many times they have read or heard about so-called “miracle foods.”

Importantly, when patients ask questions about nutrition-related news articles, they are opening the door to discussing their own dietary habits. Health care providers might consider providing simple suggestions for improving dietary intake (e.g., drink water instead of sugary beverages or eat vegetables at every meal), referral to a dietitian, and information about Internet sites that support evidence-based dietary recommendations.

To address conflicting data and nutrition recommendations, the American Diabetes Association (ADA) routinely reviews diabetes-related nutrition research and determines whether the evidence is strong enough to issue guidelines or recommend changes to the standards of care. The recently published, “Nutrition Therapy for Adults With Diabetes or

Prediabetes: A Consensus Report” (28), for example, summarized the evidence supporting a new recommendation acknowledging that several different dietary patterns are acceptable for people who have prediabetes.

## Conclusion

Everyone eats, making diet one of the most common exposures possibly influencing health and disease outcomes. Nutrition research is complicated by numerous factors inherent in this area of inquiry. Inaccurate reporting in the lay press can further complicate things and create confusion in the public. The ADA reviews data and provides consensus reports and clinical practice guidelines to reduce the confusion. Studies such as the DPP illustrate that nutrition research has enormous potential for preventing and improving the management of diabetes and improving overall health. Nutrition, with its myriad health implications, is an exciting, challenging, and ever-evolving area of research.

## DUALITY OF INTEREST

No potential conflicts of interest relevant to this article were reported.

## AUTHOR CONTRIBUTIONS

M.Z.V. researched data and wrote the manuscript. T.L.C. researched data and reviewed and edited the manuscript. M.Z.V. is the guarantor of this work and, as such, had full access to all the data and takes responsibility for the integrity of the information presented.

## REFERENCES

1. Van Horn L, Beto J, Eds. *Research: Successful Approaches in Nutrition and Dietetics*. 4th ed. Chicago, IL, Academy of Nutrition and Dietetics, 2019
2. Lovegrove J, Hodson L, Sharma S, Lanham-New S, Eds. *Nutrition Research Methodologies*. 1st ed. Hoboken, NJ, John Wiley & Sons, 2015
3. Bell KJ, Fio CZ, Twigg S, et al. Amount and type of dietary fat, postprandial glycemia, and insulin requirements in type 1 diabetes: a randomized within-subject trial. *Diabetes Care* 2020;43:59–66
4. Bell KJ, Toschi E, Steil GM, Wolpert HA. Optimized mealtime insulin dosing for fat and protein in type 1 diabetes: application of a model-based approach to derive insulin doses for open-loop diabetes management. *Diabetes Care* 2016;39:1631–1634
5. Knowler WC, Barrett-Connor E, Fowler SE, et al.; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403
6. Prentice RL, Caan B, Chlebowski RT, et al. Low-fat dietary pattern and risk of invasive breast cancer: the Women’s Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006; 295:629–642
7. Appel LJ, Moore TJ, Obarzanek E, et al.; DASH Collaborative Research Group. A clinical trial of the effects of dietary patterns on blood pressure. *N Engl J Med* 1997;336:1117–1124
8. Diabetes Prevention Program (DPP) Research Group. The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care* 2002;25:2165–2171
9. Djoussé L, Petrone AB, Hickson DA, et al. Egg consumption and risk of type 2 diabetes among African Americans: the Jackson Heart Study. *Clin Nutr* 2016;35:679–684

10. McNaughton SA, Mishra GD, Brunner EJ. Dietary patterns, insulin resistance, and incidence of type 2 diabetes in the Whitehall II Study. *Diabetes Care* 2008;31:1343–1348
11. Bureau S, Mouhoubi S, Touloumet L, et al. Are folates, carotenoids and vitamin C affected by cooking? Four domestic procedures are compared on a large diversity of frozen vegetables. *LWT—Food Science and Technology* 2015;64:735–741
12. Eckel RH, Borra S, Lichtenstein AH, Yin-Piazza SY; Trans Fat Conference Planning Group. Understanding the complexity of trans fatty acid reduction in the American diet: American Heart Association Trans Fat Conference 2006: report of the Trans Fat Conference Planning Group. *Circulation* 2007;115:2231–2246
13. Federal Register. Final determination regarding partially hydrogenated oils. 2015;80:34650–34670
14. Nutritools. Dietary assessment tools. Available from <https://www.nutritools.org/tools>. Accessed 15 February 2020
15. Hebert JR, Clemow L, Pbert L, Ockene IS, Ockene JK. Social desirability bias in dietary self-report may compromise the validity of dietary intake measures. *Int J Epidemiol* 1995;24:389–398
16. Taren DL, Tobar M, Hill A, et al. The association of energy intake bias with psychological scores of women. *Eur J Clin Nutr* 1999; 53:570–578
17. Johnson RK, Goran MI, Poehlman ET. Correlates of over- and underreporting of energy intake in healthy older men and women. *Am J Clin Nutr* 1994;59:1286–1290
18. Tomoyasu NJ, Toth MJ, Poehlman ET. Misreporting of total energy intake in older men and women. *J Am Geriatr Soc* 1999;47:710–715
19. Picó C, Serra F, Rodríguez AM, Keijer J, Palou A. Biomarkers of nutrition and health: new tools for new approaches. *Nutrients* 2019;11:e1092
20. U.S. Department of Agriculture, Agricultural Research Service. What we eat in America: food categories 2015–2016. Available from <https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group>. Accessed 2 December 2019
21. U.S. Department of Agriculture, Agricultural Research Service. What's in the food you eat search tool: find a food. Available from [https://reedir.arsnet.usda.gov/codesearchwebapp/\(S\(ijhssteqdbwyizndbb3ib2co\)\)/CodeSearch.aspx](https://reedir.arsnet.usda.gov/codesearchwebapp/(S(ijhssteqdbwyizndbb3ib2co))/CodeSearch.aspx). Accessed 2 December 2019
22. Sonavane M, Gassman NR. Bisphenol A co-exposure effects: a key factor in understanding BPA's complex mechanism and health outcomes. *Crit Rev Toxicol* 2019;49:371–386
23. Obesity and Energetics. Obesity and Energetics Offerings 10/11/19. Available from <https://www.obesityandenergetics.org/weeklyofferings/101119>. Accessed 1 December 2019
24. Williamson L. Snacking on nuts found to help prevent weight gain. *Womens Health* 29 September 2019. Available from <https://www.womenshealth.com.au/nut-intake-weight-gain-prevention>. Accessed 1 December 2019
25. Liu X, Li Y, Guasch-Ferré M, et al. Changes in nut consumption influence long-term weight change in US men and women. *BMJ Nutrition, Prevention & Health* 2019;2:90–99
26. Greenberg JA, Manson JE, Tinker L, et al. Chocolate intake and diabetes risk in postmenopausal American women. *Eur J Clin Nutr* 2017;71:1088–1093
27. Chocolate prevents diabetes! *Womens World* 13 July 2017. Available from <https://www.pressreader.com/usa/womans-world/20170731/281625305345336>. Accessed 2 March 2020
28. Evert AB, Dennison M, Gardner CD, et al. Nutrition therapy for adults with diabetes or prediabetes: a consensus report. *Diabetes Care* 2019;42:731–754