



Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>



EDITORIAL

SARS CoV-2 (COVID-19): lessons to be learned by Brazilian Physical Therapists



The current severe acute respiratory syndrome coronavirus 2 (SARS CoV-2) pandemic, also known as Coronavirus Disease 2019 (COVID-19), puts us at the forefront of scenarios with serious consequences for health systems in Brazil and in the world.¹ Current viral interstitial pneumonia has resulted in severe hypoxemic respiratory failure, which has resulted in overcrowding in intensive care units (ICUs), shortage of equipment and personnel, and significant mortality, especially in populations at risk who have chronic health conditions.² In most countries, the volume of patients has exceeded critical care capacities, with a shortage of personal protection equipment, multi-professional teams, and mechanical ventilators dominating the discussions in local hospitals and in the media.^{3,4}

While COVID-19 brought several challenges to the overall health care system, it also specifically highlighted the need for physical therapists to be properly trained and prepared to respond to such pandemic. First, it demonstrated the importance for every professional being updated, trained, and qualified in all technical skills related to infection control strategies such as wearing adequate professional clothing, effective hand washing, and use of personal protective equipment.

Second, the importance of primary care in a good health care system. Patients with at least one comorbidity have higher odds of being hospitalized and being in ICUs. Also, individuals with chronic health conditions are more likely to develop a more severe form of the disease requiring ICU admission, mechanical ventilation, and more resources. Hypertension, cardiovascular diseases, hypercholesterolemia, and diabetes are the most common comorbidities in patients who died from COVID-19.⁵ Consequently, the mortality rate (6.4%) of confirmed cases of COVID-19 in Brazil, within 52 days from the beginning of COVID-19 in every country, is around twice that of other countries, like Germany and Canada (3.0% and 4.1%, respectively), where the primary health care system is more effective in providing primary care.^{6–8} Thus, countries that have greater control over chronic health conditions may

consequently have a lower mortality rate, with a population in better health being an important benefit in pandemic situations like the one we are witnessing with COVID-19.

Third, the importance of the Cardiorespiratory and Intensive Care (CRI) Physical Therapists in the Brazilian health system. CRI Physical Therapy is an established profession worldwide; however, the number of professionals in Brazil is much higher, and dozens of thousands work in hospitals. Interestingly, the ventilatory support and the improvement in musculoskeletal dysfunctions in ICUs are assisted by two professionals (Physical Therapists and Respiratory Therapists) in most countries; however, the CRI Physical Therapists are responsible for both in Brazil. The cumulative function reinforces the importance of the CRI Physical Therapists in the ICUs, as part of the multiprofessional team, minimizing the consequences of hospitalization and facilitating the recovery of patients.

Ventilatory support in patients with COVID-19 includes several approaches: assistance in orotracheal intubation, support and management in mechanical ventilation, removal of airway secretion, changing patient's decubitus to improve gas exchange, and weaning patients from mechanical ventilation.⁹ Additionally, these patients can present with consequences due to intensive care syndrome, including prolonged intubation, continuous sedation, and use of neuromuscular blocking agents that result in muscle weakness.¹⁰ All these consequences from prolonged stay in the ICU are associated with an increase in morbidity and mortality.¹¹ Therefore, it is essential to start early rehabilitation in the ICU to promote rapid functional recovery. Some of these patients will present low exercise capacity, low physical activity level, increased sedentary behavior, dyspnea on exertion, and poorer quality of life after hospital discharge. CRI Physical Therapists will also be requested to help with the post hospital discharge rehabilitation program.

Fourth and finally, this pandemic reinforces the importance for physical therapists to remain scientifically updated. The COVID-19 pandemic resulted in a significant increase in the number of studies reporting the impact of

<https://doi.org/10.1016/j.bjpt.2020.04.004>

1413-3555/© 2020 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

the disease⁹; despite that, the pulmonary repercussion of the disease remains poorly known.¹² The difficulty in understanding how to offer optimal ventilatory support was the primary barrier for the CRI Physical Therapists. The previous professional experience was to administer ventilatory support for pneumonia based on the pathophysiology of Acute Respiratory Distress Syndrome (ARDS), where the severity of hypoxemia is associated with lung compliance. However, in COVID-19, CRI Physical Therapists faced, at least, two distinct phenotypes. Some patients present severe hypoxemia with respiratory system compliance remaining near normal, where hypoxemia seems to occur due to the loss of the hypoxic pulmonary vasoconstriction and impaired regulation of pulmonary blood flow. In another group of patients, severe hypoxemia is associated with lower compliance values, a condition similar to severe ARDS.¹² The difference between COVID-19 and ARDS required CRI Physical Therapists to be updated almost daily to understand how to give the proper ventilatory support.

In conclusion, COVID-19 imposed a huge impact on the health care system in all countries, and everyone had to respond promptly in a very short time. Brazilian hospitals entire buildings and wards have been converted in semi-intensive and ICUs, and multidisciplinary teams, including CRI Physical Therapists, Physicians, and Nurses, have to receive special training for COVID-19. The Brazilian health system presents a special environment because it can count on the presence of CRI Physical Therapists as part of the ICUs multidisciplinary team. However, this is a unique moment in human history, and the countries that have adequately invested in research, the entire health care system, and hospital infrastructure are those that will suffer less in this war.

Funding

Conselho Nacional de Pesquisa (CNPq).

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Dantas LO, Barreto RPG, Ferreira CHJ. Digital physical therapy in the COVID-19 pandemic. *Braz J Phys Ther.* 2020, <http://dx.doi.org/10.1016/j.bjpt.2020.04.006>.
2. Pelicioni PHS, Lord SR. COVID-19 will severely impact older people's lives, and in many more ways than you think! *Braz J Phys Ther.* 2020, <http://dx.doi.org/10.1016/j.bjpt.2020.04.005>.

3. Branson RD, Hess DR, Rubinson L. *SARS CoV-2: Guidance Document.* American Association for Respiratory Care; 2020. Published online March.
4. World Health Organization. *Coronavirus Disease 2019 (COVID-19) Situation Report 46;* 2020.
5. Park M, Cook AR, Lim JT, Sun Y, Dickens BL. A systematic review of COVID-19 epidemiology based on current evidence. *J Clin Med.* 2020;9(4):967.
6. Robert Koch Institut. Coronavirus Disease 2019 (COVID-19) Daily Situation Report of the Robert Koch Institute. Available from: https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Situationsberichte/2020-04-18-en.pdf?__blob=publicationFile. Updated 18.04.20.
7. DATASUS. Ministério da Saúde do Brasil. Disponível em: <https://covid.saude.gov.br/>. Atualizada em: 17/04/2020.
8. Government of Canada. Coronavirus Disease 2019 (COVID-19): Daily Epidemiology Update. Available from: <https://www.canada.ca/content/dam/phac-aspc/documents/services/diseases/2019-novel-coronavirus-infection/surv-covid19-epi-update-eng.pdf>. Updated 17.04.20.
9. Lazzeri M, Lanza A, Bellini R, et al. Respiratory physiotherapy in patients with COVID-19 infection in acute setting: a Position Paper of the Italian Association of Respiratory Physiotherapists. *Monaldi Arch Chest Dis.* 2020;90(1285):163–168.
10. Kress JP, Hall JB. ICU-acquired weakness and recovery from critical illness. *N Engl J Med.* 2014;370(17):1626–1635.
11. Borges RC, Carvalho CR, Colombo AS, da Silva Borges MP, Soriano FG. Physical activity, muscle strength, and exercise capacity 3 months after severe sepsis and septic shock. *Inten Care Med.* 2015;41(8):1433–1444.
12. Gattinoni L, Chiumello D, Caironi P, et al. COVID-19 pneumonia: different respiratory treatment for different phenotypes? *Int Care Med.* 2020, <http://dx.doi.org/10.1007/s00134-020-06033-2>.

Thiago Fernandes Pinto^{a,b,c}, Celso R.F. de Carvalho^{a,d,*}

^a Department of Physical Therapy, School of Medicine, Universidade de São Paulo (USP), São Paulo, SP, Brazil

^b Department of Physical Therapy, Universidade de Mogi das Cruzes, Mogi das Cruzes, SP, Brazil

^c Department of Physical Therapy, Centro Universitário das Américas, São Paulo, SP, Brazil

^d São Paulo Coordinator from the Associação Brasileira de Fisioterapia Cardiorrespiratória e Terapia Intensiva (Assobrafir), São Paulo, SP, Brazil

* Corresponding author at: Department of Physical Therapy, School of Medicine, Universidade de São Paulo, Rua Cipotânea, 51, Cidade Universitária, CEP: 05360-000, São Paulo, SP, Brazil.

E-mail address: cscarval@usp.br (C.R. Carvalho).