


# *Drosophila* balancer reengineering using polycistronic gRNA for CRISPR/Cas9 gene editing

Leif Benner<sup>1,2</sup>, Brian Oliver<sup>1</sup>

1. Section of Developmental Genomics, Laboratory of Cellular and Developmental 10 Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD, 20892

2. Department of Biology, Johns Hopkins University, Baltimore, MD, 21218



**Figure 1** A) U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,v,f,car</sup> cassette design. Red arrow represents transcription start site. B) Number of phenotypic mutants transmitted from *Act5c-Cas9/FM7c; U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,v,f,car</sup>/+* females. Genotypes of the progeny are *FM7c/y<sup>1</sup> cv<sup>1</sup> v<sup>1</sup> f<sup>1</sup>*. C) Mutations induced at each locus of the respective phenotypes from B. Red letters represent insertions and dashes represent deletions. Black arrows show the Cas9 cleavage site.

## Description

A number of reagents have been generated for CRISPR/Cas9 genome editing in *Drosophila* (Gratz et al. 2014; Ren et al. 2013; Kondo and Ueda 2013; Port et al. 2014; Port and Bullock 2016). The ability to efficiently express multiple gRNAs from a single vector provides the advantage of creating multiple CRISPR/Cas9 alleles at a time. An elegant design has previously shown that gRNAs separated by tRNAs allow for genomic editing at multiple loci in *Drosophila* (Port and Bullock 2016). We generated a similar polycistronic gRNA cassette under the control of the *Drosophila snRNA:U6:96Ac* (U6:3) regulatory sequences. Individual gRNAs were separated by the *Drosophila tRNA:His-GTG-1-4* tRNA (tRNA<sup>His</sup>) and targeted the genes *yellow* (*y*), *crossveinless* (*cv*), *vermillion* (*v*), *forked* (*f*), and *carnation* (*car*) along the X chromosome (A). In theory, after expression of the gRNA cassette, the tRNAs should lead to the formation of secondary RNA structures and be recognized for cleavage by RNase enzymes P and Z (Frendewey et al. 1985; Dubrovsky et al. 2004), releasing the gRNAs for incorporation into Cas9. The gRNA cassette (U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,v,f,car</sup>) was stably integrated into the *Drosophila* genome and crossed to a Cas9 line under the control of the Act5C promoter (Zhang, Koolhaas, and Schnorrer 2014). Females with the genotype *Act5c-Cas9/FM7c; U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,v,f,car</sup>/+* showed *y*<sup>-</sup>, *cv*<sup>-</sup>, and *f*<sup>-</sup> phenotypes while females with the genotype *y<sup>1</sup> w<sup>1</sup>/FM7c; U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,v,f,car</sup>/+* were wild type. We also conducted experiments with a wild type X chromosome in the place of the *FM7c* balancer chromosome. Results from these experiments were the same, with only evidence for *y*<sup>-</sup>, *cv*<sup>-</sup>, and *f*<sup>-</sup> phenotypes and all females were *v*<sup>+</sup> and *car*<sup>+</sup>. We then decided to outcross *Act5c-Cas9/FM7c; U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,v,f,car</sup>/+* females to a *y<sup>1</sup> cv<sup>1</sup> v<sup>1</sup> f<sup>1</sup>* stock and score germline transmission of editing at *y*, *cv*, and *f*, on the *FM7c* balancer chromosome (*FM7c = In(1)FM7, y<sup>31d</sup> sc<sup>8</sup> w<sup>a</sup> sn<sup>X2</sup> v<sup>Of</sup> g<sup>d</sup> B<sup>1</sup>*). We could not score *v* since *FM7c* is *v<sup>Of</sup>* and also did not score *car* since we lacked evidence for *car* phenotypes based on the wild type X experiments. Germline transmission of mutants resulted in 14% of progeny not mutant at any locus, 40% were mutant at one locus, 40% were mutant at two loci, and 6% were mutant at three loci (B). We sequenced the gRNA targeted genomic sites of *y*, *cv*, *v*, *f*, and *car* of two phenotypically *y<sup>+</sup> cv<sup>-</sup> f<sup>-</sup>* and two *y<sup>-</sup> cv<sup>-</sup> f<sup>-</sup>* balancers to compare with the original *FM7c* chromosome. All flies contained out-of-frame indels at the loci consistent with their

11/13/2018 – Open Access

phenotypes. None of the flies had mutations at *v*, and one fly had an in-frame indel at *car* (C). We are unsure why these flies did not show a *car*<sup>-</sup> phenotype, however, our sequencing results indicate that germline transmission of editing events did occur at the *car* locus. Our data suggests that gRNAs targeting *y*, *cv*, *f*, and *car* were cleaved from the cassette and loaded into Cas9 allowing for subsequent genomic editing. The ability to express multiple gRNAs from a single vector offers advantages over traditional gRNA vectors for *Drosophila* and will allow researchers to expand their experimental repertoire for CRISPR/Cas9 genome editing. Practically, polycistronic gRNA vectors can be used to quickly generate and expand new balancer genotypes, that would otherwise be difficult and time-consuming through traditional methods.

## Methods

U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,vf,car</sup> was synthesized by GenScript (Piscataway, NJ) and subcloned into pUAS-attB (Bischof et al. 2007) generating pU6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,vf,car</sup>. pU6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,vf,car</sup> was integrated into the *P{CaryP}attP40* landing site (Markstein et al. 2008) through phi-C31 mediated integration (Bischof et al. 2007). Embryo injections for transgenesis and transformant recovery was completed by BestGene (Chino Hills, CA). Sanger sequencing was completed by Genewiz (Plainfield, NJ). Flies were raised on ‘Fly Food B’ (LabExpress, Ann Arbor, MI) at 25°C. A *y*<sup>-</sup> phenotype was scored by a reduction in pigmentation of the bristles for *y*<sup>3<sup>ld</sup></sup> on FM7c. tRNA<sup>His</sup> corresponds to *tRNA:His-GTG-1-4* (FBgn0011878) and contains an additional six upstream nucleotides (GATGTA) from the annotated transcript. All sequences are available upon request.

## Reagents

FM7c = FBba0000009

FM7c, *cv*<sup>WJB</sup> *f*<sup>DPB</sup> = this study

*y*<sup>1</sup> *v*<sup>1</sup>; *P{CaryP}attP40* = BDSC 36304 (or FBst0036304)

*y*<sup>1</sup> *M{vas-int.Dm}ZH-2A w*<sup>\*</sup> = BDSC 40161 (or FBst0040161)

*y*<sup>1</sup> *M{Act5C-Cas9.P.RFP-}ZH-2A w*<sup>1118</sup> *DNAIig4*<sup>169</sup> = BDSC 58492 (or FBst0058492)

*y*<sup>1</sup> *w*<sup>1</sup>; *P{y<sup>+7.7</sup> w<sup>+mC</sup>=U6:3-tRNA<sup>His</sup>-gRNA<sup>y,cv,vf,car</sup>}attP40* = this study

## References

- Bischof, Johannes, Robert K. Maeda, Monika Hediger, François Karch, and Konrad Basler. 2007. “An Optimized Transgenesis System for *Drosophila* Using Germ-Line-Specific phiC31 Integrases.” *Proceedings of the National Academy of Sciences of the United States of America* 104 (9): 3312–17.
- Dubrovsky, Edward B., Veronica A. Dubrovskaya, Louis Levinger, Steffen Schiffer, and Anita Marchfelder. 2004. “*Drosophila* RNase Z Processes Mitochondrial and Nuclear Pre-tRNA 3’ Ends in Vivo.” *Nucleic Acids Research* 32 (1): 255–62.
- Friendewey, D., T. Dingermann, L. Cooley, and D. Soll. 1985. “Processing of Precursor tRNAs in *Drosophila*: Processing of the 3’ end Involves an Endonucleolytic Cleavage and Occurs after 5’end Maturation.” *The Journal of Biological Chemistry* 260 (1): 449–54.
- Gratz, Scott J., Fiona P. Ukken, C. Dustin Rubinstein, Gene Thiede, Laura K. Donohue, Alexander M. Cummings, and Kate M. O’Connor-Giles. 2014. “Highly Specific and Efficient CRISPR/Cas9-Catalyzed Homology-Directed Repair in *Drosophila*.” *Genetics* 196 (4): 961–71.
- Kondo, Shu, and Ryu Ueda. 2013. “Highly Improved Gene Targeting by Germline-Specific Cas9 Expression in *Drosophila*.” *Genetics* 195 (3): 715–21.
- Markstein, Michele, Chrysoula Pitsouli, Christians Villalta, Susan E. Celniker, and Norbert Perrimon. 2008. “Exploiting Position Effects and the Gypsy Retrovirus Insulator to Engineer Precisely Expressed Transgenes.” *Nature Genetics* 40 (4): 476–83.
- Port, Phillip, and Simon L. Bullock. 2016. “Augmenting CRISPR Applications in *Drosophila* with tRNA-Flanked sgRNAs.” *Nature Methods* 13 (September): 852.

11/13/2018 – Open Access

Port, Phillip, Hui-Min Chen, Tzumin Lee, and Simon L. Bullock. 2014. “Optimized CRISPR/Cas Tools for Efficient Germline and Somatic Genome Engineering in *Drosophila*.” *Proceedings of the National Academy of Sciences of the United States of America* 111 (29): E2967–76.

Ren, Xingjie, Jin Sun, Benjamin E. Housden, Yanhui Hu, Charles Roesel, Shuailiang Lin, Lu-Ping Liu, et al. 2013. “Optimized Gene Editing Technology for *Drosophila Melanogaster* Using Germ Line-Specific Cas9.” *Proceedings of the National Academy of Sciences of the United States of America* 110 (47): 19012–17.

Zhang, Xu, Wouter H. Koolhaas, and Frank Schnorrer. 2014. “A Versatile Two-Step CRISPR- and RMCE-Based Strategy for Efficient Genome Engineering in *Drosophila*.” *G3* 4 (12): 2409–18.

### **Funding**

This research was supported in part by the Intramural Research Program of the NIH, National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) to BO.

**Reviewed by** Danny Miller

**Received** 08/28/2018, **Accepted** 11/12/2018. **Published Online** 11/13/2018.

**Copyright** © 2018 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Benner, L., & Oliver, B. (2018). *Drosophila* balancer reengineering using polycistronic gRNA for CRISPR/Cas9 gene editing. <https://doi.org/10.17912/SG7D-SD61>