




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to Editors

Fighting against frailty and sarcopenia – As well as COVID-19?

Geriatric syndromes, including sarcopenia and frailty, are necessarily on the agenda - as the mortality rate of COVID-19 is higher in the elderly [1]. Herein, sarcopenia is defined as age-related loss of muscle mass and function, and frailty as multi-system impairment related with increased vulnerability to stressors [2]. Of note, sarcopenia is an essential component of physical frailty; therefore they are like the two sides of the same coin [2]. Both are associated with hospitalization, increased disability, and even mortality. Yet, frail patients are at increased risk of infectious diseases due to the increased inflammatory condition and significant immune system changes e.g. impaired proliferation of peripheral mononuclear cells and upregulated monocyte expression [3]. Likewise, the cytokine storm in COVID-19 patients is known to have poor prognosis due to the increased inflammatory response [4].

In short, we simply underscore that preventive measures for frailty/sarcopenia i.e. exercise, nutrition and vitamin (D) supplementation can fortify the fight against COVID-19 as well [5].

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to

influence the work reported in this paper.

References

- [1] Koff WC, Williams MA. Covid-19 and immunity in aging populations - a new research agenda. *N Engl J Med* 2020. <https://doi.org/10.1056/NEJMp2006761>. in press.
- [2] Cesari M, Landi F, Vellas B, Bernabei R, Marzetti E. Sarcopenia and physical frailty: two sides of the same coin. *Front Aging Neurosci* 2014;28(6):192. <https://doi.org/10.3389/fnagi.2014.00192>.
- [3] Yao X, Li H, Leng SX. Inflammation and immune system alterations in frailty. *Clin Geriatr Med* 2011;27(1):79–87. <https://doi.org/10.1016/j.cger.2010.08.002>.
- [4] Mehta P, McAuley DF, Brown M, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;395(10229):1033–4. [https://doi.org/10.1016/S0140-6736\(20\)30628-0](https://doi.org/10.1016/S0140-6736(20)30628-0).
- [5] Kara M, Ata AM, Kaymak B, Özçakar L. Ultrasound imaging and rehabilitation of muscle disorders part 2. Non-traumatic conditions. *Am J Phys Med Rehabil* 2020. <https://doi.org/10.1097/PHM.0000000000001352>. In Press.

Timur Ekiz^{a,*}, Murat Kara^b, Levent Özçakar^b

^aDepartment of Physical and Rehabilitation Medicine, Türkmenbaşı Medical Center, Adana, Turkey

^bDepartment of Physical and Rehabilitation Medicine, Hacettepe University Medical School, Ankara, Turkey

E-mail address: timurekiz@gmail.com (T. Ekiz).

* Corresponding author at: Türkmenbaşı Bulvarı Botanik Evleri Osmanbey Apt. C Blok No:3/B, 01130, Seyhan, Adana, Turkey.