


COVID-19: what if the brain had a role in causing the deaths?

C. Tassorelli^{a,b} , F. Mojoli^{c,d},
F. Baldanti^{c,e}, R. Bruno^{c,f} and
M. Benazzo^{c,g}

^aDepartment of Brain and Behavioral Sciences, University of Pavia, Pavia,
^bIRCCS Mondino Foundation, Pavia,
^cDepartment of Clinical, Surgical, Diagnostic and Pediatric Sciences, University of Pavia, Pavia, ^dDepartment of Anaesthesia and Intensive Care, Fondazione IRCCS Policlinico San Matteo, Pavia,
^eMolecular Virology Unit, Fondazione IRCCS Policlinico San Matteo, Pavia,
^fInfectious Diseases Unit, Fondazione IRCCS Policlinico San Matteo, Pavia, and ^gDepartment of Otorhinolaryngology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

Correspondence: C. Tassorelli, IRCCS Mondino Foundation, Via Mondino, 2, 27100 Pavia, Italy (tel.: 0039 (0) 382 380419; fax: 0039 (0)382 380448; e-mail: cristina.tassorelli@unipv.it).

Keywords: coronavirus, neurotropism, respiratory centres, vagus nerve

doi:10.1111/ene.14275

Received: 8 April 2020

Accepted: 18 April 2020

In the last few weeks, Italy first and then several other countries across the world have been swept up by a deadly wave of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which causes the illness named COVID-19, from the acronym CO (corona) VI (virus) D (disease) and 19 (year of the virus identification). The medical community is working day and night to assist affected people and experts in communicable diseases are striving in multiple ways to understand the progression of events leading to the lethal respiratory syndrome.

Lombardy is the most heavily affected area in the Italian coronavirus epidemic and the one with the highest number of

reported fatalities. Doctors assisting patients in the front line of the Policlinico San Matteo of Pavia, one of the Italian hospitals with the highest number of SARS-CoV-2 cases, report that several patients in the latest stages of disease, when their lung parenchyma is devastated by the infection, may not manifest dyspnoea (personal observations). Data from the Wuhan area show that more than 90% of patients in need of intensive care cannot breathe spontaneously [1]. On the other hand, patients with severely affected lungs may be paucisymptomatic and recover fully from the infection.

Thus, at least in some cases, there seems to be a discrepancy between the severity of the lung involvement and the respiratory function. Severe COVID-19 leads to death via multiple mechanisms, including myocardial injury, kidney failure, shock and disseminated intravascular coagulopathy [1,2]. However, an additional explanation worth exploring in this phase of uncertainties, also in the light of the initial evidence of neuronal localization of COVID-19, is that the respiratory failure may be driven by a dysfunction of the cardiorespiratory centres in the brainstem.

Coronaviruses (CoVs) are neurotropic [3] and SARS-CoV particles have been detected in neurons of the human brain [4,5]. Given the high level of genetic homology of CoVs, which also share many other characteristics, it is also likely that SARS-CoV-2 may gain access to the central nervous system, where it can induce neuronal injury. The peripheral invasion of nerves by SARS-CoV-2 is further suggested by the occurrence of anosmia and ageusia in more than 5% of a population of 214 COVID-19 people [6]. From the peripheral nerves the virus may have access to the central nervous system via trans-synaptic transfer, a possibility documented for other coronaviruses *in vitro* and *in vivo* [7]. Recent experimental evidence demonstrates that a human CoV strain, HCoV OC43, can travel from the nasal cavity to the olfactory bulb, then spreading to the piriform cortex and ultimately to the brainstem, via both passive diffusion and axonal transport [8,9]. Alternatively or in addition – which would increase

central nervous system vulnerability to SARS-CoV-2 diffusion – the virus may travel retrogradely along the vagus nerve, which innervates many of the visceral organs that can be invaded by the virus beyond the lungs, such as the heart and the gastrointestinal tract. Once in the vagal nerve endings, retrograde axonal transport may grant access to the brainstem.

Whatever the entry point, the final destination of SARS-CoV-2 could be the neurons of the nucleus of the solitary tract in the medulla oblongata, which form the dorsal respiratory group that generates the basic rhythm of respiration and emits repetitive bursts of inspiratory neuronal action potentials. The nucleus of the solitary tract is connected with the nucleus ambiguus, the main components of the ventral respiratory group that controls forceful inspiration and expiration. Interestingly a similar efferent pathway can also be traced for the cardiac control centre.

A recent retrospective analysis of COVID-19 manifestations in 214 subjects in the Wuhan area reported the presence of neurological symptoms in 36% of them, a percentage that rises to 46% when the most severely affected subjects are considered [10]. Thus, it is possible that the invasion of the nervous system by SARS-CoV-2 is one of the pathogenetic mechanisms leading to death. This is a call to action for using the available sophisticated neuroimaging, neurophysiological and biochemical techniques to detect and document SARS-CoV-2 invasion of the peripheral and central nervous system. These investigations will of course greatly benefit also from systematic histopathological analyses of autaptic tissues to enhance our understanding of the preferential routes of invasion and sites of possible neuronal damage. This may clarify additional modalities through which COVID-19 is killing people all over the world, possibly provide markers for the identification of the subjects who are at higher risk to evolve into the more serious manifestations of COVID-19 and, even more importantly, foster the identification of additional therapeutic strategies.

Disclosure of conflicts of interest

The authors declare no financial or other conflicts of interest.

References

1. Wang D, Hu B, Hu C, *et al.* Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020; **323**: 1061.
2. Guo T, Fan Y, Chen M, *et al.* Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020; **5**: 811–818.
3. Li YC, Bai WZ, Hirano N, *et al.* Coronavirus infection of rat dorsal root ganglia: ultrastructural characterization of viral replication, transfer, and the early response of satellite cells. *Virus Res* 2012; **163**: 628–635.
4. Ding Y, He L, Zhang Q, *et al.* Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus transmission pathways. *J Pathol* 2004; **203**: 622–630.
5. Gu J, Gong E, Zhang B, *et al.* Multiple organ infection and the pathogenesis of SARS. *J Exp Med* 2005; **202**: 415–424.
6. Vaira LA, Salzano G, Deiana G, De Riu G. Anosmia and ageusia: common findings in COVID-19 patients. *Laryngoscope* 2020; **130**: 1787.
7. Mengeling WL, Boothe AD, Ritchie AE. Characteristics of a coronavirus (strain 67N) of pigs. *Am J Vet Res* 1972; **33**: 297–308.
8. Desforges M, Le Coupanec A, Dubeau P, *et al.* Human coronaviruses and other respiratory viruses: underestimated opportunistic pathogens of the central nervous system? *Viruses* 2019; **12**: E14.
9. Dube M, Le Coupanec A, Wong AHM, *et al.* Axonal transport enables neuron-to-neuron propagation of human coronavirus OC43. *J Virol* 2018; **92**: e00404-18.
10. Mao L, Wang M, Chen S, *et al.* Neurological manifestations of hospitalized patients with COVID-19 in Wuhan, China: a retrospective case series study. *MedRxiv* 2020. <https://doi.org/10.1101/2020.02.22.20026500>