

[ ORIGINAL ARTICLE ]

## Epidemiological Features and Clinical Presentations of Acute Coronary Syndrome in Young Patients

Hiroaki Sawada, Hirohiko Ando, Hiroaki Takashima, Katsuhisa Waseda, Masahiro Shimoda, Hirofumi Ohashi, Akihiro Suzuki, Shinichiro Sakurai, Yusuke Nakano and Tetsuya Amano

### Abstract:

**Objective** Although acute coronary syndrome (ACS) is an uncommon entity in young patients, it constitutes an important problem due to the devastating effects of the disease on the more active lifestyle of young patients. At present, there are no guidelines regarding the prevention of ACS in young patients.

**Methods** We performed a retrospective study of ACS patients between 2014 and 2017. Epidemiological data, clinical findings, and short-term outcomes were evaluated between young ACS patients ( $\leq 50$  years old) and elderly ACS patients ( $> 50$  years old).

**Results** Of a total of 361 consecutive ACS patients, 37 were young ACS patients (10.2%). Compared with elderly ACS patients, young ACS patients showed a higher prevalence of males (94.6% vs. 73.8%,  $p < 0.001$ ), current smoking (70.3% vs. 29.9%;  $p < 0.001$ ), and overweight persons (67.6% vs. 27.8%,  $p < 0.001$ ). The eicosapentaenoic acid (EPA)/arachidonic acid (AA) ratio was significantly lower in young ACS patients than in elderly ACS patients [0.17 (0.12-0.25) vs. 0.25 (0.18-0.37),  $p = 0.002$ ]. The prevalence of cardio-pulmonary arrest and percutaneous cardiopulmonary support use was higher in young ACS patients than in elderly ACS patients (24.3% vs. 8.6%,  $p = 0.003$ , 16.2% vs. 3.1%,  $p < 0.001$ ).

**Conclusion** The features were markedly different between young ACS patients and elderly ACS patients. In young ACS patients, smoking, being overweight, and a low EPA/AA ratio were distinctive risk factors, and more serious clinical presentations were observed at the onset of ACS than in older patients.

**Key words:** acute coronary syndrome, young patients, smoking, EPA/AA, cardiac arrest

(Intern Med 59: 1125-1131, 2020)

(DOI: 10.2169/internalmedicine.4138-19)

### Introduction

Coronary heart disease remains a major cause of death and disability in all regions of the world (1). Although acute coronary syndrome (ACS) primarily occurs in the older age group, younger people can also be affected, and the number of ACS in younger people has been gradually increasing (2). Furthermore, it constitutes an important problem for patients and treating physicians due to the devastating effects of the disease on the more active lifestyle of younger patients. However, there are currently no guidelines or consensus regarding the prevention and treatment of ACS in young patients.


In the present study, we compared the clinical pictures of

ACS in young patients with those of elderly patients to help determine the optimal prevention and treatment strategies.

### Materials and Methods

#### Study population

This study was designed as a single-center retrospective study to evaluate the predictive factors and clinical characteristics of ACS in young patients. We analyzed the data of consecutive patients with ACS who underwent percutaneous coronary intervention (PCI) for *de novo* native coronary artery lesions between January 2014 and December 2017 at Aichi Medical University. ACS included ST segment elevation myocardial infarction, non-ST segment elevation myo-



**Figure 1.** Distinctive risk factors associated with ACS in young patients. (A) Prevalence of current smoking between young ACS patients and elderly ACS patients. (B) Prevalence of overweight between young ACS patients and elderly ACS patients. (C) Prevalence of EPA/AA  $\leq 0.3$  between young ACS patients and elderly ACS patients. ACS: acute coronary syndrome, EPA: eicosapentaenoic acid, AA: arachidonic acid

cardial infarction, or unstable angina. Patients  $< 50$  years old were classified as young ACS patients, while those  $\geq 50$  years old were classified as elderly ACS patients.

### Procedures and coronary risk factors

PCI was performed according to the standard techniques. Stenting strategy (stent type, stent size, direct stenting, or non-stent strategy), thrombus aspiration or distal protection, and post-dilatation were left to the operator's discretion. Before or during PCI, all patients were administered 200 mg of aspirin and a loading dose of a P2Y<sub>12</sub> inhibitor (300 mg of clopidogrel or 20 mg of prasugrel). On the day after the loading dose, a maintenance dose of dual antiplatelet therapy [aspirin 100 mg and a P2Y<sub>12</sub> inhibitor (75 mg of clopidogrel or 3.75 mg of prasugrel)] was administered. Hypertension was defined as a medical history of systemic hypertension or the use of antihypertensive treatment. Dyslipidemia was defined as a previous diagnosis of dyslipidemia or the use of a lipid-lowering treatment. Overweight was defined as having a body mass index (BMI)  $\geq 25$  kg/m<sup>2</sup> based on the proposed classification by the WHO (3). For the short-term outcome, the all-cause mortality within 30 days after the index PCI was evaluated.

### Statistical analyses

All statistical analyses were performed with the SPSS software program (version 22.0; IBM, Armonk, USA). Data are expressed as the mean  $\pm$  standard deviations or as medians and interquartile ranges with differences (95% confidence intervals). Categorical variables are expressed as frequencies (%). The normality of distribution was tested by the Kolmogorov-Smirnov test. Continuous variables were

compared using the unpaired Student's *t*-test, and categorical variables were compared using the chi-squared or Fisher's exact tests where appropriate. Mann-Whitney U tests were performed for non-parametric data. Statistical significance was assumed at a probability (*p*) value of  $< 0.05$ .

## Results

### Patient characteristics

A total of 361 consecutive ACS patients were evaluated in the present study. Of these, 37 were young ACS patients, accounting for 10.2% of total ACS patients, and the remaining 324 (89.8%) were elderly ACS patients (Supplementary material 1). The proportion of men was predominantly higher among the young ACS patients than among the elderly ACS patients (94.6% vs. 73.8%,  $p < 0.001$ ). In terms of conventional risk factors, there were no significant differences in the prevalence of hypertension, diabetes, and dyslipidemia between the two groups (Table 1). The prevalence of current smoking was significantly higher among young ACS patients than among elderly ACS patients (Fig. 1). Regarding anthropometric measurements, in young ACS patients, the BMI was significantly higher ( $27.8 \pm 4.7$  kg/m<sup>2</sup> vs.  $23.0 \pm 3.8$  kg/m<sup>2</sup>,  $p < 0.001$ ) as was the prevalence of overweight than in elderly ACS patients (Fig. 1). Regarding the medications administered before admission, angiotensin receptor blockers, Ca-channel antagonists, and antiplatelet agents were more commonly used in elderly ACS patients than in young ACS patients (Supplementary material 2).

**Table 1. Baseline Characteristics.**

	Young ACS (n=37)	Elderly ACS (n=324)	P
Age, years	44.9±4.1	71.1±9.8	<0.001
Male gender	35 (94.6)	239 (73.8)	0.005
Hypertension	17 (45.9)	194 (59.9)	0.10
Diabetes	11 (29.7)	96 (29.6)	0.99
Dyslipidemia	16 (43.2)	173 (53.4)	0.24
Current smoker	26 (70.3)	97 (29.9)	<0.001
Hemodialysis	1 (2.7)	10 (3.1)	0.90
Previous myocardial infarction	3 (8.1)	22 (6.8)	0.76
Height, cm	170.2±5.5	161.5±9.4	<0.001
Body weight, kg	80.2±13.9	60.5±13.3	<0.001
Body mass index, kg/m <sup>2</sup>	27.8±4.7	23.0±3.8	<0.001
HDL-cholesterol, mg/dL	38.4±7.1	42.7±12.4	0.02
LDL-cholesterol, mg/dL	104.7±30.9	101.4±31.6	0.82
Triglyceride, mg/dL	156 (114-198)	102 (73-138)	<0.001
HbA1c, %	6.5±1.8	6.3±0.9	0.17
CRP, mg/dL	0.17 (0.08-0.35)	0.18 (0.08-0.59)	0.64
EPA, µg/mL	33.8 (27.0-46.6)	47.2 (30.4-66.0)	0.02
DHA, µg/mL	112.7 (84.9-126.5)	123.8 (101.4-156.0)	0.02
AA, µg/mL	211.9 (168.4-250.3)	182.2 (149.5-217.7)	0.01
EPA/AA ratio	0.17 (0.12-0.25)	0.25 (0.18-0.37)	0.002

Values are mean±SD, n (%), or median (interquartile range).

ACS: acute coronary syndrome, EPA: eicosapentaenoic acid, DHA: docosahexaenoic acid, AA: arachidonic acid

### Laboratory findings

Low-density lipoprotein cholesterol levels were similar between the two groups. High-density lipoprotein cholesterol levels were significantly lower and triglycerides significantly higher in young ACS patients than in elderly ACS patients [Table 1; 38.4±7.1 mg/dL vs. 42.7±12.4 mg/dL,  $p=0.02$ ; 156 (114-198) mg/dL vs. 102 (73-138) mg/dL,  $p<0.001$ , respectively]. The HbA1c values were similar between the two groups. Serum eicosapentaenoic acid (EPA) and serum docosahexaenoic acid were significantly lower and serum arachidonic acid (AA) significantly higher in young ACS patients than in elderly ACS patients. Consequently, the EPA/AA ratio was significantly lower in young ACS patients than in elderly ACS patients [0.17 (0.12-0.25) vs. 0.25 (0.18-0.37),  $p=0.002$ ]. The prevalence of patients with an EPA/AA ratio  $\leq 0.3$  was significantly higher in young ACS patients than in elderly ACS patients (Fig. 1).

### Clinical presentations

Angiographic findings and clinical presentations are shown in Table 2. There was no significant difference in the classification of ACS (ST segment elevation myocardial infarction, non-ST segment elevation myocardial infarction, or unstable angina) between the two groups. Single-vessel disease predominated in young ACS patients (83.8%), whereas it was found in approximately half of the elderly ACS patients (51.5%). Regarding the clinical presentations, more serious conditions were observed in young ACS patients

than in elderly ACS patients. Patients with a higher Killip class were more frequently observed among young ACS patients than among elderly ACS patients. The prevalence of patients with cardio-pulmonary arrest (CPA) on arrival as well as those requiring percutaneous cardiopulmonary support (PCPS) was significantly higher among young ACS patients than among elderly ACS patients (Fig. 2). Even after PCI, patients with a lower thrombolysis in myocardial infarction (TIMI) flow grade were more frequently observed among young ACS patients than among elderly ACS patients. While the difference was not statistically significant, the 30-day mortality was numerically higher in young ACS patients than in elderly ACS patients.

### Discussion

The present study highlighted the differing features between young and elderly ACS patients. The main findings were as follows: 1) a higher prevalence of men, current smokers, and overweight individuals was observed among young ACS patients; 2) the EPA/AA ratio was lower in young ACS patients than in older ones; and 3) despite multivessel diseases being uncommon, serious conditions were more frequently observed in young ACS patients than in older ones.

Although ACS has a tragic impact on the daily life of younger patients, there are currently no guidelines or consensus regarding the prevention and treatment of ACS in this population. Furthermore, there have been almost no large-

**Table 2. Clinical Presentations.**

	Young ACS (n=37)	Elderly ACS (n=324)	P
Clinical presentation			0.47
STEMI	30 (81.1)	211 (65.1)	
NSTEMI	2 (5.4)	25 (7.7)	
UAP	5 (13.5)	88 (27.2)	
Systolic blood pressure, mmHg	139.6±27.8	147.0±29.6	0.24
Killip classification			<0.001
Class I	23 (62.2)	269 (83.0)	
Class II	1 (2.7)	10 (3.1)	
Class III	1 (2.7)	16 (4.9)	
Class IV	12 (32.4)	29 (9.0)	
Vessel related to the ACS			0.22
Left anterior descending artery	22 (59.5)	137 (42.3)	
Left circumflex artery	2 (5.4)	45 (13.9)	
Right coronary artery	13 (35.1)	129 (39.8)	
Left main artery	0 (0)	12 (3.7)	
Bypass graft	0 (0)	1 (0.3)	
No. of affected vessels			0.01
1	31 (83.8)	167 (51.5)	
2	6 (16.2)	103 (31.8)	
3	0 (0)	54 (16.7)	
TIMI flow grade before PCI			0.07
0	23 (62.2)	146 (45.1)	
I	0 (0)	39 (12.0)	
II	6 (16.2)	48 (14.8)	
III	8 (21.6)	91 (28.1)	
TIMI flow grade after PCI			<0.001
0	1 (2.7)	0 (0)	
I	3 (8.1)	0 (0)	
II	3 (8.1)	16 (4.9)	
III	30 (81.1)	308 (95.1)	
Maximum CK, IU/L	2,135 (792-5,383)	1,154 (307-2,762)	<0.001
30-day mortality	5 (13.5)	22 (6.8)	0.14

Values are mean±SD or n (%).

STEMI: ST segment elevation myocardial infarction, NSTEMI: non-ST segment elevation myocardial infarction, UAP: unstable angina, ACS: acute coronary syndrome, TIMI: thrombolysis in myocardial infarction, PCI: percutaneous coronary intervention, IABP: intra-aortic balloon pump counterpulsation, PCPS: percutaneous cardiopulmonary support

scale trials recently focusing on young ACS patients. Under these circumstances, the present study might therefore provide a roadmap for the more effective prevention and management of young ACS patients.

ACS is generally known to occur more often in men than in women (4). In the present study as well, an overwhelming majority of young ACS patients were men (94.6%), which is consistent with the high prevalence of men reported in the Framingham Heart Study (5). It has also been well established that cigarette smoking is a powerful risk factor for coronary artery disease (6, 7). Cigarette smoking promotes vascular endothelial dysfunction and inflammation, resulting in atherosclerosis progression and changes in plaque composition (8, 9). In public settings, the smoking rate is higher among younger people than older people, and Japan Tobacco reported that the smoking rate of Japanese men in their 30s and 40s was around 35% (10). The smok-

ing rate in young ACS patients (70.3%) in the present study is still higher than in the general population. Being overweight was also found to be characteristic of young ACS patients (67.6%) in this study and is widely recognized as a risk factor for coronary heart disease in the elderly. An autopsy study demonstrated that a high BMI was associated with both fatty streaks and raised coronary atherosclerotic lesions in young men (11).


Of note, however: hypertension, diabetes, and dyslipidemia were not distinctive features in this population when compared with elderly ACS patients. Laboratory findings showed that young ACS patients had lower high-density lipoprotein (HDL) cholesterol levels and higher triglyceride levels than elderly ACS patients, which was consistent with the results of a previous study (12). The EPA/AA ratio was also a contributor to ACS in young patients, and almost all young ACS patients (96.8%) showed an EPA/AA ratio ≤0.3.

Several epidemiologic studies have shown that a lower EPA/AA ratio was associated with unstable plaque composition (13-15), cardiovascular events (including ACS) (16-18) and fatal arrhythmic events in the early phase of AMI (19). EPA has beneficial effects on multiple atherosclerotic processes, from protecting against endothelial dysfunction to re-


ducing plaque formation, inflammation, and vulnerability (20-23). In the present study, elderly ACS patients also showed low EPA/AA ratios [0.25 (0.18-0.37)], while young ACS patients showed still lower EPA/AA ratios [0.17 (0.12-0.25)], representing an even higher risk of ACS in young patients. The EPA/AA ratio generally increases with age (24) due to differences in the consumption of dietary fish and age-related changes in serum fatty acid metabolism (25). Therefore, we cannot rule out the possibility that a lower EPA/AA ratio in young ACS patients represents the general findings in young people. This result is in line with the findings of a previous study that showed a reduced EPA/AA ratio as a contributor to the early onset of ACS (26).

Ultimately, the present study extracted three modifiable risk factors (current smoking, being overweight and a low EPA/AA ratio) as distinctive features of young ACS patients. The prevalence of these three cumulative risk factors is indicated in Fig. 3. It is noteworthy that about 90% of young ACS patients had multiple risk factors, while only 40% of elderly ACS patients had multiple risk factors. From a practical point of view, our results suggest that the modification of these risk factors can be expected to prevent the onset of ACS in younger populations. This hypothesis needs to be investigated through prospective interventional trials in the future.

Despite the number of coronary lesions being fewer, a higher Killip class and a higher prevalence of CPA and PCPS use were observed in young ACS patients than in older patients. Previous studies have shown a younger age to



**Figure 2.** Prevalence of CPA on arrival and PCPS use. (A) Prevalence of CPA on arrival between young ACS patients and elderly ACS patients. (B) Prevalence of PCPS use between young ACS patients and elderly ACS patients. CPA: cardiopulmonary arrest, PCPS: percutaneous cardiopulmonary support, ACS: acute coronary syndrome



**Figure 3.** Prevalence of cumulative three risk factors (current smoking, overweight and EPA/AA ≤ 0.3). The prevalence of three cumulative risk factors (current smoking, overweight and EPA/AA ≤ 0.3) in both young ACS patients and elderly ACS patients is shown. Patients with two or more risk factors accounted for 90.3% of young ACS patients and 36.4% of elderly ACS patients. Conversely, patients with no risk factors accounted for only 3.2% of young ACS patients. ACS: acute coronary syndrome, EPA: eicosapentaenoic acid, AA: arachidonic acid



be significantly associated with the occurrence of life-threatening arrhythmia in the setting of acute myocardial infarction (27, 28). The absence of angina before the onset of ACS, which is a characteristic finding in young ACS patients (29), is also associated with life-threatening arrhythmia (27).

### Study limitations

Several limitations associated with the present study warrant mention. This was a single-center study with a relatively small population, which might have resulted in the lack of statistical power. Consequently, the present study might have missed important features of young ACS patients. In addition, this was a post-hoc analysis; the results are therefore hypothesis-generating. Larger prospective studies are warranted to validate our results.

### Conclusion

Young ACS patients present with markedly different features from those seen in elderly ACS patients. In young ACS patients, smoking, being overweight, and a low EPA/AA ratio were distinctive risk factors, and more serious clinical presentations were observed at the onset of ACS.

**The authors state that they have no Conflict of Interest (COI).**

### Acknowledgement

We thank the patients who participated in the present study.

### References

- Roth GA, Johnson C, Abajobir A, et al. Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. *J Am Coll Cardiol* **70**: 1-25, 2017.
- Cui Y, Hao K, Takahashi J, et al. Age-specific trends in the incidence and in-hospital mortality of acute myocardial infarction over 30 years in Japan- report from the Miyagi AMI registry study. *Circ J* **81**: 520-528, 2017.
- Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* **894**: i-xii, 1-253, 2000.
- Mozaffarian D, Benjamin EJ, Go AS, et al. Executive summary: heart disease and stroke statistics-2016 update: a report from the American Heart Association. *Circulation* **133**: 447-454, 2016.
- Kannel WB, Abbott RD. Incidence and prognosis of unrecognized myocardial infarction. An update on the Framingham study. *N Engl J Med* **311**: 1144-1147, 1984.
- Yagi H, Komukai K, Hashimoto K, et al. Difference in risk factors between acute coronary syndrome and stable angina pectoris in the Japanese: smoking as a crucial risk factor of acute coronary syndrome. *J Cardiol* **55**: 345-353, 2010.
- JCS Joint Working Group, Japanese Society for Oral Health, Japanese Society of Oral and Maxillofacial Surgeons, et al. Guidelines for Smoking Cessation (JCS 2010)-digest version. *Circ J* **76**: 1024-1043, 2012.
- Chalon S, Moreno H Jr, Benowitz NL, Hoffman BB, Blaschke TF. Nicotine impairs endothelium-dependent dilatation in human veins *in vivo*. *Clin Pharmacol Ther* **67**: 391-397, 2000.
- Kumagai S, Amano T, Takashima H, et al. Impact of cigarette smoking on coronary plaque composition. *Coron Artery Dis* **26**: 60-65, 2015.
- JT's annual survey finds 17.9% of Japanese adults are smokers [Internet]. [cited 2019 Nov 1]. Available from: [https://www.jt.com/media/news/2018/pdf/20180730\\_02.pdf](https://www.jt.com/media/news/2018/pdf/20180730_02.pdf)
- McGill HC Jr, McMahan CA, Herderick EE, et al. Obesity accelerates the progression of coronary atherosclerosis in young men. *Circulation* **105**: 2712-2718, 2002.
- Lipids and lipoproteins in symptomatic coronary heart disease. Distribution, intercorrelations, and significance for risk classification in 6,700 men and 1,500 women. The Bezafibrate Infarction Prevention (BIP) Study Group, Israel. *Circulation* **86**: 839-848, 1992.
- Amano T, Matsubara T, Uetani T, et al. Impact of omega-3 polyunsaturated fatty acids on coronary plaque instability: an integrated backscatter intravascular ultrasound study. *Atherosclerosis* **218**: 110-116, 2011.
- Suzuki A, Ando H, Takashima H, et al. Effects of polyunsaturated fatty acids on periprocedural myocardial infarction after elective percutaneous coronary intervention. *EuroIntervention* **10**: 792-798, 2014.
- Wakabayashi Y, Funayama H, Ugata Y, et al. Low eicosapentaenoic acid to arachidonic acid ratio is associated with thin-cap fibroatheroma determined by optical coherence tomography. *J Cardiol* **66**: 482-488, 2015.
- Amano T, Matsubara T, Uetani T, et al. Lipid-rich plaques predict non-target-lesion ischemic events in patients undergoing percutaneous coronary intervention. *Circ J* **75**: 157-166, 2011.
- Domei T, Yokoi H, Kuramitsu S, et al. Ratio of serum n-3 to n-6 polyunsaturated fatty acids and the incidence of major adverse cardiac events in patients undergoing percutaneous coronary intervention. *Circ J* **76**: 423-429, 2012.
- Nishizaki Y, Shimada K, Tani S, et al. Significance of imbalance in the ratio of serum n-3 to n-6 polyunsaturated fatty acids in patients with acute coronary syndrome. *Am J Cardiol* **113**: 441-445, 2014.
- Hashimoto T, Ako J, Nakao K, et al. A lower eicosapentaenoic acid/arachidonic acid ratio is associated with in-hospital fatal arrhythmic events in patients with acute myocardial infarction: a J-MINUET substudy. *Heart Vessels* **33**: 481-488, 2018.
- Kurita A, Takashima H, Ando H, et al. Effects of eicosapentaenoic acid on peri-procedural (type IVa) myocardial infarction following elective coronary stenting. *J Cardiol* **66**: 114-119, 2015.
- Yamano T, Kubo T, Shiono Y, et al. Impact of eicosapentaenoic acid treatment on the fibrous cap thickness in patients with coronary atherosclerotic plaque: an optical coherence tomography study. *J Atheroscler Thromb* **22**: 52-61, 2015.
- Doi M, Nosaka K, Miyoshi T, et al. Early eicosapentaenoic acid treatment after percutaneous coronary intervention reduces acute inflammatory responses and ventricular arrhythmias in patients with acute myocardial infarction: a randomized, controlled study. *Int J Cardiol* **176**: 577-582, 2014.
- Endo J, Arita M. Cardioprotective mechanism of omega-3 polyunsaturated fatty acids. *J Cardiol* **67**: 22-27, 2016.
- Otsuka R, Kato Y, Imai T, Ando F, Shimokata H. Higher serum EPA or DHA, and lower ARA compositions with age independent fatty acid intake in Japanese aged 40 to 79. *Lipids* **48**: 719-727, 2013.
- Rees D, Miles EA, Banerjee T, et al. Dose-related effects of eicosapentaenoic acid on innate immune function in healthy humans: a comparison of young and older men. *Am J Clin Nutr* **83**: 331-342, 2006.
- Yagi S, Aihara K-i, Fukuda D, et al. Reduced ratio of eicosapentaenoic acid and docosahexaenoic acid to arachidonic acid is associated with early onset of acute coronary syndrome. *Nutr J* **14**: 2015.
- Jabbari R, Engström T, Glinge C, et al. Incidence and risk factors

of ventricular fibrillation before primary angioplasty in patients with first ST-elevation myocardial infarction: a nationwide study in Denmark. *J Am Heart Assoc* **4**: e000738, 2015.

- 28.** Bougouin W, Marijon E, Puymirat E, et al. Incidence of sudden cardiac death after ventricular fibrillation complicating acute myocardial infarction: a 5-year cause-of-death analysis of the FAST-MI 2005 registry. *Eur Heart J* **35**: 116-122, 2014.
- 29.** Doughty M, Mehta R, Bruckman D, et al. Acute myocardial in-

farction in the young-The University of Michigan experience. *Am Heart J* **143**: 56-62, 2002.

The Internal Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

---

© 2020 The Japanese Society of Internal Medicine  
*Intern Med* 59: 1125-1131, 2020