

RESEARCH LETTER

Open Access



Prehospital pulse oximetry: a red flag for early detection of silent hypoxemia in COVID-19 patients

Romain Jouffroy, Daniel Jost*  and Bertrand Prunet

The COVID-19 pandemic, which has been expanding since the first cases in Asia in late 2019, may result in acute respiratory failure (ARF) with severe hypoxemia [1–3]. In prehospital settings, the paucity of clinical respiratory signs has made assessing the severity of some COVID-19 patients challenging. Indeed, even though hypoxic ARF generally leads to an increase in respiratory rate (RR) [4], in some COVID-19 patients, a persistent normal RR was inconsistent with the severity of hypoxia.

Based on retrospective data, we aimed to describe the discrepancy between prehospital initial RR (RRi) and initial SpO₂ (SpO₂i; i.e., before oxygen supplementation, FiO₂ = 21%) in COVID-19 patients suffering from ARF.

We retrospectively examined consecutive COVID-19 patients suffering from ARF who were treated by the Paris Fire Brigade's basic life-support (BLS) teams in the prehospital setting. Data were provided from primary home care providers. Based on a previous study [5], we used the SpO₂i/RRi ratio as an estimator of the discrepancy insofar as a low numerator is associated with hypoxia, whereas a high denominator is typically associated with respiratory failure.

After having measured the SpO₂i/RRi values in COVID-19 patients, we compared them to those of non-COVID-19 patients (i.e., patients with other causes of ARF treated by the BLS teams over the previous 3 years in the same period).

Continuous data were described as median (interquartile range) and were compared by applying the Kruskal-Wallis test. The French Society of Anaesthesia and

Intensive Care approved the trial protocol on April 7, 2020 (IRB 00010254-2020-055).

The study examined 1201 patients who experienced COVID-19 between March 13 and 29, 2020. The median SpO₂i/RRi value was significantly higher than that of patients treated in the previous 3 years (5 [4, 5] in 2020 versus 3.4 [2.4–4.5] in 2019, 3.3 [2.2–4.4] in 2018, and 3.5 [2.5–4.6] in 2017, $p < 0.001$, Fig. 1).

In summary, this retrospective study based on prehospital first responder data highlighted a relatively higher discrepancy between SpO₂i and RRi in COVID-19 ARF patients, in comparison with previous non-COVID-19 ARF patients. Without a systematic SpO₂i measurement, a normal breathing rate could mask profound hypoxia and make severity assessment in COVID-19 patients all the more difficult in an out-of-hospital setting.

Despite differences in worldwide prehospital emergency medical services, pulse oximetry is an accessible tool that prehospital healthcare providers can easily use.


In conclusion, prehospital pulse oximetry might be used as a red flag for early detection of “silent hypoxemia” in COVID-19 patients. The prehospital SpO₂i/RRi ratio needs further investigation because it might help to identify non-clinically obvious ARFs.

* Correspondence: Daniel.jost@pompierparis.fr

Paris Fire Brigade, Emergency Medicine Department, 1 Place Jules Renard, 75017 Paris, France



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



Acknowledgements

Thanks to the Paris Fire Brigade Basic Life Support Teams for their daily engagement during the pandemic.

Authors’ contributions

All authors have made substantial contributions to the conception and design of the letter, drafting the article or revising it critically for relevant intellectual content, and final approval of the version to be submitted. The authors read and approved the final manuscript

Funding

None

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

As a retrospective study design, the study protocol has been submitted to the Ethics Committee of “The French Society of Anaesthesia and Intensive Care.” The trial protocol was approved on April 7, 2020 (IRB 00010254-2020-055).

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Received: 20 May 2020 Accepted: 27 May 2020

Published online: 08 June 2020

References

1. Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis.* 2020;14(7):3099-3107.
2. Onder G, Rezza G, Brusaferro S. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA.* 2020. <https://doi.org/10.1001/jama.2020.4683> Accessed on May 15, 2020.
3. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020;382:1708–20.
4. Braman SS. The regulation of normal lung function. *Allergy Asthma Proc.* 1995;16:223–6.
5. Roca O, Messika J, Caralt B, et al. Predicting success of high-flow nasal cannula in pneumonia patients with hypoxemic respiratory failure: the utility of the ROX index. *J Crit Care.* 2016;35:200–5.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.