



# Pathogenic variants in *KPTN* gene identified by clinical whole-genome sequencing

Isabelle Thiffault,<sup>1,2,3</sup> Andrea Atherton,<sup>4</sup> Bryce A. Heese,<sup>4</sup> Ahmed T. Abdelmoity,<sup>4</sup> Kailash Pawar,<sup>4</sup> Emily Farrow,<sup>1,3,4</sup> Lee Zellmer,<sup>1</sup> Neil Miller,<sup>1</sup> Sarah Soden,<sup>1,3,4</sup> and Carol Saunders<sup>1,2,3</sup>

<sup>1</sup>Center for Pediatric Genomic Medicine, Children's Mercy Hospital, Kansas City, Missouri 64108, USA; <sup>2</sup>Department of Pathology and Laboratory Medicine, Children's Mercy Hospitals, Kansas City, Missouri 64108, USA; <sup>3</sup>University of Missouri–Kansas City School of Medicine, Kansas City, Missouri 64108, USA; <sup>4</sup>Department of Pediatrics, Children's Mercy Hospitals, Kansas City, Missouri 64108, USA

**Abstract** Status epilepticus is not rare in critically ill intensive care unit patients, but its diagnosis is often delayed or missed. The mortality for convulsive status epilepticus is dependent on the underlying aetiologies and the age of the patients and thus varies from study to study. In this context, effective molecular diagnosis in a pediatric patient with a genetically heterogeneous phenotype is essential. Homozygous or compound heterozygous variants in *KPTN* have been recently associated with a syndrome typified by macrocephaly, neurodevelopmental delay, and seizures. We describe a comprehensive investigation of a 9-yr-old male patient who was admitted to the intensive care unit, with focal epilepsy, static encephalopathy, autism spectrum disorder, and macrocephaly of unknown etiology, who died of status epilepticus. Clinical whole-genome sequencing revealed compound heterozygous variants in the *KPTN* gene. The first variant is a previously characterized 18-bp in-frame duplication (c.714\_731dup) in exon 8, resulting in the protein change p.Met241\_Gln246dup. The second variant, c.394 + 1G > A, affects the splice junction of exon 3. These results are consistent with a diagnosis of autosomal recessive *KPTN*-related disease. This is the fourth clinical report for *KPTN* deficiency, providing further evidence of a wider range of severity.

Corresponding author:  
csaunders@cmh.edu

© 2020 Thiffault et al. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial License, which permits reuse and redistribution, except for commercial purposes, provided that the original author and source are credited.

**Ontology terms:** complex febrile seizures; delayed fine motor development; delayed gross motor development; epileptic encephalopathy; intellectual disability, severe; macrocephaly at birth; polymorphic focal epileptiform discharges

Published by Cold Spring Harbor Laboratory Press

doi:10.1101/mcs.a003970

[Supplemental material is available for this article.]

## INTRODUCTION


In recent years, next-generation sequencing (NGS) technologies have revolutionized approaches in clinical genetics. Whole-exome sequencing (WES) or whole-genome sequencing (WGS) allows diagnoses in many patients with complex phenotypes and unusual clinical presentations. As the cost of NGS falls, it has become feasible to use this powerful technology in clinical care, simultaneously unraveling variations in about 19,000 genes. Technological advances have led to the ability to sequence, analyze, and interpret entire genome data in a timely manner, clearly changing the diagnostic paradigm and proving to be cost-effective in many cases (Soden et al. 2014). Clinical diagnostic sequencing currently focuses on identifying causal mutations in the exome (~1% of the genome), where most disease-causing mutations are known to occur. WGS permits analysis of coding regions as well as regulatory elements that control gene expression; however, noncoding variants

remained largely unexplored in clinical diagnostics because of the interpretive challenges (Warman Chardon et al. 2015). Although it lacks the depth of coverage of an exome, WGS can have more reliable and uniform sequence coverage, particularly in regions of the genome with low sequence complexity or high GC-rich content. Both WGS and WES approaches have benefits and limitations, but the limitation of NGS gene panels and WES should be considered prior to clinical testing (Williams et al. 2008; Soden et al. 2014; Ankala et al. 2015; Warman Chardon et al. 2015). In this context, the present work demonstrates the application of clinical WGS in the pediatric population in which we provided a fast, accurate, and cost-effective molecular diagnosis in a pediatric patient with a genetically heterogeneous phenotype.

## RESULTS

### Clinical Presentation

A 9-yr-old Caucasian male was admitted to the Children’s Mercy Hospital pediatric intensive care unit (PICU) for status epilepticus and further evaluation of a suspected underlying genetic condition because of his history of macrocephaly, intractable epilepsy, autism, severe developmental delays, hypotonia, and hypoglycemia. Dysmorphic features include frontal bossing, sunken eye sockets, downslanting palpebral fissures, small ears, thin upper lip, and small nose (Fig. 1). The prenatal history was unremarkable: Delivery was at full term, and neonatal development was normal. He was the third child for his parents together, born to a 26-yr-old mother and 30-yr-old father. Family history was noncontributory. At birth, he weighed 2.71 kilograms and was 49.5 cm long. His head circumference was noted to be



**Figure 1.** Our patient was a 9-yr-old male with epilepsy, static encephalopathy, autism spectrum disorder, and (A) macrocephaly of unknown etiology who died of status epilepticus. (B) Several facial dysmorphisms were found, including frontal bossing, sunken eye sockets, downslanting palpebral fissures, small ears, thin upper lip, and small nose. (C) At 4 yr old, he was normal height and weight, but his head circumference measured 57 cm (>98th percentile).

greater than the 99th percentile at birth. At 4 yr old, he was of average height and weight, but his head circumference measured 57 cm (>98th percentile, +4SD). He had poor tracking and eye contact but a normal ophthalmologic exam. A head CT was completed at 2 mo of age and reported to be normal. At 3 mo of age, he started to have partial seizures but EEG was normal. At 5 mo of age he was noted to have hepatosplenomegaly accompanied by hypoglycemic episodes. Developmentally, he smiled socially at 2 mo, rolled at 8 mo, sat without assistance at 10–11 mo, crawled at 14 mo, and walked at 2.5 yr. A brain MRI was normal at 4 mo of age; however a second, performed at 9 yr of age, was concerning for a possible arteriovenous malformation, with abnormal hyperintense T2/FLAIR signals in the subcortical white matter of the right anteromedial temporal lobe. An extensive etiologic workup was normal, including high-resolution karyotype, CGH microarray, *PTEN* sequencing, fragile X, enzyme analysis for Gaucher, mucopolysaccharidosis I and II, plasma amino acids, pristanic and phytanic acids, very long chain fatty acid (VLCFA), and urine organic acids. Mild elevations of C14 species on an acylcarnitine profile in 2010 were detected in conjunction with ketotic hypoglycemia and were likely associated with physiologic response to fasting. Previously, the seizures occurred at a frequency of 1–2 per hour, consisting of focal tonic-clonic seizures involving his left or right side of the body but occasionally progressed to status epilepticus, and a pentobarbital infusion was started with the goal to acquire burst suppression. Unfortunately, this was unsuccessful, and he passed away at age 9.

### Genetic Analysis

Clinical WGS was performed on the patient and WES was performed on his healthy mother, following informed consent. The patient was compound heterozygous for two pathogenic variants in the *KPTN* gene (Table 1). This gene was ranked 351st in the Phenomizer gene list. The first variant identified was an 18-bp in-frame duplication (c.714\_731dup ACCGACCACATCTGCAGA; rs587777148) in exon 8, resulting in the protein change p.Met241\_Gln246dup. This variant has been previously reported in *trans* with a second truncating variant in multiple affected individuals from two Amish families (Baple et al. 2014). Prior *in vitro* transfection studies indicated the mutant protein is mislocalized and accumulates in neurons of affected individuals, leading to a dominant negative effect (Baple et al. 2014). The p.Met241\_Gln246dup variant is not found in population databases such as dbSNP, Exome Variant Database, or ExAC but was found in 1 of 560 Amish control chromosomes (Baple et al. 2014); 136/275574 in gnomAD data set (0.05%). The second variant, c.394 + 1G > A, affects the splice junction of exon 3. This variant has not been reported in affected individuals but is predicted to cause aberrant splicing. This variant was observed in ~0.01% individuals of European ancestry in the NHLBI Exome Sequencing Project and Exome Aggregation Consortium (ExAC); 19/277090 in gnomAD data set (0.007%). The c.394 + 1G > A variant, but not the c.714\_731dup variant, was inherited from the mother, suggesting these variants are *in trans*. This genotype was confirmed by Sanger sequencing and is consistent with a diagnosis of autosomal recessive *KPTN*-related disease. In addition, two variants of unknown significance were reported ((*CHD2*-NM\_001271.3:c.5268G > C (p.Gln1756His); *CTDP1*-NM\_004715.4:c.1219T > C (p.Trp407Arg)). No incidental findings were reported for this case. The turnaround time was 33 d.

### DISCUSSION

The use of NGS techniques by clinical laboratories has risen tremendously and has greatly facilitated the elucidation of the etiologic diagnosis in patients suspected of having a genetic disease. The various approaches, including WES, WGS, and targeted panels, all have benefits and limitations. Targeted panels offer a specific list of genes relevant to the clinical

**Table 1.** Variant table

Gene	Chromosome	HGVS DNA reference	HGVS protein reference	Variant type	Predicted effect (substitution, deletion, etc.)	dbSNP/dbVar ID	Genotype	ClinVar ID	Parent of origin	Comments
KPTN	19:47986551-47986551 C>T	NM_007059.2:c.394+1G>A	p.?	LoF	n/a	rs373139784	Het.	499654	Maternally inherited	gnomAD Swedish, (12/26104), 0.046%
KPTN	19:47983175-47983176 → ACCGACCACATCT GCAGA	NM_007059.2:c.714_731dupTCTGCAGATGTGG	p.Leu239_Val244dup	In-frame	blorum: 4	rs1399298568	Het.	100680	Paternally inherited	gnomAD Swedish, (46/25962), 0.177%
CHD2	15:93567716-93567716 G>C	NM_001271.3:c.5268G>C	p.Gln1756His	Missense	SIFT: tolerated_low_confidence (0.33)					
PolyPhen-2: possibly_damaging (0.496)	rs201950393	Het.	ClinVar 377653	Maternally inherited		gnomAD Ashkenazi Jewish, (9/10362), 0.0869%				
CTDP1	18:77474679-77474679 T>C	NM_004715.4:c.1219T>C	p.Trp407Arg	Missense	SIFT: tolerated (0.59)					
PolyPhen-2: benign (0.007)	rs149090172	Het.	n/a	Maternally inherited		gnomAD Other, (12/6882), 0.174%				

phenotype in question and often boast higher coverage than WES/WGS. However, whether this improves sensitivity, particularly in Mendelian disorders, is not clear. In addition, a wide range of interlaboratory variability exists for gene lists offered for the same condition. For instance, commercial clinical NGS panels for macrocephaly include anywhere from 12 to 44 genes. Although targeted panels may appear to be a more economical approach to WES/WGS, this is only the case if the gene relevant to the patient being tested is present on the panel, which is often difficult to know a priori, particularly in patients with nonspecific symptoms such as intellectual disability or seizures. In many cases, serial testing of additional genes and panels quickly surpasses the expense of WES/WGS (Soden et al. 2014). In addition, targeted panels may fail to incorporate newly discovered disease-associated genes. For example, new genes associated with both syndromic and nonsyndromic macrocephaly have been identified, further expanding the genetic heterogeneity to more than 165 conditions, including 17 metabolic disorders, associated with macrocephaly (Williams et al. 2008; Keppler-Noreuil et al. 2014; Koh et al. 2014; Marchese et al. 2014; Sugathan et al. 2014; Nevado et al. 2015; Nguyen et al. 2015). One such gene, *KPTN*, encoding kaptin, was first reported in January of 2014 in patients with a syndrome typified by macrocephaly, neurodevelopmental delay, and seizures (MRT41; OMIM# 615637) (Baple et al. 2014; Pajusalu et al. 2015). *KPTN* is not currently offered in any clinical panels for macrocephaly in the United States. Of the clinical laboratories listed in the NextGxDx website, only two of 16 offer *KPTN* as part of an autism/intellectual disability panel, and only two of 17 include it as part of an epilepsy panel. The comparison of such gene lists is difficult for clinicians and the curation is onerous for the clinical laboratory to manage. Clinical WES/WGS removes the guesswork as far as which gene to include, as all genes relevant to the patient's phenotype are queried in the analysis process. In the current case, by using clinical WGS in a 9-yr-old Caucasian male admitted in pediatric intensive care unit for status epilepticus, a diagnosis of *KPTN*-related disease was made.

A comparison of the individual clinical signs in patients reported with *KPTN*-related disease are shown in Table 2. The most unifying features are macrocephaly (92%), developmental delay (100%), and intellectual disability (100%). Recurrent dysmorphic features included frontal bossing, abnormal head shape, and prominent chin. With the exception of the p.Met241\_Gln246dup, which is predicted to be nonfunctional by producing a misfolded altered protein, all patients have truncating variants (p.S259\*; p.S223Qfs\*50; p.S200Ifs\*55); no patients were homozygous for the p.Met241\_Gln246dup. RT-PCR or western blot experiments were not performed to assess if those variants result in nonsense-mediated mRNA decay or whether a truncated protein (lacking the carboxy-terminal amino acids) is produced. The second variant uncovered in our patient, c.394 + 1G > A, is located upstream of the two previously reported truncating variants. This is the first splicing variant reported in *KPTN*, it is likely to result in nonsense-mediated mRNA decay, which may explain the more severe outcome. Additional external factors such as modifier genes could also influence the phenotype. Of the 12 previous patients described, seven have survived into their second and third decades, with two having expired at ages 29 and 30 to a head injury and pneumonia (Table 2; Baple et al. 2014; Pajusalu et al. 2015; Lucena et al. 2020). At this time, limited data exist for genotype–phenotype correlations; further phenotyping and natural history of disease study are needed to understand the clinical spectrum and prognosis of *KPTN*-related syndrome.

This is the fourth clinical report for *KPTN*-deficiency, the 13th patient but the first to die in childhood with status epilepticus, providing further evidence of a wider range of severity.

Several studies have shed new light on the molecular and cellular processes that orchestrate the human neuronal circuitry and is defective in neurological disorders (Ropers et al. 2011; Riviere et al. 2012; Baple et al. 2014; Koh et al. 2014; Sugathan et al. 2014).

**Table 2.** A comparison of clinical findings of affected individuals reported with pathogenic variants in the KPTN gene

Origin	Amish, Ohio				Amish, Ohio				Estonia			Mixed, Kansas*			
	Homozygous p.S259*				p.S259* /p.M241Q246Dup				Homozygous p.S223Qfs*50			c.394+1G>A /p.M241Q 246Dup		#	%
Patients	1	2	3	4	5	6	7	8	9	10	11	12	12	n = 12	
Gender	M	M	M	M	M	F	F	M	F	M	F	M	M		
Age at evaluation (years)	28.2	29	30	16.5	13.2	22.7	24.9	1.4	7.8	32	24	4	4		
Age at death (years)	n.a	29 (head injury)	30 (pneumonia)	n.a	n.a	n.a	n.a	n.a	n.a	n.a	n.a	9	9		
<b>Growth parameters</b>															
Birth weight, kg (SD)	2.95 (0.7)	2.92 (0)	3.46 (+1.3)	1.59 (-0.2)	3.35 (-0.4)	2.89 (+1.2)	3.16 (-0.5)	3.2 (+2.0)	2.75 (+1.1)	?	?	?	3.13		
Birth OFC, cm (SD) or birth macrocephaly reported	?	40.6 (+2.9)	35.6 (+1.75)	?	51 (+3.9)	?	?	macrocephaly	macrocephaly	34 (N)	37 (N)	macrocephaly	5	41.7	
Height, cm (SD) at evaluation	166.7 (-1.6)	?	165.1 (-1.8)	169 (-0.7)	161.6 (+0.6)	156.2 (-1.3)	160.0 (-0.6)	?	123.5 (-0.4)	N	N	100.0 (25th)			
Weight, kg (SD)	121.2 (+3.4)	?	66.2 (-0.2)	63.1 (+0.1)	51.5 (+0.8)	107.9 (+3.6)	82 (+2.2)	?	?	N	N	16.8 (50th)			
OFC, cm (SD)	62 (+3.0)	?	63.5 (+3.6)	62.5 (+3.4)	61 (+3.3)	63 (+5.4)	60 (+3.2)	52.5 (+3.2)	55.5 (+2.1)	63 (+4.5)	60 (+4.0)	57.0 (98th)	11	91.7	
Parental OFC, cm (SD)	?	Mother 55.5 (+0.1); Father 60 (+1.7)	?	Mother 55.5 (+0.1); Father 60 (+1.7)	?	Mother 58.5 (2.3)	?	Mother 57 (+1.2); Father 59.5 (+1.5)	Mother 57 (+1.2); Father 59.5 (+1.5)	?	?	N			
<b>Development</b>															
Walked (years)	1	1	1.33	1.9	4	3.8	2.4	>2.2	2.2	n.a	n.a	2.5			
Expressive and receptive	■	■	■	■	■	■	■	■	■	■	■	■	■	12	100.0
Language deficit	■	■	■	■	■	■	■	■	■	■	■	■	■	12	100.0
Intellectual disability	■	■	■	■	■	■	■	■	■	■	■	■	■	12	100.0
Hearing	N	N	N	N	N	N	N	N	N	N	N	N	N	0	0.0
<b>Neurology</b>															
Childhood hypotonia	-	-	-	■	■	■	■	■	■	?	?	■	■	7	58.3
Seizures (onset)	AS/ GTCS (3 mo)	AS/GTCS (7 yr)	GTCS (7 yr)	GTCS (7 yr)	GTCS (7 yr)	GTCS (7 yr)	GTCS (7 yr)	GTCS (7 yr)	GTCS (7 yr)	GTCS (10 mo)	GTCS (10 mo)	GTCS (3 mo)	5	41.7	
Neuroimaging	N	?	Ventriculomegaly	?	?	?	?	Widening of metopic suture	N	N	N	Abnormal**	3	25.0	
<b>Behavioral characteristics</b>															
Repetitive speech	■	-	-	-	■	■	■	■	-	■	-	■	■	6	50.0
Stereotypies	-	-	■	■	■	■	■	-	-	■	-	■	■	7	58.3
Hyperactivity	-	-	-	■	-	-	-	-	-	■	-	■	■	3	25.0
Autistic features	?	?	?	?	?	?	?	?	?	■	■	■	■	3	25.0
Anxiety	■	■	■	■	■	■	■	■	■	■	■	■	■	10	83.3

(Continued on next page.)

**Table 2.** (Continued)

Origin	Amish, Ohio			Amish, Ohio			Amish, Ohio			Estonia		Mixed, Kansas*	
	Homozygous p.S259*			p.S259* /p.M241Q246Dup			p.S259* /p.M241Q246Dup			Homozygous p.S223Qfs*50		c.394 +1G > A /p.M241Q 246Dup	
Genotype	1	2	3	4	5	6	7	8	9	10	11	12	n = 12
Patients													#
													%
Phobia	-	-	-	-	-	-	-	■	-	?	?	-	1 8.3
<b>Physical anomalies</b>													
Frontal bossing	■	■	■	■	■	■	■	■	■	■	■	■	12 100.0
High palate	?	?	?	?	?	?	?	?	?	■	■	-	2 16.7
Hypertelorism	-	-	-	■	-	-	-	■	-	-	-	-	2 16.7
Plagiocephaly	-	-	-	■	■	-	■	■	-	-	-	-	4 33.3
Prominent chin	■	■	■	■	-	-	-	-	■	■	■	■	7 58.3
Hepatosplenomegaly	-	-	-	-	-	-	■	-	-	-	-	■	2 16.7
Splenomegaly	-	-	-	-	-	-	-	■	-	-	-	-	1 8.3
Liver cirrhosis	-	-	-	-	-	-	■	-	-	-	-	-	1 8.3
Anemia	-	-	-	-	-	-	-	■	-	-	-	-	1 8.3
Recurrent Infections	-	-	■	-	-	-	■	■	-	?	?	■	4 33.3
5th finger clinodactily	■	■	■	-	-	-	-	-	-	-	-	-	3 25.0
Fetal finger pad	-	-	-	-	-	-	-	-	■	-	--	-	1 8.3
Reference	Baple et al. 2014	Baple et al. 2014	Baple et al. 2014	Baple et al. 2014	Baple et al. 2014	Baple et al. 2014	Baple et al. 2014	Baple et al. 2014	Baple et al. 2014	Pajusalu et al. 2015	Pajusalu et al. 2015	n.a.	

(F) Female, (M) male, (OFC) occipitofrontal circumference, (SD) standard deviation score, (N/P) not performed, (■) indicates presence of the clinical features in an affected individual, (-) indicates absence of the clinical features in an affected individual, (?) indicates the presence or absence of clinical features in an affected individual is unknown, (AS) absence of seizures, (GTC) generalized tonic-clonic seizures, (N) normal, (n.a.) not applicable.



Evidence implicating defects in proteins involved in actin dynamics, microtubule homeostasis, and vesicle trafficking in developmental disorders via effects on neuronal development and migration and brain architecture has accumulated over the past decade. The role of microtubule homeostasis (*LIS1* and *DCX*) or tubulin (*TUBA1A*, *TUBA8*, *TUBB2A*, *TUBB4A*, *TUBB2B*, *TUBB3*, and *TUBB* and *TBCE*) genes has been established in several brain disorders. More recently, the actin-encoding genes *ACTB* and *ACTG1* have been shown to be involved in brain malformations causing Baraitser–Winter syndrome (OMIM # 243310) (Riviere et al. 2012). Additional intellectual disability genes involved in actin dynamics and vesicle trafficking include *STXBP1* and *SYP* (Barcia et al. 2013; Kato 2015). Patients with *KPTN* deficiency consistently exhibit macrocephaly, intellectual disability, and developmental delay. Furthermore, *KPTN* is a novel actin binding protein, enriched in neuronal growth cones and cortical sites of neurons at early developmental stages, likely playing a central role in modulating neuron morphology and growth. Although genes affecting many of these pathways are associated with hearing loss, to date, no patients with *KPTN*-related disease, including our patient, have been reported with deafness. As with several other published studies (Riviere et al. 2012; Soden et al. 2014; Ankala et al. 2015; Warman Chardon et al. 2015; Willig et al. 2015), WGS was found to be a rapid and cost-efficient approach for molecular diagnostic of a genetically heterogeneous condition (Thiffault et al. 2019). Thus, this report confirms the occurrence of *KPTN*-related syndrome outside of the Amish population and demonstrates the variability in the phenotypic spectrum and severity. Pathological mechanisms of abnormal neuronal actin cytoskeleton and discrepancy between the underlying phenotypes caused by *KPTN* deficiency remain to be elucidated.

## METHODS

Genomic DNA was extracted from peripheral blood mononuclear cells using a Chemagen MSM1 robot (PerkinElmer). WGS was prepared using the Kapa Hyper library prep omitting PCR. Sequencing was completed on an Illumina HiSeq 2500 instrument utilizing paired end 2 × 125 base pair reads with v4 Chemistry (Illumina). The proband's sample was sequenced to a depth of 115.16 Gb for a mean coverage of ~37× (Supplemental Table S1). Bidirectional sequence was assembled, aligned to reference gene sequences based on human genome build GRCh37/UCSC hg19, and analyzed using custom-developed software, RUNES, and VIKING (Saunders et al. 2012; Soden et al. 2014; Willig et al. 2015). Variants were filtered to a 1% minor allele frequency and then prioritized by the American College of Medical Genetics (ACMG) categorization (Richards et al. 2015), OMIM identity, and phenotypic assessment. Alignments were viewed using Integrative Genomic Viewer software version 2.3.8 (IGV; Broad Institute). A candidate gene list was generated by Phenomizer using Human Phenotype Ontology (Kohler et al. 2014) terms: Macrocephaly (HPO:0000256), Muscular hypotonia (HPO:0001252), Autism (HPO:0000717), Seizures (HPO:0001250), Global developmental delay (HPO:0001263), and Abnormal facial shape (HPO: 0001999), with a cutoff at *P*-value of 0.5. This gene list contained approximately 700 genes and was imported into VIKING to guide the analysis; however, a separate analysis, not limited by this gene list, was performed in parallel to look for relevant pathogenic genotypes in genes not included in the HPO gene list. Pathogenic, likely pathogenic, and variants of unknown significance in HPO genes were reported; likely benign and benign variants are not reported but are made available upon request. Incidental findings in the 56 genes recommended by ACMG were requested by the family and analyzed for pathogenic and likely pathogenic variants only.



## ADDITIONAL INFORMATION

### Data Deposition and Access

Our patient consent does not permit patient sequence data to be uploaded to a data repository. The variants reported have been deposited in the ClinVar database (<https://www.ncbi.nlm.nih.gov/clinvar/>) and can be found under accession numbers VCV000100680.3, VCV000499654.1, and VCV000377653.5.

### Ethics Statement

Written informed consent was obtained from the patient's legal guardians for publication of this case report and governed by the Institutional Review Board of Children's Mercy Hospitals and Clinics. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

### Acknowledgments

We are very grateful to this family for allowing this case to be published. We thank our colleagues in the Center for Pediatric Genomic Medicine and Children's Mercy Hospital in Kansas City. This work was not supported by a grant.

### Competing Interest Statement

The authors have declared no competing interest.

### Referees

Sander Pajusalu  
Anonymous

Received January 7, 2019;  
accepted in revised form  
April 16, 2020.

### Author Contributions

C.J.S., E.F., I.T., L.Z., and S.S. conceived and designed the experiments; C.J.S., E.F., and I.T. performed the experiments; N.M. and E.F. contributed reagents/materials/analysis tools; I.T. and C.J.S. wrote the paper; E.F., S.S., L.Z., B.H., A.A., and N.M. reviewed the manuscript; K.P. reviewed the manuscript and contributed to the clinical investigation of the patient; and A.A., B.H., and A.T.A. contributed to the recruitment and clinical investigations of the patient for the study.

## REFERENCES

- Ankala A, da Silva C, Gualandi F, Ferlini A, Bean LJ, Collins C, Tanner AK, Hegde MR. 2015. A comprehensive genomic approach for neuromuscular diseases gives a high diagnostic yield. *Ann Neurol* **77**: 206–214. doi:10.1002/ana.24303
- Baple EL, Maroofian R, Chioza BA, Izadi M, Cross HE, Al-Turki S, Barwick K, Skrzypiec A, Pawlak R, Wagner K, et al. 2014. Mutations in *KPTN* cause macrocephaly, neurodevelopmental delay, and seizures. *Am J Hum Genet* **94**: 87–94. doi:10.1016/j.ajhg.2013.10.001
- Barcia G, Barnerias C, Rio M, Siquier-Pernet K, Desguerre I, Colleaux L, Munnich A, Rotig A, Nabbout R. 2013. A novel mutation in *STXBP1* causing epileptic encephalopathy (late onset infantile spasms) with partial respiratory chain complex IV deficiency. *Eur J Med Genet* **56**: 683–685. doi:10.1016/j.ejmg.2013.09.013
- Kato M. 2015. Genotype-phenotype correlation in neuronal migration disorders and cortical dysplasias. *Front Neurosci* **9**: 181. doi:10.3389/fnins.2015.00181
- Keppler-Noreuil KM, Sapp JC, Lindhurst MJ, Parker VE, Blumhorst C, Darling T, Tosi LL, Huson SM, Whitehouse RW, Jakkula E, et al. 2014. Clinical delineation and natural history of the *PIK3CA*-related overgrowth spectrum. *Am J Med Genet A* **164A**: 1713–1733. doi:10.1002/ajmg.a.36552
- Koh JY, Lim JS, Byun HR, Yoo MH. 2014. Abnormalities in the zinc-metalloprotease-BDNF axis may contribute to megalencephaly and cortical hyperconnectivity in young autism spectrum disorder patients. *Mol Brain* **7**: 64. doi:10.1186/s13041-014-0064-z
- Kohler S, Doelken SC, Mungall CJ, Bauer S, Firth HV, Bailleul-Forestier I, Black GC, Brown DL, Brudno M, Campbell J, et al. 2014. The Human Phenotype Ontology project: linking molecular biology and disease through phenotype data. *Nucleic Acids Res* **42**: D966–D974. doi:10.1093/nar/gkt1026
- Lucena PH, Armani-Franceschi G, Bispo-Torres AC, Bandeira ID, Lucena MFG, Maldonado I, Veiga MF, Miguel D, Lucena R. 2020. *KPTN* gene homozygous variant-related syndrome in the northeast of Brazil: a case report. *Am J Med Genet A* **182**: 762–767. doi:10.1002/ajmg.a.61492

- Marchese M, Conti V, Valvo G, Moro F, Muratori F, Tancredi R, Santorelli FM, Guerrini R, Sicca F. 2014. Autism-epilepsy phenotype with macrocephaly suggests *PTEN*, but not *GLIALCAM*, genetic screening. *BMC Med Genet* **15**: 26. doi:10.1186/1471-2350-15-26
- Nevado J, Rosenfeld JA, Mena R, Palomares-Bralo M, Vallespin E, Angeles Mori M, Tenorio JA, Gripp KW, Denenberg E, Del Campo M, et al. 2015. *PIAS4* is associated with macro/microcephaly in the novel interstitial 19p13.3 microdeletion/microduplication syndrome. *Eur J Hum Genet* doi:10.1038/ejhg.2015.51
- Nguyen LS, Schneider T, Rio M, Moutton S, Siquier-Pernet K, Verny F, Boddaert N, Desguerre I, Munich A, Rosa JL, et al. 2015. A nonsense variant in *HERC1* is associated with intellectual disability, megalencephaly, thick corpus callosum and cerebellar atrophy. *Eur J Hum Genet* doi:10.1038/ejhg.2015.140
- Pajusalu S, Reimand T, Ounap K. 2015. Novel homozygous mutation in *KPTN* gene causing a familial intellectual disability-macrocephaly syndrome. *Am J Med Genet A* **167A**: 1913–1915. doi:10.1002/ajmg.a.37105
- Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, et al. 2015. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* **17**: 405–424. doi:10.1038/gim.2015.30
- Riviere JB, van Bon BW, Hoischen A, Kholmanskikh SS, O’Roak BJ, Gilissen C, Sullivan CT, Christian SL, Abdul-Rahman OA, Atkin JF, et al. 2012. De novo mutations in the actin genes *ACTB* and *ACTG1* cause Baraitser–Winter syndrome. *Nat Genet* **44**: 440–444. doi:10.1038/ng.1091
- Ropers F, Derivery E, Hu H, Garshasbi M, Karbasiyan M, Herold M, Nürnberg G, Ullmann R, Gautreau A, Sperling K, et al. 2011. Identification of a novel candidate gene for non-syndromic autosomal recessive intellectual disability: the WASH complex member *SWIP*. *Hum Mol Genet* **20**: 2585–2590. doi:10.1093/hmg/ddr158
- Saunders CJ, Miller NA, Soden SE, Dinwiddie DL, Noll A, Alnadi NA, Andraws N, Patterson ML, Krivohlavek LA, Fellis J, et al. 2012. Rapid whole-genome sequencing for genetic disease diagnosis in neonatal intensive care units. *Sci Transl Med* **4**: 154ra135. doi:10.1126/scitranslmed.3004041
- Soden SE, Saunders CJ, Willig LK, Farrow EG, Smith LD, Petrikin JE, LePichon JB, Miller NA, Thiffault I, Dinwiddie DL, et al. 2014. Effectiveness of exome and genome sequencing guided by acuity of illness for diagnosis of neurodevelopmental disorders. *Sci Transl Med* **6**: 265ra168. doi:10.1126/scitranslmed.3010076
- Sugathan A, Biagioli M, Golzio C, Erdin S, Blumenthal I, Manavalan P, Ragavendran A, Brand H, Lucente D, Miles J, et al. 2014. *CHD8* regulates neurodevelopmental pathways associated with autism spectrum disorder in neural progenitors. *Proc Natl Acad Sci* **111**: E4468–E4477. doi:10.1073/pnas.1405266111
- Thiffault I, Farrow E, Zellmer L, Berrios C, Miller N, Gibson M, Caylor R, Jenkins J, Faller D, Soden S, et al. 2019. Clinical genome sequencing in an unbiased pediatric cohort. *Genet Med* **21**: 303–310. doi:10.1038/s41436-018-0075-8
- Warman Chardon J, Beaulieu C, Hartley T, Boycott KM, Dymont DA. 2015. Axons to exons: the molecular diagnosis of rare neurological diseases by next-generation sequencing. *Curr Neurol Neurosci Rep* **15**: 64. doi:10.1007/s11910-015-0584-7
- Williams CA, Dagli A, Battaglia A. 2008. Genetic disorders associated with macrocephaly. *Am J Med Genet A* **146A**: 2023–2037. doi:10.1002/ajmg.a.32434.
- Willig LK, Petrikin JE, Smith LD, Saunders CJ, Thiffault I, Miller NA, Soden SE, Cakici JA, Herd SM, Twist G, et al. 2015. Whole-genome sequencing for identification of Mendelian disorders in critically ill infants: a retrospective analysis of diagnostic and clinical findings. *Lancet Respir Med* **3**: 377–387. doi:10.1016/S2213-2600(15)00139-3