


Article

CORDITE: The Curated CORona Drug InTERactions Database for SARS-CoV-2



Roman Martin,
Hannah F. Löchel,
Marius Welzel,
Georges Hattab,
Anne-Christin
Hauschild,
Dominik Heider

dominik.heider@uni-marburg.de

HIGHLIGHTS

CORDITE aggregates all available knowledge on potential drugs for SARS-CoV-2

CORDITE represents the largest curated database for drugs and interactions

CORDITE makes the data accessible via a webserver and an open API

CORDITE enables easy meta-analyses, design of new clinical studies, or literature search

Martin et al., iScience 23,
101297
July 24, 2020 © 2020 The
Authors.
<https://doi.org/10.1016/j.isci.2020.101297>



Article

CORDITE: The Curated CORona Drug InTERactions Database for SARS-CoV-2

Roman Martin,^{1,2} Hannah F. Löchel,^{1,2} Marius Welzel,¹ Georges Hattab,¹ Anne-Christin Hauschild,¹ and Dominik Heider^{1,3,*}

SUMMARY

Since the outbreak in 2019, researchers are trying to find effective drugs against the SARS-CoV-2 virus based on de novo drug design and drug repurposing. The former approach is very time consuming and needs extensive testing in humans, whereas drug repurposing is more promising, as the drugs have already been tested for side effects, etc. At present, there is no treatment for COVID-19 that is clinically effective, but there is a huge amount of data from studies that analyze potential drugs.

We developed CORDITE to efficiently combine state-of-the-art knowledge on potential drugs and make it accessible to scientists and clinicians. The web interface also provides access to an easy-to-use API that allows a wide use for other software and applications, e.g., for meta-analysis, design of new clinical studies, or simple literature search. CORDITE is currently empowering many scientists across all continents and accelerates research in the knowledge domains of virology and drug design.

INTRODUCTION

Coronaviruses can cause major outbreaks with fatal pneumonia, e.g., the severe acute respiratory syndrome coronavirus (SARS-CoV) in 2003 with a fatality rate of around 10% (Lee et al., 2003) or the Middle East respiratory syndrome coronavirus (MERS-CoV) in 2012 with a fatality rate of 35% (de Groot et al., 2013). In 2019, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) caused the disease COVID-19 and led to a pandemic (Zhu et al., 2020). The common symptoms of COVID-19 include fever, cough, shortness of breath, and dyspnea. A more severe progression of the COVID-19 disease may lead to pneumonia and severe acute respiratory syndromes. Unfortunately, there are no specific drugs or vaccines approved for the treatment of COVID-19 yet (Huang et al., 2020).

Traditionally, de novo drug design and drug repurposing are the common strategies scientists use to combat viral diseases. However, designing a new drug is very time consuming and needs extensive testing before clinical use in humans. In contrast, the repurposing of approved drugs or drugs in clinical trials is more promising, in particular as these have already been tested for side effects, toxicity, bioavailability, etc. At the moment, there is a huge amount of data from studies that analyze potential drugs based on computer simulations, *in vitro* studies, case studies, or clinical trials. However, none of the proposed treatments for COVID-19 have been proven to be clinically effective yet.

Potential drugs can target viral proteins or human proteins. At the time of writing, the most important viral target molecules that have been investigated for drug repurposing are SARS-CoV-2 spike protein, main protease (3CLpro, Mpro), papain-like proteinase (PLpro), and RNA-dependent RNA polymerase (RdRp) (Ou et al., 2020; Zhang et al., 2020), whereas the main targets on the human cell membrane are angiotensin-converting enzyme 2 (ACE2) and transmembrane protease serine 2 (TMPRSS2) (Aronson and Ferner, 2020). Besides these targets, several other candidates have been investigated as potential targets. Such investigations occurred on both viral and human sides, e.g., the methyltransferase and furin, respectively (Bestle et al., 2020; Khan et al., 2020). Also, several potential drugs were announced at the beginning of the pandemic, which later proved to be insufficiently effective or even harmful to patients, such as hydroxychloroquine or chloroquine phosphate. For instance, Gautret et al., 2020 reported a significant decrease in viral load in patients treated with hydroxychloroquine compared with controls. However, others have shown that these potential drugs can have severe side effects such as cardiac toxicity and also increased

¹Department of Mathematics and Computer Science, Philipps-University of Marburg, Hans-Meerwein-Str. 6, 35032 Marburg, Germany

²These authors contributed equally

³Lead Contact

*Correspondence:
dominik.heider@uni-marburg.de

<https://doi.org/10.1016/j.isci.2020.101297>






Figure 1. Overview of the Replication Cycle of SARS-CoV2 Inside a Human Cell

The main targets that are investigated for drug repurposing are reported.

mortality (Mehra et al., 2020). Nevertheless, some of those studies showing higher mortality were later retracted because of a lack of transparency concerning the data (Mehra et al., 2020b). Other potential drug candidates, such as a combination of lopinavir with ritonavir (also known as Kaletra), were first considered promising candidates (Jeon et al., 2020) but later shown to be ineffective for SARS-CoV-2 (Cao et al., 2020). Nevertheless, many new drug candidates have been proposed by using different computational approaches, which address different viral and human targets (Hufsky et al., 2020; Sadegh et al., 2020).

The SARS-CoV-2 virus uses the human receptor ACE2 for cell entry, which is also used by SARS (Gralinski and Menachery, 2020; Wan et al., 2020; Xu et al., 2020; Zhou et al., 2020). It is expressed as a membrane-bound protein in the lungs, heart, kidneys, stomach, spleen, intestine, bone marrow, kidney, liver, brain, testis, and placenta (Shenoy et al., 2011; Donoghue et al., 2000; Soler et al., 2008). Moreover, Hoffmann et al. (2020) showed that SARS-CoV-2 depends on TMPRSS2, which therefore can serve as a target in therapy as well. TMPRSS2 primes the spike protein of SARS-CoV-2 to enable cell entry on ACE2. The authors further suggested that ACE2 binding to the spike protein might have a higher affinity compared with SARS. This may explain the infection of the upper respiratory system by SARS-CoV-2, which has not been observed in SARS, because in the upper respiratory system less ACE2 is expressed.

Structurally, each virion is approximately 50–200 nm in diameter (Chen et al., 2020). Like other coronaviruses, SARS-CoV-2 has four structural polyproteins, known as the S (spike), E (envelope), M (membrane), and N (nucleocapsid) proteins. The N protein holds the RNA genome, whereas the S, E, and M proteins together are part of the viral envelope. Upon successful attachment of the virus to the membrane of the host cell via the spike protein, the virus fuses with the membrane (Wu et al., 2020; Hoffmann et al.,




Figure 2. Depiction of All Processing Steps within the CORDITE Workflow.

(A) Literature aggregation.

(B) database curation.

(C) web interface.

2020). Next, the endosome sees a drop in pH and the genomic RNAss(+) is released into the cell's cytoplasm (Yang et al., 2004). As shown in Figure 1, upon uncoating and release of the RNA + sense viral genome in the cytoplasm, the host ribosome translates the viral RNA replicase or the viral RNA-dependent RNA polymerase (RdRp) (Ahn et al., 2012; Yin et al., 2020). This triggers proteolysis using the main protease of SARS-CoV-2 (also named 3CLpro) and the papain-like proteinase (PLpro) and enables the replicate-transcriptase complex to transcript subgenomic RNAs(+) (Anand et al., 2003; Chen et al., 2014). Pre-genomic RNAss(–) are replicated into genomic RNAss(+), whereas the former translation of subgenomic RNA(+) results in the aforementioned polyproteins. Essentially, the virus RNA uses the host cell to create new viral RNA genomes and to assemble new viral particles. These are the aforementioned S, E, M, and N structural proteins. They transit to the endoplasmic reticulum and are recruited for the assembly and building of new virions into the Golgi apparatus and burgeon out of the apparatus to then form mature virions (Guo et al., 2020). The N protein allows the viral genome to be extremely tightly packed into future viral capsids. In the end, new virions are released out of the host cell to infiltrate other cells in the host organism (He et al., 2020). A simplified illustration of the replication cycle of the SARS-CoV-2 virus is shown in Figure 1.

The knowledge from the life cycle of the SARS-CoV-2 virus offers a great potential to find suitable drugs for the already known targets. To combine the large amounts of current knowledge, we centralize the aggregated information from published articles and preprints about potential drugs, targets, and their interactions into CORDITE (CORona Drug InTEractions database), which includes a web interface and an application programming interface (API).

RESULTS

CORDITE integrates many functionalities to enable users to access, sort, and download relevant data to conduct meta-analyses, to design new clinical trials, or even to conduct a curated literature search. CORDITE automatically incorporates publications from PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/>), bioRxiv (<https://www.biorxiv.org/>), chemRxiv (<https://www.chemrxiv.org/>), and medRxiv (<https://www.medrxiv.org/>) that report information on computational, *in vitro*, or case studies on potential drugs for COVID-19 (see Figure 2). Besides original research, we also included reviews and comments in the database. The information from the articles and preprints are manually curated by moderators and can be accessed via the web interface or the open API. Users can directly access the publications, interactions,

Type	Numbers
Interactions	849
Targets	25
Drugs	639
Publications	324
Clinical trials	247

Table 1. Summary Table of all Entry Types at Date of Submission

drugs, targets, and clinical trials. The interactions and the effectiveness of certain drugs are also shown as positive/negative results in a transparent way. Thus, the user can easily see the contradictory results of different studies and can take these into account for, e.g., meta-analyses. This access may offer easier integration for future software or apps that may need curated data.

Moreover, we include registered clinical trials from the National Institutes of Health (<https://clinicaltrials.gov/>) for COVID-19. The CORDITE database is updated through manual curation. The current summary for interactions, drugs, etc., is reported in [Table 1](#). An introductory video for the use of CORDITE can be found on the website.

DISCUSSION

CORDITE provides an easy-to-use web interface and also access to an easy-to-use API that allows a wide use for other software and applications, e.g., for meta-analysis, design of new clinical studies, or simple literature search. CORDITE is currently the largest curated database for drug interactions for SARS-CoV-2, and therefore it offers the opportunity to accelerate research in the knowledge domains of virology and drug design. Because of the popularity and impact of the CORDITE website and database, we plan to continue hosting and managing CORDITE throughout the COVID-19 pandemic. Moreover, we will continuously improve the website and database to establish a standard tool for future diseases. We believe our efforts are essential to help researchers and clinicians to get an overview of existing studies and to accelerate the global search for new treatments.

Limitations of the Study

CORDITE aggregates available knowledge on drugs for COVID-19; however, in particular the preprints have not been peer reviewed. Thus, CORDITE cannot guarantee the reliability of the studies and their results, in particular the effectiveness of the drugs.

Resource Availability

Lead Contact

Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Dominik Heider (dominik.heider@uni-marburg.de).

Materials Availability

This study did not generate new unique data.

Data and Code Availability

This study did not generate data. The curated data can be accessed freely via the API.

Availability: CORDITE is available at <https://cordite.mathematik.uni-marburg.de>.

METHODS

All methods can be found in the accompanying [Transparent Methods supplemental file](#).

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.isci.2020.101297>.

ACKNOWLEDGMENTS

The authors thank Hagen Dreßler for technical assistance.

AUTHOR CONTRIBUTIONS

R.M. developed and implemented the database and the API. H.F.L. developed and implemented the web interface and curated approximately 2.3% of the data. M.W. developed the automated literature crawler and curated approximately 11.3% of the data. G.H. created [Figure 1](#) and the visual identity of the web interface and contributed to the initial draft. A.-C.H. contributed to the project design, created [Figure 2](#), and curated approximately 7.4% of the data. D.H. designed and supervised the project, wrote the manuscript, and curated approximately 79.0% of the data. All authors revised the final manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: May 4, 2020

Revised: June 9, 2020

Accepted: June 15, 2020

Published: July 24, 2020

REFERENCES

- Ahn, D.G., Choi, J.K., Taylor, D.R., and Oh, J.W. (2012). Biochemical characterization of a recombinant SARS coronavirus nsp12 RNA-dependent RNA polymerase capable of copying viral RNA templates. *Arch. Virol.* 157, 2095–2104.
- Anand, K., Ziebuhr, J., Wadhwani, P., Mesters, J.R., and Hilgenfeld, R. (2003). Coronavirus main proteinase (3CLpro) structure: basis for design of anti-SARS drugs. *Science* 300, 1763–1767.
- Aronson, J.K., and Fenner, R.E. (2020). Drugs and the renin-angiotensin system in covid-19. *BMJ* 369, m1313.
- Bestle, D., Heindl, M.R., Limburg, H., Lam, van T.V., Pilgram, O., Moulton, H., Stein, D.A., Hardes, K., Eickmann, M., Dolnik, O., et al. (2020). TMPRSS2 and furin are both essential for proteolytic activation and spread of SARS-CoV-2 in human airway epithelial cells and provide promising drug targets. *BioRxiv*. <https://doi.org/10.1101/2020.04.15.204205>.
- Cao, B., Wang, Y., Wen, D., Liu, W., Wang, J., Fan, G., Ruan, L., Song, B., Cai, Y., Wei, M., et al. (2020). A trial of lopinavir-ritonavir in adults hospitalized with severe covid-19. *N. Engl. J. Med.* 382, 1787–1799.
- Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., Qiu, Y., Wang, J., Liu, Y., Wei, Y., et al. (2020). Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 395, 507–513.
- Chen, X., Yang, X., Zheng, Y., Yang, Y., Xing, Y., and Chen, Z. (2014). SARS coronavirus papain-like protease inhibits the type I interferon signaling pathway through interaction with the STING-TAK1 complex. *Protein Cell* 5, 369–381.
- de Groot, R.J., Baker, S.C., Baric, R.S., Brown, C.S., Drosten, C., Enjuanes, L., Fouchier, R.A., Galiano, M., Gorbalevna, A.E., Memish, Z.A., et al. (2013). Middle East respiratory syndrome coronavirus (MERS-CoV): announcement of the coronavirus study group. *J. Virol.* 87, 7790–7792.
- Donoghue, M., Hsieh, F., Baronas, E., Godbout, K., Gosselin, M., Stagliano, N., Donovan, M., Woolf, B., Robison, K., Jeyaseelan, R., et al. (2020). A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circ. Res.* 87, E1–E9.
- Gautret, P., Lagier, J.C., Parola, P., Hoang, V.T., Meddeb, L., Mailhe, M., Doudier, B., Courjon, J., Giordanengo, V., Vieira, V.E., et al. (2020). Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int. J. Antimicrob. Agents*. <https://pubmed.ncbi.nlm.nih.gov/32205204/>.
- Gralinski, L.E., and Menachery, V.D. (2020). Return of the coronavirus: 2019-nCoV. *Viruses* 12, E135.
- Guo, Y.R., Cao, Q.D., Hong, Z.S., Tan, Y.Y., Chen, S.D., Jin, H.J., Tan, K.S., Wang, D.Y., and Yan, Y. (2020). The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak—an update on the status. *Mil. Med. Res.* 7, 11.
- He, F., Deng, Y., and Li, W. (2020). Coronavirus disease 2019: what we know? *J. Med. Virol.* 92, 719–725.
- Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.H., Nitsche, A., et al. (2020). SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 181, 271–280.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., et al. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395, 497–506.
- Hufsky, F., Lamkiewicz, K., Almeida, A., Aouacheria, A., Arighi, C., Bateman, A., Baumbach, J., Beerenwinkel, N., Brandt, C., Cacciabue, M., et al. (2020). Computational strategies to combat COVID-19: useful tools to accelerate SARS-CoV-2 and coronavirus research. *Preprint*. <https://doi.org/10.20944/preprints202005.0376.v1>.
- Jeon, S., Ko, M., Lee, J., Choi, I., Byun, S.Y., Park, S., Shum, D., and Kim, S. (2020). Identification of antiviral drug candidates against SARS-CoV-2 from FDA-approved drugs. *Antimicrob. Agents Chemother.* 64, e00819-20.
- Khan, R.J., Jha, R.K., Amera, G.M., Jain, M., Singh, E., Pathak, A., Singh, R.P., Muthukumaran, J., and Singh, A.K. (2020). Targeting SARS-CoV-2: a systematic drug repurposing approach to identify promising inhibitors against 3C-like proteinase and 2'-O-ribose methyltransferase. *J. Biomol. Struct. Dyn.* 20, 1–14.
- Lee, N., Hui, D., Wu, A., Chan, P., Cameron, P., Joyn, G.M., Ahuja, A., Yung, M.Y., Leung, C.B., To, K.F., et al. (2003). A major outbreak of severe acute respiratory syndrome in Hong Kong. *N. Engl. J. Med.* 348, 1986–1994.
- Mehra, M.R., Desai, S.S., Ruschitzka, F., and Patel, A.N. (2020a). Hydroxychloroquine or chloroquine with or without a macrolide for treatment of COVID-19: a multinational registry analysis. *Lancet*. [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)31180-6/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31180-6/fulltext).
- Mehra, M.R., Ruschitzka, F., and Patel, A.N. (2020b). Retraction—hydroxychloroquine or

chloroquine with or without a macrolide for treatment of COVID-19: a multinational registry analysis. *Lancet.* [https://doi.org/10.1016/S0140-6736\(20\)31324-6](https://doi.org/10.1016/S0140-6736(20)31324-6).

Ou, X., Liu, Y., Lei, X., Li, P., Mi, D., Ren, L., Guo, L., Guo, R., Chen, T., Hu, J., et al. (2020). Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nat. Commun.* 11, 1620.

Sadegh, S., Matschinske, J., Blumenthal, D.B., Galindez, G., Kacprowski, T., List, M., Nasirigerdeh, R., Oubounyt, M., Pichlmair, A., Rose, T.D., and Salgado-Albarrán, M. (2020). Exploring the SARS-CoV-2 virus-host-drug interactome for drug repurposing. *arXiv.* <https://arxiv.org/abs/2004.12420>.

Shenoy, V., Qi, Y., Katovich, M.J., and Raizada, M.K. (2011). ACE2, a promising therapeutic target for pulmonary hypertension. *Curr. Opin. Pharmacol.* 11, 150–155.

Soler, M.J., Barrios, C., Oliva, R., and Batlle, D. (2008). Pharmacologic modulation of ACE2 expression. *Curr. Hypertens. Rep.* 10, 410–414.

Wan, Y., Shang, J., Graham, R., Baric, R.S., and Li, F. (2020). Receptor recognition by the novel coronavirus from wuhan: an analysis based on decade-long structural studies of SARS coronavirus. *J. Virol.* 94, e00127–20.

Wu, C., Liu, Y., Yang, Y., Zhang, P., Zhong, W., Wang, Y., Wang, Q., Xu, Y., Li, M., Li, X., et al. (2020). Analysis of therapeutic targets for SARS-CoV-2 and discovery of potential drugs by computational methods. *Acta Pharm. Sin B* 10, 766–788.

Xu, X., Chen, P., Wang, J., Feng, J., Zhou, H., Li, X., Zhong, W., and Hao, P. (2020). Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci. China Life Sci.* 63, 457–460.

Yang, Z.Y., Huang, Y., Ganesh, L., Leung, K., Kong, W.P., Schwartz, O., Subbarao, K., and Nabel, G.J. (2004). pH-dependent entry of severe acute respiratory syndrome coronavirus is mediated by the spike glycoprotein and enhanced by dendritic cell transfer through DC-SIGN. *J. Virol.* 78, 5642–5650.

Yin, W., Mao, C., Luan, X., Shen, D.D., Shen, Q., Su, H., Wang, X., Zhou, F., Zhao, W., Gao, M., and Chang, S. (2020). Structural basis for inhibition of the RNA-dependent RNA polymerase from SARS-CoV-2 by remdesivir. *Science* 368, 1499–1504.

Zhang, L., Lin, D., Sun, X., Curth, U., Drosten, C., Sauerhering, L., Becker, S., Rox, K., and Hilgenfeld, R. (2020). Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved α -ketoamide inhibitors. *Science* 368, 409–412.

Zhou, P., Yang, X.L., Wang, X.G., Hu, B., Zhang, L., Zhang, W., Si, H.R., Zhu, Y., Li, B., Huang, C.L., et al. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579, 270–273.

Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., et al. (2020). A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.* 382, 727–733.