

# A Systematic Review of Vancomycin Dosing in Patients with Hematologic Malignancies or Neutropenia

This article was published in the following Dove Press journal:  
*Infection and Drug Resistance*

Na He <sup>1,2</sup>  
Fei Dong<sup>3</sup>  
Wei Liu <sup>1</sup>  
Suodi Zhai<sup>1</sup>

<sup>1</sup>Department of Pharmacy, Peking University Third Hospital, Beijing, People's Republic of China; <sup>2</sup>Department of Pharmacy Administration and Clinical Pharmacy, School of Pharmaceutical Science, Peking University, Beijing, People's Republic of China; <sup>3</sup>Department of Hematology, Peking University Third Hospital, Beijing, People's Republic of China

**Objective:** To provide a comprehensive review of vancomycin dosing in patients with hematologic malignancies or neutropenia.

**Methods:** PubMed, Embase and the Cochrane Library were searched through April 2, 2020. Original studies relevant to vancomycin dosing regimen in adults with hematologic malignancies or neutropenia were included. No restriction was applied in study design and language. A descriptive analysis was performed.

**Results:** Twenty-three studies were included eventually, of which eighteen were case series studies, four were cohort studies and another one was a randomized controlled trial. Five case series studies made a clinical audit of conventional vancomycin dosing in patients with malignancies or neutropenia, showing that the proportion of patients with sub-therapeutic trough levels remained high, ranging from 32% to 88%. Seven case series studies and four cohort studies demonstrated that vancomycin clearance (CL<sub>va</sub>) tended to be higher in patients with hematologic malignancies or neutropenia, whereas volume of distribution (V) seemed to be comparable to the control group. Five studies proposed individualized initial dosing regimen per the pharmacokinetic changes; however, no prospective validation has been conducted in clinical setting. Additionally, four case series studies suggested that the correlation between vancomycin clearance and estimated creatinine clearance was relatively poor, bringing a great challenge to proper dosing strategy. A randomized controlled trial stated that therapeutic drug monitoring (TDM) of vancomycin could decrease the incidence of nephrotoxicity in immunocompromised febrile patients with hematologic malignancies.

**Conclusion:** The available evidence indicates that conventional vancomycin dosing leads to suboptimal concentration in patients with hematologic malignancy or neutropenia. TDM accompanied by pharmacokinetic interpretation can decrease the risk of nephrotoxicity. The individualization of the initial dosing regimen and mechanisms of augmented clearance require further research.

**Keywords:** vancomycin, hematologic malignancy, neutropenia, pharmacokinetics, evidence-based practice

## Introduction

A proper dosing regimen is the cornerstone of antimicrobial therapy, which has a great impact on treatment outcome, development of drug resistance as well as dose-dependent toxicity. Traditionally, the use of reduced doses in patients with renal impairment has been widely accepted. Dosage adjustments for patients with renal failure have been listed in labels of various medications and relevant clinical guidelines.<sup>1,2</sup> However, more and more studies have underlined the

Correspondence: Suodi Zhai; Wei Liu  
Department of Pharmacy, Peking University Third Hospital, 49 Huayuan North Road, Haidian District, Beijing 100191, People's Republic of China  
Tel +86 10-82266686; +86 10-82265740  
Email zhaisuodi@163.com; andthen0023@163.com

existence of augmented renal clearance (ARC), especially in critically ill patients,<sup>3</sup> patients with brain injury<sup>4</sup> and neurosurgery,<sup>5</sup> which could result in antibiotics' sub-therapeutic concentrations and poorer outcomes. In this case, an assumption could be made that dosing regimens should be optimized according to the degree of increase in renal function, similar to the downward dose adjustments in patients with renal dysfunction.

Risk of infections will increase in patients with neutropenia, which occurs frequently after chemotherapy for cancer, especially hematologic malignancies.<sup>6</sup> Therefore, the administration of optimal antibiotics was recommended in clinically or microbiologically documented infections.<sup>7</sup> Additionally, patients with hematologic malignancies or neutropenia have been reported to have enhanced renal clearance,<sup>8,9</sup> which would affect the systematic exposure of antibiotics predominantly excreted through urine, including the commonly used anti-pseudomonas beta-lactams, aminoglycosides and vancomycin. Hence, the optimization of dosing regimens' might also be required under the circumstance.

To our knowledge, vancomycin is one of the most well-studied antibiotics with respect to therapeutic drug monitoring (TDM).<sup>2,10,11</sup> In spite of the potential changes in pharmacokinetic parameters and possible clinical failure proposed in patients with hematologic malignancies or neutropenia,<sup>8,9</sup> neither increased dosing regimen nor TDM of vancomycin has been recommended in these patients, implying that the evidence was insufficient or the integration of evidence into practice should be strengthened. Notably, no comprehensive review has been conducted on this issue.

The objective of this study was to gain an in-depth understanding of the current status of vancomycin dosing regimen, pharmacokinetics and optimization of vancomycin dosing in patients with hematologic malignancies or neutropenia, which could be of great value for clinical practice and identifying knowledge gaps for future research.

## Methods

We conducted this systematic review using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.<sup>12</sup>

## Data Sources and Searches

PubMed, Embase and the Cochrane Library were searched from their respective inception to July 26th., 2018. A complementary search was also performed to identify the most recent articles (published before April 2, 2020). The search terms included hematologic malignancy, neutropenia and vancomycin. Both mesh terms and text words were used. The search strategy is detailed in [Tables S1–S3](#). Reference lists of the retrieved articles and related reviews were also examined manually for additional studies.

## Eligibility Criteria

All records that comprised of adult patients with hematologic malignancies or neutropenia were included. When the proportion of hematologic malignancies or neutropenia was greater than 80% in one individual arm, the arm could be assumed to be patients with hematologic malignancies or neutropenia, respectively. Furthermore, all the patients were required to receive intravenous vancomycin. Outcomes should involve at least one of the followings: vancomycin serum concentration, pharmacokinetic (PK) parameters, vancomycin dosing, clinical response and nephrotoxicity. The exclusion criteria were as follows: (1) insufficient clinical data; (2) study types were cases, reviews or editorials; (3) the analysis was not relevant to vancomycin dosing regimen; (4) duplicate publication. No restriction was applied in language.

## Study Selection

Two reviewers (N. H. and W. L.) screened titles and abstracts per the eligibility criteria to identify potential publications independently at first. Then, the full text was assessed for final inclusion. Any disagreement was resolved by discussion between the 2 reviewers or by consulting a third reviewer (S. Z.).

## Data Extraction

A pre-specified data form was used to extract the following information: study characteristics (the first author's name, year of publication, study design, country, sample size), patients' baseline characteristics (characteristics of patients included, proportion of patients with neutropenia, gender, age, weight, renal function), vancomycin dosing, timing of vancomycin serum concentration sampling, outcomes of interest. The data

extraction was performed by one reviewer (N. H.) and checked by another reviewer (W. L.). Discrepancies were addressed by discussion between two reviewers or consultation with the third reviewer (S. Z.) if necessary.

## Quality Assessment

The methodological quality of each included study was assessed by 2 reviewers (N. H. and X. L.) independently, and disagreements were resolved by discussion. The potential risk of bias in the randomized controlled trials was assessed using Cochrane risk of bias.<sup>13</sup> The quality of cohort studies was assessed per the Newcastle-Ottawa Scale (NOS) scale.<sup>14</sup> Concerning case series studies, we used National Institutes of Health (NIH) Quality Assessment Tool (<https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>). As no validated tool for pharmacokinetic studies was available, we used the ClinPK Statement, a reporting guideline for clinical pharmacokinetic studies to assess their quality.<sup>15</sup>

## Data Analysis


To summarize all the information concerning vancomycin dosing in patients with hematologic

malignancies or neutropenia, a descriptive analysis was performed.

## Results

Of the 6404 potentially relevant published reports identified, 46 reports proved potentially eligible after duplicates removed and abstracts screened. On full-text screening, 23 studies were ultimately included in the systematic review (Figure 1). The list of the excluded studies in the process of full-text screening is detailed in Table S4.

The basic characteristics of the included studies can be found in Table S5. All the 23 studies<sup>16–38</sup> were published in English, of which three<sup>16,18,19</sup> were conference abstracts. One study<sup>33</sup> was a simulation study without an actual clinical data. Two studies<sup>20,32</sup> adopted the same set of data with different analyses methods. Therefore, twenty-one sets of clinical data were finally included. In 18 studies,<sup>17,18,20,22–31,34–38</sup> vancomycin was infused intermittently, and 2 studies adopted continuous infusion,<sup>19,21</sup> whereas the remaining 1 conference abstract did not report the specific dosing regimen.<sup>16</sup> For single-arm studies, nine studies included neutropenic hematologic patients consecutively,<sup>16,18,20,21,26,28,31,34,35</sup> while 6 studies included patients with hematologic



**Figure 1** Flow chart of study selection.

malignancies without distinguishing neutropenia from non-neutropenia.<sup>17,19,22,27,29,38</sup> Additionally, another two studies included hospitalized patients and took neutropenia as a risk factor.<sup>30,36</sup> Concerning comparative cohort studies, there were two studies comparing neutropenic patients with non-neutropenic patients,<sup>23,25</sup> and another 2 studies focused on the difference between hematologic malignant patients and control groups.<sup>24,37</sup>

Concerning the quality assessment of these included studies, 4 studies were not assessed, of which 3<sup>16,18,19</sup> were conference abstracts and one<sup>33</sup> was a simulation pharmacokinetic study. The detailed results of quality assessment are shown in [Tables S6–S9](#). Overall, included studies were of adequate quality.

All the included studies were classified according to their objectives as follows:

## Clinical Audit of Vancomycin Dosing

Seven studies<sup>16–22</sup> aimed to make a clinical audit of conventional vancomycin dosing in patients with hematologic malignancies or neutropenia. The characteristics and summary of results in each study are listed in [Table 1](#). Six studies reported the proportion of patients with sub-therapeutic concentrations in routine clinical care, and five of which reported value ranging from 32% to 88%. However, Vazin et al<sup>22</sup> did not report the specific vancomycin dosing, yielding result (3.6%) that differed significantly from other 5 studies. Furthermore, Vermis et al<sup>19</sup> stated that to attain therapeutic vancomycin levels, vancomycin maintenance dose (41.7 mg/kg/d vs. 32.7 mg/kg/d) was significantly higher when ARC (estimated  $CL_{CR}$  greater than 120 mL/min) was present in hematologic malignant patients. Overall, vancomycin concentrations following conventional dosage were insufficient in patients with hematologic malignancies or neutropenia.

## The Potential Change in Pharmacokinetic Parameters

### Comparative Studies Between Patients with Hematologic Malignancies or Neutropenia and Control Groups

Two studies<sup>23,25</sup> were comparative cohort studies between neutropenic patients and non-neutropenic patients, and another 2 studies<sup>24,37</sup> focused on the difference between patients with hematologic malignancies and the control group. The characteristics of the four studies are summarized in [Table 2](#).

Although both neutropenic patients and the control group applied the standard dosage and consistent sampling time in Choi et al,<sup>23</sup> the median serum vancomycin concentration was lower in neutropenic patients than the control group (9.1 mg/L vs. 12.1 mg/L,  $P < 0.0001$ ). Multiple logistic regression analysis still revealed a significant association between sub-therapeutic vancomycin concentration (trough serum concentration  $< 10$  mg/L) and neutropenia (odds ratio [OR]: 1.75;  $P = 0.029$ ).

Additionally, Haeseker et al<sup>25</sup> primarily investigated neutropenia and hematologic malignancy's effect on pharmacokinetic parameters, whereas Al-Kofide et al<sup>24</sup> and Izumisawa et al<sup>37</sup> focused on the effect of hematologic malignancy. Concerning specific pharmacokinetic parameters, vancomycin clearance ( $CL_v$ ) was higher in patients with hematologic malignancies or neutropenia ([Table 3](#)). However, the results for volume of distribution ( $V$ ) were still conflicting ([Table 3](#)). Notably, five patients in Haeseker et al<sup>25</sup> received vancomycin in both neutropenic and nonneutropenic period and presented a reversible augmented  $CL_v$  in the nonneutropenic period ( $91 \pm 26$  mL/min vs.  $45 \pm 10$  mL/min,  $P = 0.009$ ).

### Development and Validation of PK Models

Although seven studies<sup>20,26–29,36,38</sup> calculated vancomycin's pharmacokinetic parameters in patients with hematologic malignancies or neutropenia, only three studies<sup>29,36,38</sup> used non-linear mixed effects modelling. The characteristics and PK parameter of studies included are listed in [Table 4](#), showing a marked difference in  $CL_v$  from those reported for patients with non-hematologic malignancy and non-neutropenia.<sup>39,40</sup> Notably, one study<sup>36</sup> included neutropenia as one of the covariates affecting vancomycin clearance, of which vancomycin clearance is increased in patients with neutropenia by 27.7%. Nevertheless,  $V$  seemed to be comparable to normal controls without hematologic malignancy and neutropenia.<sup>39,40</sup> Additionally, all the PK parameters had great inter-individual variation among patients with hematologic malignancies or neutropenia.

## The Potential Effect of Neutropenia on Creatinine Clearance ( $CL_{CR}$ )

Four studies<sup>25,28,30,31</sup> reported the potential effect of neutropenia on  $CL_{CR}$ . Hirai et al<sup>30</sup> conducted a single-center retrospective study in 292 patients with normal serum

creatinine concentration, and demonstrated that febrile neutropenia was an independent risk factor of ARC (OR: 2.76; 95% CI: 1.11–6.67;  $P = 0.0254$ ). However, Haeseke et al<sup>25</sup> showed that the estimated  $CL_{CR}$  was not significantly different between patients with neutropenia and non-neutropenia (Table 2).

Three studies evaluated the correlation between  $CL_{va}$  and estimated  $CL_{CR}$  solely. Soto et al<sup>31</sup> included 45 neutropenic ( $<1000/\text{mm}^3$ ) hematologic patients and demonstrated that the correlation coefficient between  $CL_{va}$  ( $106 \pm 37$  mL/min) and estimated  $CL_{CR}$  ( $84.7 \pm 32$  mL/min) was 0.42. Le Normand et al<sup>28</sup> illustrated a poor correlation in neutropenic patients ( $100/\text{mm}^3$ ) as well ( $n = 10$ ,  $r = 0.281$ ). According to Hirai et al,<sup>30</sup> the non-ARC patients showed a significant correlation between  $CL_{CR}$  and  $CL_{va}$  ( $r = 0.8726$ ,  $P < 0.0001$ ); however, no such relationship was observed in patients with ARC ( $r = 0.1029$ ,  $P = 0.4866$ ).

Above all, although  $CL_{CR}$  possibly has an increase in patients with neutropenia, estimated  $CL_{CR}$  itself could not identify the specific patients with ARC, which brought difficulty to the prediction of  $CL_{va}$ .

## Optimization of Initial Vancomycin Dosing Regimen

Six studies<sup>24,25,32,33,36,38</sup> were relevant to the optimization of initial vancomycin dosing regimen. Taghizadeh-Ghehi et al<sup>32</sup> evaluated the applicability of the most cited vancomycin one-compartment models developed in common patients using data from their recent study.<sup>20</sup> They demonstrated that none of the seven pharmacokinetic models performed well to calculate initial vancomycin dosage in Iranian patients underwent hematopoietic stem cell transplantation. Using a published population pharmacokinetic (PPK) model<sup>29</sup> in patients with hematologic malignancies, Fernandez et al<sup>33</sup> performed Monte Carlo simulation to calculate vancomycin dosages required in the specific subpopulation. When standard vancomycin dosing (2000 mg/d) was given, cumulative fraction of response (CFR) for *S. aureus* was 90.4%, 47.3% and 31.2% for  $CL_{CR}$  values of  $<60$ , 60–120 and  $>120$  mL/min, respectively. If a CFR of 80% was considered to be clinically appropriate, vancomycin doses of 3000 and 4000 mg/d for a  $CL_{CR}$  60–120 and  $>120$  mL/min should be used. Okada et al<sup>38</sup> also proposed a vancomycin dosing nomogram in patients undergoing

allogeneic hematopoietic stem-cell transplantation based on PPK model and Monte Carlo simulation. Suggested vancomycin dosing is 1g per 12 hours when  $CL_{CR}$  ranging from 75 to 90 mL/min, 0.75 g per 8 hours when  $CL_{CR}$  ranging from 90 to 120 mL/min, 1g per 8 hours when  $CL_{CR}$  ranging from 120 to 175 mL/min, and 1.25 g per 8 hours for  $CL_{CR}$  greater than 175 mL/min. Based on individualized pharmacokinetic parameters calculated by Al-Kofide et al,<sup>24</sup> the actual dosing regimen for cancer patients should be 60 mg/kg/day, which doubled the required dose for the general population (30 mg/kg/d). Haeseke et al<sup>25</sup> demonstrated that to achieve the same  $AUC_{24}$ , the mean dosage in patients with neutropenia was significantly higher than the control group ( $2017 \pm 720$  vs  $1521 \pm 727$  mg,  $P < 0.001$ ). In this case, they concluded that the daily dose should be increased with 33% in patients with neutropenia (from 15 mg/kg twice daily to 13 mg/kg three times daily). Similarly, another study<sup>36</sup> suggested a 25% increase for vancomycin dosing in neutropenic patients. However, the dosing algorithms aforementioned were inconsistent to some extent and have not been validated in the prospective clinical setting. Hence, no simple upward dose adjustment can be put up with great validity and the initial dosing recommendation still remains investigational.

## Evaluation and Implementation of Vancomycin TDM

Two aspects of TDM have been explored before, including the target trough concentration and the evaluation of TDM-guided vancomycin therapy. Suzuki et al<sup>35</sup> retrospectively included 63 febrile neutropenic patients with hematologic malignancies and investigated the association of first trough concentration at steady state with clinical efficacy and nephrotoxicity. They proposed that the cut-off value of vancomycin trough concentration should be around 11.5 mg/L in these patients.

To assess the effectiveness and safety of vancomycin TDM and pharmacokinetic interpretation, Fernandez et al<sup>34</sup> performed a prospective randomized study in 70 immunocompromised febrile patients with hematologic malignancies. Although there was no significant difference in clinical response rate and duration of fever between TDM-guided group ( $n=37$ ) and control group ( $n=33$ ), the incidence of nephrotoxicity significantly decreased (13.5% vs. 42.4%,  $P < 0.05$ ).

**Table 1** Studies with Clinical Audit of Vancomycin Dosing

Author (Year)	Country	Study Design	Characteristics of Patients Included	Patients with Neutropenia (%)	Sample Size	Age (Years)	Gender (M/F)
Hochart 2011 <sup>21</sup>	France	Single-center retrospective study	Acute myeloid leukemia patients with febrile neutropenia	100%	54 (67 vancomycin treatment courses, VTCs)	50 ± 13.6	27/27
O'Donnell 2011 <sup>16</sup> (conference abstract)	UK	Single-center retrospective study	Bone marrow transplant patients who experienced an episode of febrile neutropenia	100%	12	NR	NR
Donovan 2012 <sup>18</sup> (conference abstract)	United states	Retrospective study	Neutropenic adult patients	100%	198	NR	NR
Vazin 2012 <sup>22</sup>	Iran	Prospective study	Patients in a hematology-oncology ward who received at least 3 successive doses of vancomycin and had serum vancomycin concentrations at steady state	88%	58	36.58 ± 14.33	44/14
Ghehi 2013 <sup>20</sup>	Iran	Single-center prospective study	Adults receiving vancomycin for neutropenic fever after HSCT	100%	46	32.9 ± 12.45	30/16
Luo 2014 <sup>17</sup>	Canada	Single-center prospective study	Leukemia/bone marrow transplant outpatients (at least two doses of vancomycin)	42%	48	54.5 <sup>b</sup>	24/24
Vermis 2014 <sup>19</sup> (conference abstract)	Belgium	Single-center retrospective study	Patients with hematologic malignancies	72%	96 (112 VTCs)	NR	NR

**Notes:** <sup>a</sup>Proportion of patients with normal renal function; <sup>b</sup>median; <sup>c</sup>ARC was defined as calculated creatinine clearance exceeding 120 mL/min (Cockcroft–Gault formula).  
**Abbreviations:** NR, not reported; HSCT, hematopoietic stem cell transplantation; VTC, vancomycin treatment course; ARC, augmented renal clearance.

## Discussion

### Brief Summary of the Systematic Review

Several descriptive studies demonstrated that their routine vancomycin dosing was inadequate for effective antimicrobial therapy. Regarding the alterations in PK parameters, studies showed that CL<sub>va</sub> tended to be higher in patients with hematologic malignancies or

during febrile neutropenia, whereas V seemed to be comparable to the control groups. Although several pharmacokinetic models have been developed and a few dosing regimens have been proposed, there is still no consensus on initial vancomycin dosing in patients with hematologic malignancies or neutropenia. The available evidence indicates that TDM and optimal

Weight (kg)	Renal Function	Vancomycin Dosing Regimen	Timing of Serum Vancomycin	Summary of Results
73 ± 18.1	107.5 ± 35.4 mL/min	Continuous infusion; loading dose: 15.5 ± 3.3 mg/kg; maintenance dose: 35.4 ± 6.9 mg/kg/d	At 24 hours for patients with a loading dose and 48 hours for patients without any loading dose.	<ul style="list-style-type: none"> <li>the target serum level for continuous infusion was greater than 20 mg/L, and only 6 (12%) cases achieved the target</li> </ul>
NR	NR	NR	At 24 hours, then twice weekly.	<ul style="list-style-type: none"> <li>32% of trough levels were subtherapeutic (&lt; 5 mg/L).</li> </ul>
NR	NR	Intermittent infusion; 15 – 20 mg/kg/dose and administration times are determined by renal function	NR	<ul style="list-style-type: none"> <li>25.3% of patients achieved therapeutic trough concentrations (15 – 20 mg/L)</li> </ul>
68.05 ± 12.61	57/58 (98.2%) <sup>a</sup>	Intermittent infusion; Correcting dosage based on creatinine clearance was given to 10 (17.23%) of the patients	Blood samples were taken from the patients who received vancomycin for 3 consecutive days, and just before the administration of the next dose.	<ul style="list-style-type: none"> <li>vancomycin trough serum concentration range was 15.59 ± 13.02 mg/L</li> <li>subtherapeutic trough level (&lt; 10 mg/L) was detected in 3.6% of patients</li> <li>53.3% had a level above the maximum therapeutic concentration</li> </ul>
74.8 ± 16.6	102.5 ± 35.33 mL/min	Intermittent infusion; 31.9 (±10.5) mg/kg/d	Within 30 minutes prior to the fourth dose	<ul style="list-style-type: none"> <li>25 (54.3%) patients had trough concentrations of &lt;10 mg/L</li> <li>6 patients (13%) had trough levels of &lt; 5 mg/L</li> </ul>
NR	77 μmol/L <sup>b</sup>	Intermittent infusion; once-daily (2073 ± 338 mg/d)	NR	<ul style="list-style-type: none"> <li>10 (21%) patients had therapeutic vancomycin trough concentrations (i.e., greater than 10 mg/L)</li> </ul>
NR	NR	Continuous infusion; loading dose: 15 mg/kg, maintenance dose: 30 mg/kg/d	NR	<ul style="list-style-type: none"> <li>ARC<sup>c</sup> was observed in 73 VTC with an average renal clearance of 147.0 mL/min versus 79.0 mL/min.</li> <li>Therapeutic vancomycin levels (20 mg/L) were obtained on day 5 (median) with an average vancomycin maintenance dose of 41.7 mg/kg/day when ARC was present versus 32.7 mg/kg/day on day 3.</li> </ul>

pharmacokinetic interpretation can help in decreasing the risk of nephrotoxicity.

## Implications for Clinical Practice

In view that standard dosing is inadequate for some patients with hematologic malignancies or neutropenia, improper dosing should be considered as a possible rea-

son when clinical improvement was not achieved in these patients with suspected or documented Gram-positive infection. Therefore, optimization of dosing regimen must be considered in both initial dosing and dose adjustment. However, it still remains a question of how to identify the patients with ARC accurately, which makes the individualization of initial dosing difficultly. For

**Table 2** The Characteristic of Comparative Studies

Author (Year)	Country	Study Design	Characteristics of Patients Included	Sample Size	Grouping		Gender (M/F)	Age (Years)	
					Study Group	Control Group		Study Group	Control Group
<b>Neutropenic patients vs non-neutropenic patients</b>									
Haeseker 2014 <sup>25</sup>	Netherlands	Single-center prospective study	Adults received vancomycin intravenously and had at least two plasma samples	171	Neutropenia (< 500/mm <sup>3</sup> ): n=56	Non-neutropenia: n=115	104/67	55 ± 13	61 ± 14
				68 (a subset of patients with hematologic malignancies)	Neutropenia (< 500/mm <sup>3</sup> ): n=55	Non-neutropenia: n=13	NR	NR	NR
Choi 2017 <sup>23</sup>	Korea	Single-center retrospective study	Adults receiving routine TDM of vancomycin (trough and peak).	1307	Neutropenia (< 500/mm <sup>3</sup> ) n=162	Non-neutropenia: n=1145	728/579	54 (37–65) <sup>#</sup>	56 (45–64) <sup>#</sup>
<b>Patients with hematologic malignancies vs non-cancer patients</b>									
Al-Kofide 2009 <sup>24</sup>	Saudi Arabia	Single-center retrospective study	Adults receiving vancomycin therapy	31	Cancer patients (proportion of patients with hematologic malignancies was 88.9%): n=18	Patients without cancer: n=13	NR	48.5 ± 20.2	43.4 ± 22.1
Izumisawa 2019 <sup>37</sup>	Japan	Retrospective cohort study	Adults receiving > 3 days of vancomycin therapy	522	Hematologic malignancy patients: n=261	Non-malignancy patients: n=261	321/201	65.6 ± 13.6	67.2 ± 16.9

**Notes:** <sup>#</sup>Median (interquartile range); <sup>a</sup>proportion of patients with hematologic malignancy; <sup>b</sup>proportion of patients with neutropenia; <sup>c</sup>the absolute count of neutrophils.

**Abbreviations:** NR, not reported; V, volume of distribution; V<sub>ss</sub>, volume of distribution at steady state; CL<sub>va</sub>, vancomycin clearance; t<sub>1/2</sub>, half-life; TDM, therapeutic drug monitoring.

example, Soto et al<sup>31</sup> and Le Normand et al<sup>28</sup> demonstrated that the correlation between estimated CL<sub>CR</sub> and CL<sub>va</sub> was poor. Haeseker et al<sup>41</sup> also showed that CL<sub>va</sub> algorithms based on estimated CL<sub>CR</sub> were unsuitable in these patients. Additionally, Taghizadeh-Ghehi et al<sup>32</sup> demonstrated that none of the seven most cited vancomycin one-compartment models performed well to calculate initial dosage. In this case, TDM of vancomycin could be valuable in patients with hematologic malignan-

cies or neutropenia. Without performing TDM, the extremely high dosing could not be administered. Furthermore, previous studies demonstrated that pharmacokinetic dosing programs using measured vancomycin serum levels could predict vancomycin levels with acceptable accuracy and precision.<sup>42</sup> Therefore, we recommend, when possible, TDM-guided therapy to optimize vancomycin therapy in patients with hematologic malignancy or neutropenia. Above all, the systematic review



Weight (kg)		Patients with Hematologic Malignancies/ Neutropenia n(%)		Renal Function		Vancomycin Dosing Regimen	Timing of Serum Vancomycin	Determination of Pharmacokinetic Parameters	Outcomes
Study Group	Control Group	Study Group	Control Group	Study Group	Control Group				
NR	NR	55/56 (98.2%) <sup>a</sup>	13/115 (11.3%) <sup>a</sup>	113 ± 57 mL/min	107 ± 78 mL/min	Intermittent infusion; an initial loading dose of 15 mg/kg + dose individualization based on TDM and renal function.	Two plasma samples (peak and trough concentration)	Maximum a posteriori (MAP) Bayesian estimation (MW/Pharm 3.60, Mediware, the Netherlands)	CL <sub>Va</sub> , V
NR	NR	55 (100%) <sup>a</sup>	13 (100%) <sup>a</sup>	114 ± 57 mL/min	111 ± 58 mL/min				
62.0 (56.0–70.0) <sup>#</sup>	60.0 (53.0–68.7) <sup>#</sup>	135 (83.3%) <sup>a</sup>	184 (16.1%) <sup>a</sup>	0.6 (0.5–0.8) <sup>#</sup> mg/dL	0.7 (0.5–0.9) <sup>#</sup> mg/dL	Intermittent infusion; 1000 mg vancomycin every 12 h	Steady-state serum vancomycin concentration (after at least the fourth dose)	Posterior Bayesian estimation (Abbott's PKS software)	Serum trough vancomycin concentration at steady state, $t_{1/2}$
66.7 ± 17.1	68.9 ± 14	NR <sup>b</sup>	NR <sup>b</sup>	105.4 ± 62.3 mL/min	87.2 ± 27.5 mL/min	Intermittent infusion; Initial vancomycin dosing regimens were chosen by attending physicians	Peak and trough vancomycin serum concentration (after the third dose or at steady state)	Pharmacokinetic equations	CL <sub>Va</sub> , V, $t_{1/2}$
55.0 ± 10.3	56.2 ± 13.1	1.47 ± 2.46 × 10 <sup>3</sup> /μL <sup>c</sup>	7.80 ± 4.66 × 10 <sup>3</sup> /μL <sup>c</sup>	77.0 ± 29.2 mL/min	74.1 ± 35.6 mL/min	Intermittent infusion; Initial dosing was not pre-specified	After ≥ 3 days following the start of administration	Bayesian estimation using TDM software Ver 3.3	Trough concentration, CL <sub>Va</sub> , V <sub>ss</sub> , $t_{1/2}$

underlines the necessity to perform vancomycin TDM in patients with hematologic malignancies or neutropenia, which might be overlooked previously.

## Implications for Further Research

According to the comprehensive systematic review, several knowledge gaps have been identified, and are summarized as follows:

- The mechanism of the altered PK parameters warrants investigation, which could help us judge whether the phenomenon was deceptive or not.

Two scenarios should be considered to clarify the mechanism of the altered PK parameters. On the one hand is the further research in clinical settings. First, most of the studies did not distinguish whether the change in pharmacokinetic parameters was due to

**Table 3** The Pharmacokinetic Parameters in Comparative Studies

Author (Year)	Sample Size	Vancomycin Clearance (mL/min)			T <sub>1/2</sub> of Vancomycin (h)			V (L)		
		Hematologic Malignancies or Neutropenia	Control Group	P value	Hematologic Malignancies or Neutropenia	Control Group	P value	Hematologic Malignancies or Neutropenia	Control Group	P value
<b>Neutropenic patients vs. non-neutropenic patients</b>										
Haeseker 2014 <sup>25</sup>	171 68 (subset of patients with hematologic malignancies)	67 ± 26 68 ± 26	50 ± 22 53 ± 16	<0.001 0.024	- -	- -	- -	62 ± 32 62 ± 32	56 ± 29 59 ± 18	0.304 0.691
Choi 2017 <sup>23</sup>	1307	-	-	-	7.4 h (5.9–10.7 h) #	8.9 h (6.9–12.0 h) #	-	-	-	-
<b>Patients with hematologic malignancies vs. non-cancer patients</b>										
Al-Kofide 2009 <sup>24</sup>	31	110.1 ± 42	71.2 ± 22.2	0.005	8.6 ± 7.1	5.4 ± 1.9	0.111	70 ± 45	31.1 ± 8.3	0.002
Izumisawa 2019 <sup>37</sup>	522	0.055 ± 0.017 L/h/kg	0.051 ± 0.019 L/h/kg	<0.05	32.7 ± 13.0	37.1 ± 18.8	<0.05	1.81 ± 0.57 L/kg	1.84 ± 0.62 L/kg	>0.05

**Notes:** -, not reported; #, median (interquartile range).

hematologic malignancy or neutropenia. The effect of hematologic malignancy can be complicated by neutropenia and vice versa. Only Haeseker et al<sup>25</sup> demonstrated that the augmented clearance was associated with neutropenia rather than hematologic malignancies with a limited sample size. As previous studies showed vancomycin clearance was higher in patients with hematologic malignancies than solid tumors<sup>43</sup> and the difference between solid malignancies and control groups was attenuated,<sup>44</sup> we assume that the phenomenon might be explained by different proportions of patients with neutropenia. In other words, it is the neutropenia that affects pharmacokinetic changes per se. Conventional doses of vancomycin may not offer adequate systematic exposure in febrile neutropenic patients rather than hematologic malignancy without neutropenia. Further studies are needed to elucidate the exact effect between hematologic malignancies and neutropenia. Second, studies focused on non-neutropenic immunocompromised states are limited currently. Whether the pharmacokinetic changes exist in non-neutropenic immunocompromised states can help in interpreting the mechanism. Third, none of the studies evaluated the change in measured  $CL_{CR}$ , which might also be helpful for understanding the mechanism and identifying patients with ARC.

On the other hand, the physiological mechanism responsible for ARC has not been well-defined, which can be investigated using *in vitro* studies and animal models. Several assumptions have been put up, including: (1) possible changes in renal function and urine flow can be induced by cancer, systematic inflammation and increased intravenous fluid; (2) tubular secretion apart from glomerular filtration; (3) non-renal elimination of vancomycin, such as hepatic conjugation; (4) cancer and neutropenia could enhance vascular permeability, which would induce increased vancomycin extravasation and low serum concentrations.<sup>20,23-25,28,29</sup> The above assumptions require further exploration.

- No PPK model has been developed in patients with hematologic malignancies and concomitant neutropenia, which could help with determining individualized initial dose.
- The optimization of vancomycin dosing in patients with hematologic malignancies or

neutropenia requires further research. For example, prospective validation of vancomycin initial dosing regimens, stages of enhanced  $CL_{CR}$  and a consensus of initial dosing strategies are urgently needed worldwide.

- Few studies on the PKs of vancomycin in patients with hematologic malignancies or neutropenia reported clinical outcomes. Taking safety endpoints as an example, some studies illustrated that patients with hematologic malignancies may be vulnerable to nephrotoxicity.<sup>45,46</sup> In this case, the target trough concentration for these patients might be different from other patients and require further research. Indeed, it should be noted that the establishment of the relationship between accelerated vancomycin elimination and outcomes of clinical effectiveness is difficult due to the complexity of these patients.

## Strengths and Limitations

To our knowledge, this is the first comprehensive review concerning vancomycin dosing optimization in patients with hematologic malignancy or neutropenia. Additionally, the characteristics and the key results of each individual study are presented in [Tables 1–4](#), which can provide specific details for physicians, pharmacists as well as researchers. However, as the available evidence to date was limited and diverse, no quantitative analysis was performed. Clinical heterogeneity existed across studies. For example, patients included had different types of hematologic malignancies and no consistent definition of neutropenia has been applied among studies. Nevertheless, we consider that this systematic review maps the relevant literature on this topic, allowing us to pay attention to the optimization of vancomycin dosing in patients with hematologic malignancy or neutropenia.

## Conclusion

The available evidence indicates that conventional vancomycin dosing leads to suboptimal concentration in patients with hematologic malignancy or neutropenia. TDM accompanied by pharmacokinetic interpretation can decrease the risk of nephrotoxicity. The individualization of the initial dosing regimen and mechanisms of augmented clearance require further research.

**Table 4** Characteristics and Results of Studies for Developing PK/PPK Models

Author (Year)	Country	Study Design	Characteristics of Patients Included	Patients with Neutropenia (%)	Sample Size	Number of serum Concentrations	Age	Gender (M/F)	Weight (kg)
<b>Pharmacokinetic analysis</b>									
Kureishi 1990 <sup>26</sup>	Canada	Single-center prospective study	Patients with acute leukemia and had absolute granulocyte below 500/mm <sup>3</sup>	100%	25	NR	NR	NR	NR
Le Normand 1994 <sup>28</sup>	France	Single-center prospective study	Patients with hematologic malignancies who were neutropenic (100/mm <sup>3</sup> )	100%	10	130	36.2 (range: 18–50)	4/6	64.6 ± 10.4
Jarkowski 2011 <sup>27</sup>	United states	Single-center prospective study	Acute myeloid leukemia patients receiving vancomycin	NR	25	NR	59.12 ± 16.26	17/8	86.05 ± 19.42
Ghehi 2013 <sup>20</sup>	Iran	Single-center prospective study	Patients with neutropenic fever after HSCT	100%	20	40	29.9 ± 9.5	NR	72.5 ± 15.2 (ABW)
<b>Population pharmacokinetic model</b>									
Buelga 2005 <sup>29</sup>	Spain	Single-center retrospective study	Adult inpatients with an underlying hematologic malignancy	43.7%	215	1004	51.5 ± 15.9	119/96	64.7 ± 11.3
Okada 2018 <sup>38</sup>	Japan	Single-center retrospective study	Patients undergoing allo-HSCT who received preventive treatment with vancomycin	NR	75	227	49 (range: 17–69)	49/26	59.4 (range: 39.4–104.5)
Bury 2019 <sup>36</sup>	The Netherlands	Retrospective matched cohort study	Intravenous vancomycin therapy for ≥ 2 days and at least one available vancomycin concentration	26.7%	116	742	61.4 ± 13.4	67/49	NR

**Abbreviations:** V, volume of distribution; CL, clearance; Ke, elimination rate constant;  $t_{1/2}$ , half-life; NR, not reported; ABW, adjusted body weight; TBW, total body weight;  $CL_{CR}$ , creatinine clearance; Vc, volume of central compartment; Vss, steady-state volume of distribution; Vp, distribution volume of peripheral compartment.

Renal Function	Vancomycin Dosing	Timing of Vancomycin Sampling	Pharmacokinetic Modeling Method	Model	Pharmacokinetic Parameters			
					V	CL	Ke (h <sup>-1</sup> )	t <sub>1/2</sub> (h)
NR	Intermittent infusion; 15 mg/kg q12h	Prior to infusion and at 1 and 3 h post-infusion daily during the first 3 days and every 3 to 7 days thereafter	Equations with two steady-state samples	One-compartment model	0.61 ± 0.21 L/kg	NR	NR	5.6 ± 1.8
141.2 ± 36.2 mL/min	Intermittent infusion; 1000 mg every 12 h	The first dose: prior to injection, at the end of the infusion, and 11 samples collected until 11 h after the end of the infusion	G-Pharm computer program	Two-compartment model	V <sub>c</sub> : 22.9 ± 11.4 L	158 ± 51 mL/min	NR	2.94 ± 0.84
85.72 ± 37.28 mL/min/1.73m <sup>2</sup>	Intermittent infusion; 1970.00 ± 605.19 mg/d	Three samples: 1 h, 3–8 h, and 8–24 h post-infusion	Maximum a priori Bayesian estimation using Adapt 5	Two-compartment model	V <sub>c</sub> : 0.23 L/kg V <sub>ss</sub> : 0.60 L/kg	0.14 L/h/kg	NR	NR
104.7 ± 37.0 mL/min	Intermittent infusion; 31.9 (±10.5) mg/kg/d (69.6%:1g q12h; 17.4%:1g q8h)	First steady-state trough (within 30 minutes prior to the fourth dose), peak concentration, random sample	Equations with two steady-state samples	One-compartment model	0.60 (0.44–0.76) L/kg	0.090 (0.071–0.109) L/h/kg 109.7 (82.7–136) mL/min	0.16 (0.13–0.19)	4.9 (3.8–6.0)
89.4 ± 39.2 mL/min	Intermittent infusion	Blood sampling was ordered as required clinically	Nonlinear mixed-effect modeling approach (NONMEM)	One-compartment model	CL (L/h): 1.08 × CL <sub>CR</sub> (Cockcroft and Gault) (L/h); CV <sub>CL</sub> : 28.16% V (L) = 0.98 × TBW; CV <sub>V</sub> : 37.15%.			
113 (range: 47–253) mL/min	Intermittent infusion initial dosage of 1 g/12 hours (if the CL <sub>CR</sub> was >75 mL/min/1.73 m <sup>2</sup> ).	Immediately before administering vancomycin, 1 hour after drug administration and at some other points as necessary	Nonlinear mixed-effect modeling approach (NONMEM)	Two-compartment model	V <sub>c</sub> (L) = 39.2 × (TBW/59.4) <sup>0.78</sup> ; CV <sub>V<sub>c</sub></sub> = 14.2% CL (L/h) = 4.25 × (CL <sub>CR</sub> /113) <sup>0.70</sup> ; CV <sub>CL</sub> = 25.2% V <sub>p</sub> (L) = 56.1; CV <sub>V<sub>p</sub></sub> = 66.9%			
Median 92.7 mL/min	Intermittent infusion The specific dosing was not pre-specified	NR	Nonlinear mixed-effect modeling (NONMEM)	Two-compartment model	CL(L/h) = 3.22 + (1 + 0.00834 × (CL <sub>CR</sub> - 104)) × 1.277 <sup>NEUTROPENIA</sup> ; CV <sub>CL</sub> = 33.0%			

## Acknowledgments

Thanks for Xiaotong Li's help for risk of bias assessment during the revision process.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Matzke GR, Aronoff GR, Atkinson AJ, et al. Drug dosing consideration in patients with acute and chronic kidney disease—a clinical update from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int*. 2011;80(11):1122–1137. doi:10.1038/ki.2011.322
- Matsumoto K, Takesue Y, Ohmagari N, et al. Practice guidelines for therapeutic drug monitoring of vancomycin: a consensus review of the Japanese Society of Chemotherapy and the Japanese Society of Therapeutic Drug Monitoring. *J Infect Chemother*. 2013;19(3):365–380. doi:10.1007/s10156-013-0599-4
- Bilbao-Meseguer I, Rodriguez-Gascon A, Barrasa H, Isla A, Solinis MA. Augmented renal clearance in critically ill patients: a systematic review. *Clin Pharmacokinet*. 2018;57(9):1107–1121. doi:10.1007/s40262-018-0636-7
- Udy AA, Jarrett P, Lassig-Smith M, et al. Augmented renal clearance in traumatic brain injury: a single-center observational study of atrial natriuretic peptide, cardiac output, and creatinine clearance. *J Neurotrauma*. 2017;34(1):137–144. doi:10.1089/neu.2015.4328
- Kim AJ, Lee JY, Choi SA, Shin WG. Comparison of the pharmacokinetics of vancomycin in neurosurgical and non-neurosurgical patients. *Int J Antimicrob Agents*. 2016;48(4):381–387. doi:10.1016/j.ijantimicag.2016.06.022
- Keng MK, Sekeres MA. Febrile neutropenia in hematologic malignancies. *Curr Hematol Malig Rep*. 2013;8(4):370–378. doi:10.1007/s11899-013-0171-4
- Freifeld AG, Bow EJ, Sepkowitz KA, et al. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the infectious diseases society of america. *Clin Infect Dis*. 2011;52(4):e56–e93.
- Goulenok T, Fantin B. Antimicrobial treatment of febrile neutropenia: pharmacokinetic-pharmacodynamic considerations. *Clin Pharmacokinet*. 2013;52(10):869–883. doi:10.1007/s40262-013-0086-1
- Lortholary O, Lefort A, Tod M, Chomat AM, Darras-Joly C, Cordonnier C. Pharmacodynamics and pharmacokinetics of antibacterial drugs in the management of febrile neutropenia. *Lancet Infect Dis*. 2008;8(10):612–620. doi:10.1016/S1473-3099(08)70228-7
- Ye ZK, Chen YL, Chen K, et al. Therapeutic drug monitoring of vancomycin: a guideline of the division of therapeutic drug monitoring, chinese pharmacological society. *J Antimicrob Chemother*. 2016;71(11):3020–3025. doi:10.1093/jac/dkw254
- Rybak M, Lomaestro B, Rotschafer JC, et al. Therapeutic monitoring of vancomycin in adult patients: a consensus review of the American Society of Health-System Pharmacists, the Infectious Diseases Society of America, and the Society of Infectious Diseases Pharmacists. *Am J Health Syst Pharm*. 2009;66(1):82–98. doi:10.2146/ajhp080434
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097
- Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0. Cochrane Collaboration; 2011:1–8.
- Wells G, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality if nonrandomized studies in meta-analyses. Available from: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.htm](http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm). Accessed May 11, 2020.
- Kanji S, Hayes M, Ling A, et al. Reporting guidelines for clinical pharmacokinetic studies: the ClinPK statement. *Clin Pharmacokinet*. 2015;54(7):783–795. doi:10.1007/s40262-015-0236-8
- O'Donnell R. Audit of BMT antibiotic febrile neutropenia guidelines. *Bone Marrow Transplant*. 2011;46:S433–S434.
- Luo C, Hussaini T, Lalaria K, Yeung J, Lau TT, Broady RC. Evaluation of a Once-Daily Vancomycin regimen in an outpatient leukemia/bone marrow transplant clinic (OD-VANCO Study). *Can J Hosp Pharm*. 2014;67(4):280–285. doi:10.4212/cjhp.v67i4.1372
- Donovan KA, Sklenicka J. Is a standard vancomycin dosing protocol appropriate to achieve therapeutic trough levels in adult neutropenic patients? *Pharmacotherapy*. 2012;32(10):e237–e238.
- Vermis K, Steel E, Vandenbroucke J. Prevalence of augmented renal clearance in haematological patients and the impact on vancomycin dosing. *J Oncol Pharm Pract*. 2014;20(3):7.
- Ghehi MT, Rezaee S, Hayatshahi A, et al. Vancomycin pharmacokinetic parameters in patients undergoing Hematopoietic Stem Cell Transplantation (HSCT). *Int J Hematol Oncol Stem Cell Res*. 2013;7(4):1–9.
- Hochart C, Berthon C, Corm S, et al. Vancomycin serum concentration during febrile neutropenia in patients with acute myeloid leukemia. *Med Mal Infect*. 2011;41(12):652–656. doi:10.1016/j.medmal.2011.09.014
- Vazin A, Japoni A, Shahbazi S, Davarpanah MA. Vancomycin utilization evaluation at hematology-oncology ward of a teaching hospital in iran. *Iran J Pharm Res*. 2012;11(1):163–170.
- Choi MH, Choe YH, Lee SG, Jeong SH, Kim JH. Neutropenia is independently associated with sub-therapeutic serum concentration of vancomycin. *Clin Chim Acta*. 2017;465:106–111. doi:10.1016/j.cca.2016.12.021
- Al-Kofide H, Zaghoul I, Al-Naim L. Pharmacokinetics of vancomycin in adult cancer patients. *J Oncol Pharm Pract*. 2010;16(4):245–250. doi:10.1177/1078155209355847
- Haeseker MB, Croes S, Neef C, Bruggeman CA, Stolk LM, Verbon A. Vancomycin dosing in neutropenic patients. *PLoS One*. 2014;9(11):e112008. doi:10.1371/journal.pone.0112008
- Kureishi A, Jewesson PJ, Bartlett KH, Cole CD, Chow AW. Application of a modified bioassay for monitoring serum teicoplanin and vancomycin in febrile neutropenic patients. *Antimicrob Agents Chemother*. 1990;34(9):1642–1647. doi:10.1128/AAC.34.9.1642
- Jarkowski IA, Forrest A, Sweeney RP, et al. Characterization of vancomycin pharmacokinetics in the adult acute myeloid leukemia population. *J Oncol Pharm Pract*. 2012;18(1):91–96. doi:10.1177/1078155211402107
- Le Normand Y, Milpied N, Kergueris MF, Harousseau JL. Pharmacokinetic parameters of vancomycin for therapeutic regimens in neutropenic adult patients. *Int J Biomed Comput*. 1994;36(1–2):121–125. doi:10.1016/0020-7101(94)90102-3
- Buelga DS, Fernandez de Gatta MM, Herrera EV, Dominguez-Gil A, Garcia MJ. Population pharmacokinetic analysis of vancomycin in patients with hematological malignancies. *Antimicrob Agents Chemother*. 2005;49(12):4934–4941. doi:10.1128/AAC.49.12.4934-4941.2005
- Hirai K, Ishii H, Shimoshikiryō T, et al. Augmented renal clearance in patients with febrile neutropenia is associated with increased risk for subtherapeutic concentrations of vancomycin. *Ther Drug Monit*. 2016;38(6):706–710. doi:10.1097/FTD.0000000000000346
- Soto J, Alsar MJ, Chantal P, Sacristan JA. Correlation of vancomycin clearance and creatinine clearance: unreliability for predicting initial dosing in neutropenic haematological patients. *J Antimicrob Chemother*. 1993;32(6):920–922. doi:10.1093/jac/32.6.920
- Taghizadeh-Ghehi M, Rezaee S, Gholami K, Hadjibabaie M. Predictive performance of vancomycin population pharmacokinetic models in Iranian patients underwent hematopoietic stem cell transplantation. *J Res Pharm Pract*. 2015;4(3):129–134. doi:10.4103/2279-042X.162357

33. Fernandez de Gatta MM, Santos BD, Sanchez NA, Dominguez-Gil A, Garcia MJ. Vancomycin dosage optimization in patients with malignant haematological disease by pharmacokinetic/pharmacodynamic analysis. *Clin Pharmacokinet.* 2009;48(4):273–280. doi:10.2165/00003088-200948040-00005
34. Fernandez de Gatta MM, Calvo MV, Hernandez JM, Caballero D, San MJ, Dominguez-Gil A. Cost-effectiveness analysis of serum vancomycin concentration monitoring in patients with hematologic malignancies. *Clin Pharmacol Ther.* 1996;60(3):332–340. doi:10.1016/S0009-9236(96)90060-0
35. Suzuki Y, Tokimatsu I, Morinaga Y, et al. A retrospective analysis to estimate target trough concentration of vancomycin for febrile neutropenia in patients with hematological malignancy. *Clin Chim Acta.* 2015;440:183–187. doi:10.1016/j.cca.2014.11.027
36. Bury D, Ter Heine R, van de Garde E, et al. The effect of neutropenia on the clinical pharmacokinetics of vancomycin in adults. *Eur J Clin Pharmacol.* 2019;75(7):921–928. doi:10.1007/s00228-019-02657-6
37. Izumisawa T, Kaneko T, Soma M, et al. Augmented renal clearance of vancomycin in hematologic malignancy patients. *Biol Pharm Bull.* 2019;42(12):2089–2094. doi:10.1248/bpb.b19-00652
38. Okada A, Kariya M, Irie K, et al. Population pharmacokinetics of vancomycin in patients undergoing allogeneic hematopoietic stem-cell transplantation. *J Clin Pharmacol.* 2018;58(9):1140–1149. doi:10.1002/jcph.1106
39. Matzke GR, Zhanel GG, Guay DR. Clinical pharmacokinetics of vancomycin. *Clin Pharmacokinet.* 1986;11(4):257–282. doi:10.2165/00003088-198611040-00001
40. Matzke GR, McGory RW, Halstenson CE, Keane WF. Pharmacokinetics of vancomycin in patients with various degrees of renal function. *Antimicrob Agents Chemother.* 1984;25(4):433–437. doi:10.1128/AAC.25.4.433
41. Haeseker M, Croes S, Neef C, Bruggeman C, Stolk L, Verbon A. Evaluation of vancomycin prediction methods based on estimated creatinine clearance or trough levels. *Ther Drug Monit.* 2016;38(1):120–126. doi:10.1097/FTD.0000000000000250
42. Fernandez de Gatta MM, Fruns I, Dominguez-Gil A. Individualizing vancomycin dosing regimens: an evaluation of two pharmacokinetic dosing programs in critically ill patients. *Pharmacotherapy.* 1994;14(2):196–201.
43. Krivoy N, Peleg S, Postovsky S, Ben AM. Pharmacokinetic analysis of vancomycin in steady state in pediatric cancer patients. *Pediatr Hematol Oncol.* 1998;15(4):333–338. doi:10.3109/08880019809014017
44. Omote S, Yano Y, Hashida T, et al. A retrospective analysis of vancomycin pharmacokinetics in Japanese cancer and non-cancer patients based on routine trough monitoring data. *Biol Pharm Bull.* 2009;32(1):99–104. doi:10.1248/bpb.32.99
45. Toro JJ, Diaz-Duque AE, Gass MJ, Haile DJ, Gushiken F. Acute renal injury and vancomycin levels in patients undergoing autologous hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant.* 2017;23(3):S139. doi:10.1016/j.bbmt.2016.12.269
46. Mae H, Ooi J, Takahashi S, et al. Early renal injury after myeloablative cord blood transplantation in adults. *Leuk Lymphoma.* 2008;49(3):538–542. doi:10.1080/10428190701824577

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>