

ARTICLE

Open Access

Neonatal sevoflurane exposure induces impulsive behavioral deficit through disrupting excitatory neurons in the medial prefrontal cortex in mice

Linghua Xie¹, Yue Liu², Yuhan Hu³, Bei Wang², Zhirui Zhu², Yilei Jiang², Yaojun Suo², Miaofeng Hu², Jing Gao², Rahim Ullah⁴ and Zhiyong Hu¹ 

Abstract

Sevoflurane, in particular multiple exposures, has been reported to cause the abnormal neurological development including attention-deficit/hyperactivity disorder (ADHD). This study is to investigate ADHD-like impulsivity in adult mice after repeated sevoflurane exposures at the neonatal stage. Six-day-old pups were exposed to 60% oxygen in the presence or absence of 3% sevoflurane for 2 h and the treatment was administrated once daily for three consecutive days. To assess the impulsivity, the cliff avoidance reaction (CAR) was carried out at the 8th week. Our results showed that repeated sevoflurane treatment increased the number of jumps and shortened the jumping latency in the CAR test. The cortices were harvested for immunostaining to detect c-Fos and calmodulin-dependent protein kinase IIa (CaMKIIa) expression in the medial prefrontal cortex (mPFC). We found that mPFC neurons, especially excitatory neurons, were highly activated and related to impulsive behavior. The activation viruses (AAV-CaMKIIa-hM3Dq) were injected to evaluate the effects of specific activation of mPFC excitatory neurons on impulsive behavior in the presence of clozapine-N-oxide (CNO). Likewise, the inhibitory viruses (AAV-CaMKIIa-hM4Di) were injected in the sevoflurane group to explore whether the mPFC excitatory neuronal inhibition reduced the impulsivity. Our results revealed that chemogenetic activation of mPFC excitatory neurons induced impulsive behavior whereas inhibition of mPFC excitatory neurons partially rescued the deficit. These results indicate that repeated sevoflurane exposures at the critical time induce impulsive behavior accompanied with overactivation of mPFC excitatory neurons in adult stages. This work may further extend to understand the ADHD-like impulsive behavior of the anesthetic neurotoxicity.

Introduction

Attention-deficit/hyperactivity disorder (ADHD) is a chronic neurodevelopmental disorder manifesting primarily as attention deficit, hyperactivity, and impulsivity¹. ADHD affects 5–8% of school-age children, in which 50–80% cases persist till adulthood². Adult ADHD has complex manifestations. Currently, the core symptom used in adult ADHD diagnosis is impulsivity, defined as

poorly conceived control deficit^{3,4}. Impulsivity increases the risk of pathological gambling, substance abuse, and relationship failure. As the prevalence rate of adult ADHD is soaring over the world, therefore, neurobiological and psychosocial factors related to the etiology and symptom control of ADHD have been studied in depth—exemplified by genes, neurotransmitters, neural pathways, drugs, and education.

Patients with ADHD show remarkable variation in complicating factors and neuropsychological weaknesses⁵. Previous studies suggested that the impaired development of dopaminergic neurons is related to ADHD development⁶. In addition to disorders in neurotransmitter system, other studies indicated that ADHD is caused by

Correspondence: Zhiyong Hu (huzhiyong777@zju.edu.cn)

¹Department of Anesthesiology, The First Affiliated Hospital, The Children's Hospital, Zhejiang University School of Medicine, Hangzhou, China

²Department of Anesthesiology, The Children's Hospital, Zhejiang University School of Medicine, Hangzhou, China

Full list of author information is available at the end of the article

© The Author(s) 2020



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

abnormal structural and functional synapses in mPFC^{6,7}, which is responsible for behavioral inhibition and executive functions. Recently, clinical studies reported that repeated exposures to inhalational anesthetics in children is associated with the development of ADHD^{6,8–10}. This finding was further supported by an animal study, which revealed that ketamine administration caused ADHD-like hyperactive behavior¹¹. However, whether anesthetics have a long-term effect on ADHD development, particularly impulsive behavior, remains uncertain. Given the potential issue of public mental health and social stability, identifying the links between general anesthesia and impulse control disorder is now of paramount importance. Sevoflurane, the most commonly used pediatric anesthetic, has been shown to act as a neurotoxin affecting synaptogenesis and neuronal morphology in animal studies^{12,13}. Therefore, we hypothesize that exposure of the immature brain to sevoflurane may induce ADHD-like behavior conditions in later adulthood.

In this study, we exposed neonatal mice to sevoflurane and found it induced ADHD-like impulsive behavior in the cliff avoidance reaction (CAR) test in adult age. We revealed that sevoflurane permanently affected the mPFC excitatory neurons, which were involved in impulsive behavior. We also demonstrated that chemogenetic inhibition of the mPFC excitatory neurons rescued sevoflurane-induced impulsive behavior. These results suggest that sevoflurane-induced impulsive behavior is another type of long-term outcome to anesthetic neurotoxicity. Furthermore, excitatory neuronal activation in the mPFC may serve as a pathological mechanism for anesthetics-induced ADHD in adults.

Materials and methods

Animals

All procedures were performed following the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and approved by the Animal Advisory Committee at Zhejiang University. C57BL/6J mice were group-housed in a temperature-controlled room at the Animal Facility of Zhejiang University under a 12 h/12 h light/dark cycle. Food and water were available ad libitum. Male and female mice were allowed to meet for one night and then the females were housed individually. Time mated birth was monitored and pups were collected for further experiments.

Exposure to sevoflurane

Animal groups were determined in a pilot study and randomly assigned to the sevoflurane group and control group before the experiment. At postnatal day 6 (P6), pups were placed in a translucent plastic chamber (20 × 10 × 10 cm³) with a layer of soda-lime and wet cotton on the bottom floor. The temperature of the chamber floor

was maintained at 32 °C using a heating pad under the chamber. The chamber was connected to a sevoflurane vaporizer (Vapor 2000; Medical Systems, Inc., Germany) and continuously flushed with 60% oxygen balanced nitrogen (FIO₂ = 0.6) at 2 L/min for 2 h with (the sevoflurane group) or without (the control group) 3% sevoflurane. The treatment was administered once per day for 3 consecutive days. Mice were allowed to grow up to 8 weeks. Female mice were excluded to avoid the possible confounding effects of the estrous cycle and only males were used for further experiments.

Cliff avoidance reaction (CAR)

One cohort of mice ($n = 24$) was evaluated for the CAR test. CAR is a behavioral paradigm used to investigate the ADHD-like impulsive behavior in mature rodents^{14–16}. It depends on the natural tendency of animals to avoid falling from a height, which is more than twice the length of the animal¹⁷. On account of a control deficit and risk-taking choice, the impulsive mice were more active in exploring the cliff, which led to a jump from the high platform at a higher frequency. According to a previous study¹⁸, a round platform (an inverted glass beaker with 11 cm diameter and 15 cm height) placed in the center of the open field apparatus (45 × 45 × 45 cm³) was used for the CAR test. The behavioral experiments were conducted in a quiet room illuminated by dim light between 9 and 12 am. Before the tests, the mice ($n = 24$) were handled in a friendly manner once/day for 3 consecutive days to decrease the stress-related neuronal activation. On the test day, mice were first allowed to habituate in the test room for more than 30 min. Each mouse was then gently placed on the platform and the duration that it remained on the platform (from the initial placement until falling onto the floor) was recorded. During the whole 15 min testing period, the mice were repeatedly placed on the platform once they fell off, and the cumulative cliff-falling events were documented. The latency is defined as the duration when mice stayed on the platform within the 15 min testing period. The apparatus was cleaned with 75% ethanol between tests to prevent olfactory influence.

Elevated plus-maze (EPM) test

Another cohort of mice ($n = 16$) were evaluated for the EPM test. The EPM consists of two open arms and two closed arms (30 × 5 cm²) in an arrangement that the two arms of each type are opposite to each other. The maze was placed at the height of 65 cm above the ground. A video camera was fixed above the maze to record the movements of mice. Each mouse was placed in the center area facing the open arms and allowed to explore freely for 5 min. Time spent in the open arms, the number of entries into the open arms and freezing time were calculated using the video-based ANY-maze system

(Stoelting, USA). The apparatus was cleaned with 75% ethanol between tests.

Immunohistochemistry

90 min after completing the CAR test, 6 mice were sacrificed under terminated anesthesia the CAR test according to a random number table, and then perfused with 4% paraformaldehyde (PFA). Their brains were harvested for immunohistochemistry analysis. Coronal brain sections (40- μ m thickness) were cut by a freezing microtome (CM30503, Leica Biosystems, Germany). After 2-h blocking in 5% normal donkey serum (NDS; 017-000-121, Jackson ImmunoResearch Laboratories, Inc. USA), 1% bovine serum albumin (BSA; A2153, Sigma-Aldrich, USA), and 0.3% Triton X-100 (Sigma-Aldrich, USA), the sections were incubated with the primary antibodies (mouse anti-c-Fos [1:1,000; Abcam, ab208942] and/or rabbit anti-CaMKII α [1:1000; Abcam, ab52476]) overnight at 4°C. Brain sections were then incubated with secondary antibodies (Alexa Fluor 488 anti-mouse IgG [1:1,000; A-21202, Thermo Fisher Scientific] or Alexa Fluor 546 anti-rabbit IgG [1:1,000; A-10040, Thermo Fisher Scientific]). The processed brain sections were then mounted onto glass slides with 60% glycerol in PBS. Images were captured using a Nikon A1 laser-scanning confocal microscope (Nikon, Japan). The number of immune-positive cells was counted and analyzed by ImageJ (NIH, USA). DAPI (4',6-dimethyl-2-phenylindole; 1:3,000; C1002; Beyotime) was used simultaneously to identify nuclei.

Surgery and chemogenetic manipulation

Under sodium pentobarbital anesthesia (50 mg/kg, intraperitoneal injection (i.p.)), other cohorts were mounted in a stereotaxic apparatus (RWD Life Science, Shenzhen, China). After adjusting the position of the skull horizontally under a stereomicroscope (RWD Life Science, Shenzhen, China)¹⁹, a small craniotomy was made with a thin drill over the medial prefrontal cortex (typical coordinate: 1.9 mm anterior to Bregma; 0.34 mm lateral to the midline). Adeno-associated virus (AAV) carrying fusion genes for hM3D (AAV-CaMKII α -hM3D(Gq)-mCherry), hM4D (AAV-CaMKII α -hM4D(Gi)-mCherry), or mcherry (AAV-CaMKII α -mcherry) were injected using a 10 μ L micro-syringe (#701, Hamilton, USA) at a rate of 60 nL/min by micro-syringe pump (kdScientific, USA). Over a 10-min period, 80 nL viruses were injected at a depth of 2.4 mm from the Bregma. Mice with viral injections in the incorrect area or expressions beyond the targeted area (mPFC) were excluded from the study. The hM3Dq was injected unilaterally for activation ($n = 7$) and hM4Di ($n = 8$) was bilaterally injected for inhibition. The control adenovirus (mCherry only) was also injected unilaterally or bilaterally for corresponding control

experiments. The viruses hM3Dq (AAV_{2/9}, 3.6×10^{12} genomic copies per ml), hM4Di (AAV_{2/9}, 3.6×10^{12} genomic copies per mL), or mcherry (AAV_{2/9}, 3.54×10^{12} genomic copies per mL) were made by HanBio (Shanghai, China).

The mice injected with the above viruses were received a saline i.p. to test the baseline of the CAR deficit. After that, mice were given CNO (1 mg/kg i.p.; clozapine-N-oxide; HY17366, MedchemExpress, USA) for virus-infected neuron manipulation.

Electrophysiological recordings of brain slices

After a 3–4-week viral incubation, coronal sections including the mPFC were cut at a thickness of 300 μ m using a vibratome (VT 1200S, Leica, Germany) in ice-cold cutting solution (234 mM Sucrose, 5 mM KCl, 1.25 mM NaH₂PO₄, 5 mM MgSO₄, 26 mM NaHCO₃, 25 mM dextrose, 1 mM CaCl₂, oxygenated with 95% O₂/ 5% CO₂) as described previously¹⁹. Next, slices were incubated in the aerated (95% O₂/5% CO₂) artificial cerebrospinal fluid (ACSF: 124 mM NaCl, 2 mM KCl, 1.25 mM KH₂PO₄, 2 mM MgSO₄, 2 mM CaCl₂, 26 mM NaHCO₃, and 10 mM D-(C) glucose, pH 7.4.) to recover for 30 min at 32°C, and subsequently incubated for 1 h at room temperature. Brain slices were then transferred to a recording chamber. To test function of expressed hM3Dq or hM4Di protein, whole-cell patch-clamp recordings were applied with glass pipettes backfilled with an artificial intracellular fluid (120 mM K-gluconate, 15 mM KCl, 10 mM HEPES, 4 mM Mg-ATP, 0.3 mM Tris-GTP, 0.5 mM EGTA, adjusted to pH 7.3 with KOH, 285–290 mOsm). Pipettes were connected to the headstage of a Heka EPC 10 amplifier (Heka Elektronik, USA). Cells expressing viral-encoded fluorescent markers (mCherry) were visualized under microscope with infrared differential interference contrast optics (BX51WI, Olympus, Japan). Current-clamp recording was used to measure evoked action potentials in both CNO activation and inhibition experiment. After applying currents in steps of 20 pA (from 0 to 360 pA) neurons were recovered for 5 min. The brain slices were then perfused with ACSF containing 5 μ M CNO. The same current-clamp procedure was performed 10 min after CNO perfusion.

Statistical analysis

The sample size was estimated by software PASS.11 (NCSS, USA) according to the preliminary experimental results. Experienced researchers conducted independently the different experiments and blinded data collectors analyzed these results. All data are shown as mean \pm SEM unless otherwise specified. Before analysis, all data underwent the Kolmogorov–Smirnov normality test. Comparisons between two groups were made with unpaired or paired Student's *t*-test if data are normally

distributed; otherwise, Mann–Whitney U test or Wilcoxon signed-rank tests were used. Variance was found to be similar between the groups as tested using the Levene test of homogeneity of variances. Two-way ANOVA followed by Bonferroni's post-hoc test was used for multiple comparisons, as appropriate. p -value less than 0.05 was considered to be statistically significant, defined as $*p < 0.05$; $**p < 0.01$; $***p < 0.001$; and $****p < 0.0001$. Calculation was performed in GraphPad Prism TM 8.0 (version 8.0; GraphPad Software Inc., USA) and SPSS 22.0 (IBM Corp., USA).

Result

General data

During sevoflurane exposure, pups were in pink color and respiration was maintained well and there were no deaths throughout the whole process.

Postnatal sevoflurane exposure showed ADHD-like impulsive behavior and anxiety-free in adulthood

To investigate whether neonatal sevoflurane exposure induces voluntary risk-taking behavior in adulthood, the CAR test was performed on 8-week-old mice that have been treated either with sevoflurane or mock-control at the neonatal stage. An inverted glass beaker was placed in the center area of the open field apparatus, mice were carefully placed on the platform and the CAR test was initiated (Fig. 1a). After an adaptation period in the test room, each mouse was evaluated for 15 min and their movements were monitored by a video camera (Fig. 1b; Supplemental Video 1, 2). Sevoflurane-treated mice showed marked abnormalities with significantly increased jumping events ($p < 0.0001$, Fig. 1c) and decreased latency of falling from the platform compared with controls ($p < 0.0001$, Fig. 1d). A previous study showed that anxiety or fear itself could also result in impulsive-like behavior²⁰. To exclude the contribution of anxiety and/or fear to impulsive-like behavior in our animal model, an EPM test was performed. The EPM test showed no significant difference in the time spent in open arms ($p = 0.067$, Fig. 1e) and entries to the open arms ($p = 0.0594$, Fig. 1f) between the groups. These data suggest that sevoflurane exposure has no impact on inducing anxiety. Moreover, the sevoflurane group exhibits significantly less freezing time ($p = 0.0104$, Fig. 1g). These results indicate that postnatal exposure to sevoflurane can induce ADHD-like impulsive behavior.

Enhanced mPFC neuronal activities are associated with the sevoflurane-induced impulsivity


c-Fos has been reported as a reliable marker for neuronal activation and its expression is induced by various behavioral stimuli²¹. Therefore, we labeled brain sections with c-Fos to identify specific areas responsible for CAR

behavior in our animal model. We counted the c-Fos expressing neurons in various brain regions, including mPFC, caudate putamen (CPU), anterior paraventricular thalamic nucleus (aPVT), basolateral amygdaloid nucleus (BLA) and bed nucleus of the stria terminalis (BNST) 90 min after the CAR test (Fig. 2a). Interestingly, sevoflurane-treated mice showed a notable increased expression of c-Fos in mPFC compared with control ($F(7, 52) = 35.5, p = 0.0006$, Fig. 2b). We found there were more CPU c-Fos positive cells in the sevoflurane-treated mice compared with control ($p = 0.0382$, Fig. 2b). No significant difference in c-Fos positive cells was identified between aPVT and posterior paraventricular thalamic nucleus (pPVT) between sevoflurane-treated mice and control (Fig. 2b). Furthermore, the c-Fos expression in anxiety- and fear-related brain regions, including BNST²² and BLA²³ showed no significant difference between sevoflurane-treated mice and control, which is also consistent with the EPM results (Fig. 2b). It also showed that c-Fos expression in CA3 (an area of the hippocampus) and the hypothalamus (HP) has no significant difference between the treatment group and control (Fig. 2b).

Different neuronal subtypes in mPFC play various roles in encoding and regulating behaviors²⁴. To examine what are the specific types of neurons that involve in impulsive behaviors, we performed double-immunofluorescent staining of CaMKII α and c-Fos (Fig. 2c) for sevoflurane-treated mice and control. We found that sevoflurane-treated mice are possessing a significantly greater population of c-Fos positive cells expressing CaMKII α ($p = 0.0042$, Fig. 2d). In the sevoflurane-treated group, 84.7 % of c-Fos positive cells were colocalized with the CaMKII α^+ neurons (Fig. 2d). Together these results suggest that excitatory neurons in mPFC were over-activated in sevoflurane-treated mice.

Chemogenetic activation of mPFC excitatory neurons induces ADHD-like impulsive behavior

It is known that mPFC is responsible for risk-decision making⁴. To test whether mPFC excitatory neural activities are involved in ADHD-like impulsive behavior, the DREADD (designer receptors exclusively activated by designer drugs)-based tools were used. We unilaterally injected excitatory virus (hM3Dq-mCherry) or non-functional control virus (mCherry) in mPFC (Fig. 3a). To ensure that the virus is effective, the whole-cell recording was performed in mPFC neurons. We confirmed that CNO administration activated hM3Dq-expressing neuronal activity in mPFC (Fig. 3b) by inducing a decreased spike threshold ($p = 0.0478$, Fig. 3c) and increased spike number under current step injections (Fig. 3d). Furthermore, we injected (i.p) CNO (1 mg/kg) into both mCherry-injected and hM3Dq-injected mice and traced the movements of hM3Dq-injected mice and



control (Fig. 3e). We observed a significant increase in jumping events ($p = 0.0006$, Fig. 3f) and less latency ($p = 0.0056$, Fig. 3g) in the hM3Dq-injected group compared with the control group (mCherry-injected). Consistently, the activation of mPFC after CNO administration significantly increased the number of jumping events ($p = 0.0156$, Fig. 3h) and decreased the latency compared with the baseline level of CAR assessed by saline i.p. injection ($p = 0.0156$, Fig. 3i). Finally, we evaluated the anxiety level of mice using EPM test and found that time spent in the open arms ($p = 0.6904$, Fig. 3j) and the number of entries to the open arms ($p = 0.9485$, Fig. 3k) showed no significant difference between hM3Dq-injected mice and control. No significant difference was found in freezing time between the two groups ($p = 0.7575$, Fig. 3l). Together, our data indicate that chemogenetic activation of the mPFC neurons was sufficient to induce impulsive behavior deficit without affecting the freezing time or anxiety level.

Chemogenetic inhibition of neurons in the mPFC attenuates the sevoflurane-induced impulsive performance

If the amplified activity of the mPFC excitatory neurons is responsible for impulsivity, then inhibition of these

neurons may rescue the phenotype. To prove our hypothesis, we bilaterally injected hM4Di-mCherry and mCherry encoding virus into the mPFC in the sevoflurane-treated mice (Fig. 4a). The effectiveness of the viruses was also verified by electrophysiological recordings. CNO administration inhibited spiking in the hM4Di-expressing mPFC neurons (Fig. 4b), resulting in an increased spike threshold ($p = 0.0013$, Fig. 4c) and a reduction in spike number under the current step injection (Fig. 4d). Movements of mice were traced after CNO i.p. injection (Fig. 4e). We found that inhibition of the mPFC neurons dramatically reduced the jumping events ($p = 0.0006$, Fig. 4f) and increased the latency ($p = 0.0023$, Fig. 4g). Consistently, the CNO injection into the sevoflurane-treated mice showed fewer jumping events ($p = 0.0078$, Fig. 4h) and longer latency ($p = 0.0156$, Fig. 4i), compared with the baseline of the CAR assessed by saline i.p. injection. The administration of CNO has no effect on anxiety level (time in open arms, $p = 0.095$; open arms entries, $p = 0.6412$; Fig. 4j, k) or fear behavior ($p = 0.0803$, Fig. 4l). Our data suggest that inhibition of mPFC excitatory neurons ameliorate ADHD-like impulsive behavior for sevoflurane-treated mice.




Fig. 2 Enhanced mPFC neuronal activities are associated with the sevoflurane-induced impulsivity. **a** Representative images show the c-Fos positive cells in mPFC, CPU, aPVT, BLA, BNST from control (top), and sevoflurane (bottom) mice. Scale bar, 100 μm. mPFC, medial prefrontal cortex; CPU, caudate putamen, striatum; aPVT anterior paraventricular thalamic nucleus; pPVT posterior paraventricular thalamic nucleus; BLA basolateral amygdaloid nucleus; BNST bed nucleus of the stria terminalis. (dotted line show the boundary of brain area). **b** Quantification of c-Fos positive cells in the mPFC, CPU, aPVT, pPVT, BNST, BLA, CA3, and HP, respectively. HP, hypothalamus area. The number of c-Fos expression in the mPFC is markedly higher for sevoflurane mice ($n = 6$, Two-way ANOVA, Bonferroni multiple comparison post hoc tests). **c** Representative micrograph showing the c-Fos⁺ neurons (green), CaMKIIα⁺ neurons (red) and colocalized cells. Scale bar (left) 100 μm; (right) 50 μm. (arrows = double-labeled cells). **d** Quantification of the number of c-Fos⁺ neurons that colocalized with CaMKIIα⁺ neurons (top). Percentage of CaMKIIα⁺ cells expressing c-Fos in the sevoflurane mice (bottom). ($n = 6$; Unpaired Student's *t* test). Data = mean ± SEM; n.s. no significance, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Discussion

These results demonstrated that repeated exposure to sevoflurane at the developmental stage causes ADHD-like

impulsive behavior coupled with overactivated excitatory mPFC neurons later in adulthood. We also showed that selective activation of excitatory neurons potentially




Fig. 3 (See legend on next page.)

(see figure on previous page)

Fig. 3 Chemogenetic activation of mPFC excitatory neurons induces ADHD-like impulsive behavior. **a** Schematic diagram (coronal section) showing the target of viral injection in the mPFC (top). Location of unilaterally viral hM3D (Gq) expression (red) in mPFC under control of the CaMKII α promoter (bottom). **b** Current–voltage relationship of a representative mPFC neuron recorded before and during 5 μ M CNO perfusion. Raw traces show individual voltage responses to a series of 600-ms current pulses from 0 to 360 pA with 20 pA steps. **c** CNO decreased the minimal injected current to induce action potential (AP). ($n = 10$, Paired Student's t test). **d** The number of induced action potentials at different current steps. ($n = 10$, two-way ANOVA, Bonferroni multiple comparison post hoc tests.). **e** Representative movement traces showing the locations of mcherry-expressing (top) and hM3Dq-expressing in the air-exposure mice (bottom) in a cliff avoidance reaction test following CNO (1 mg/kg) administration. **f, g** Quantification of the number of cumulative cliff jumping events (left) and the latency (right) of the first jump from the cliff in the mcherry group and hM3Dq group ($n = 7$ per group; Mann–Whitney U test for the number; Unpaired Student's t test for the latency). **h, i** Quantification of the number of cumulative cliff jumping events (left) and the latency (right) of the first jump from the cliff in the baseline group and hM3Dq group. ($n = 7$; Wilcoxon matched-pairs signed-rank test). **j–l** Quantification of the EPM results of the time spent in the open arms, open arms entries, freezing time of hM3Dq and mcherry group after chemogenetic activation of mPFC glutamatergic neurons. ($n = 7$; Unpaired Student's t test). Data = mean \pm SEM; n.s. no significance, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

promoted impulsive behavior and chemogenetic inhibition of these neurons abolished the sevoflurane-induced impulsive behavior in adult mice. Our findings potentially provide insights for understanding the underlying mechanisms for anesthetics-related ADHD.

It has been reported that repeated, but not a single, anesthetic exposure during early postnatal development results in neurological impairment^{25–27}, including neurocognitive impairments^{28,29}. However, recent clinical studies indicated that children who received repeated anesthetic exposure had deficits in executive function and motor processing (ADHD)^{30–32}. In this study, our finding suggests such ADHD-like impulsive behavior in adulthood may also due to exposure to sevoflurane during an early age (Fig. 1c, d). Some studies used 5-choice serial reaction time (5-CSRTT) to measure the impulsivity in rats, which required a long time training and learning³³. Still, it is well established that neonatal exposure to sevoflurane is likely to impair animals' memory or learning ability. We think the CAR test, which obtained impulsive-related measures from a free-exploration procedure without learning tasks, was more suitable for this type of study.

A previous study demonstrated that anxiety or fear itself can also result in impulsive-like behavior²⁰. Therefore, to exclude fear- or anxiety-like behavior, we performed the elevated plus maze (EPM) test and found no anxiety alternation in both sevoflurane-treated group and viruses-injected mice. Intriguingly, the sevoflurane group showed less freezing time during the EPM test (Fig. 1g), which reflected sevoflurane-treated mice exhibited less fear. The reason may be that early exposure to sevoflurane has at least two behavioral consequences: increased impulsivity and fearlessness. Importantly, impulsivity and fear are clearly controlled by two separate circuits, as silencing the mPFC CaMKII α ⁺ neurons hardly reverses shortened freezing time shown in mice exposed to sevoflurane. This might also explain why the freezing time had no

significant alternation (Figs. 3l and 4l) when the mPFC neuronal activities were manipulated.

The mPFC encodes and regulates the highest cognitive functions including impulse controlling^{34,35}. The mPFC development is a continuous process and retains its neuroplasticity throughout the life²⁴. Thus, sevoflurane could induce neurodevelopmental diseases by regulating synaptogenesis and synaptic development in the mPFC³⁶, which suggests that mPFC neurons are vulnerable to sevoflurane-induced neurotoxicity. Indeed, our study found that neonatal exposure to sevoflurane-induced ADHD-like impulsive behavior at the adult stage through over-activating the mPFC excitatory neurons (Fig. 2b–d). Another clinical study suggested that glutamate level and the excitatory transmission have a modulatory role in ADHD adults³⁷. We think that the dysfunctional glutamate system in the mPFC may be a pivotal contributor to adult ADHD development.

Based on our observation, the number of c-Fos⁺/CaMKII α ⁺ neurons in the mPFC is significantly increased in mice subjected to neonatal sevoflurane exposure. We believe that the effect is mainly due to the activation of the excitatory neurons. In general, anesthetics act on γ -aminobutyric acid (GABA) receptors to inhibit the neuronal activity and/or attenuating excitatory activity through the glutamate receptors. Prolonged inhibition of neuronal activity has been shown to induce homeostatic upregulation in neuronal excitability³⁸. This intrinsic homeostatic plasticity triggered by relatively long-term sevoflurane exposure may be manifested during a critical developmental period. It will be interesting to examine whether the intrinsic biophysical properties of pyramidal neurons in the miniature excitatory postsynaptic current (mEPSC) in mice with early sevoflurane exposure are enhanced in future studies. Indeed, sevoflurane has been reported to enhance neuronal excitability through elevating corticosteroid levels³⁹ and influencing the T-type Ca²⁺ channels⁴⁰.




Fig. 4 (See legend on next page.)

(see figure on previous page)

Fig. 4 Chemogenetic inhibition of neurons in the mPFC attenuates the sevoflurane-induced impulsive performance. **a** Schematic diagram showing the target of viral injection (top). Location of the bilaterally viral hM4D (Gi) expression (red) in mPFC under control of the CaMKII α promoter (bottom). **b** Current–voltage relationship of a representative mPFC neuron recorded before and during 5 μ M CNO perfusion. Raw traces show individual voltage responses to a series of 600-ms current pulses from 0 to 360 pA with 20 pA steps. **c** The minimal injected current to induce action potential (AP) is increased by CNO. ($n = 10$; Paired Student's t test). **d** The number of induced action potentials at different current steps. ($n = 10$; two-way ANOVA, Bonferroni multiple comparison post hoc tests). **e** Representative movement traces showing the locations of mcherry group (top) and hM4Di group (bottom) sevoflurane-exposure mice in a CAR test following CNO (1 mg/kg) administration. **f, g** Quantification of the number of cumulative jumping events (left) and the latency (right) of the mcherry group and hM4Di group. ($n = 6, 8$ for mcherry, hM4Di group respectively; Mann–Whitney U test). **h, i** Quantification of the number of cumulative jumping events (left) and the latency (right) of the baseline group and hM4Di group. ($n = 8$ for each group; Wilcoxon matched-pairs signed-rank test). **j–l** Quantification of the EPM results of the time spent in the open arms, open arms entries, freezing time of hM4Di and mcherry group after chemogenetic inhibition of mPFC glutamatergic neurons. ($n = 6, 8$ for mcherry, hM4Di group, respectively; Unpaired Student's t test). Data = mean \pm SEM; n.s. no significance, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Both excitatory and inhibitory synaptic inputs have been shown to be regulated by sevoflurane during postnatal development. Lengthy sevoflurane exposure at P7 has been reported to compromise astrocyte development, which arrests excitatory synapse maturation and results in decreased mEPSC frequency in cortical pyramidal neurons⁴¹. Long-term reduction in excitatory inputs may cause homeostatic scaling of excitatory synapses and indirectly enhance neuronal excitability. However, sevoflurane exposure at P16–17 only transiently increased mEPSC frequency and decreased miniature inhibitory postsynaptic current (mIPSC) frequency in mPFC pyramidal neurons in male mice⁴². Thus, it is essential to explore the detailed mechanism underlying early sevoflurane-induced alteration of synaptic integration in postsynaptic cells in future studies. Taken together, we believe that enhanced neuronal excitability in the mPFC in sevoflurane-treated animals may be due to a combination of direct changes in intrinsic properties and altered synaptic integrations of cortical neurons. Besides, the mPFC is composed primarily of glutamatergic neurons (80–90% of neurons) and inhibitory interneurons (10–20% of neurons)⁴³. In our current study, about 15.3% of c-Fos positive cells in the mPFC were not excitatory neurons in the sevoflurane-treated mice during the impulsivity-related behavioral test (Fig. 2d), which indicates that few inhibitory neurons might be activated to balance the overactivation of excitatory neurons in the mPFC.

There are several limitations in this study. The mPFC has complex afferent and efferent projections to mediate the symptoms of psychiatric disorders⁴⁴. Thus, how the downstream regions (e.g., thalamus, striatum, amygdala) of mPFC and their neural circuits are involved in processing behavioral changes in the sevoflurane-treated mice are unknown and warrant further study. Anesthesia and surgery are always going together. Our animals were surgical free and hence how surgery and associated

trauma contribute to this behavioral change is also unknown and awaits further study.

Conclusion

This study demonstrated that repeated sevoflurane exposure during the early development period induces ADHD-like impulsive behavior in later adulthood. This disorder is related to the overactivation of mPFC neurons and can be attenuated by the inhibition of mPFC excitatory neurons. These results may suggest that sevoflurane-induced impulsive behavior is another long-term outcome caused by the anesthetic neurotoxicity and the excitatory neuronal activation in the mPFC may serve as a pathological mechanism for anesthetics-induced ADHD in adults.

Acknowledgements

Thanks for the technical support by the Core Facilities, Zhejiang University School of Medicine. The authors thank Prof. Yi Shen and Zhenxin Yuan, Zhejiang University, for their help and advice during the study. Thanks for foundation support by grants-in-aid for scientific research from the National Natural Science Foundation of China (81471240; 81641042; 81901937) and the Science and Technology Agency, Zhejiang, China (2014C33170).

Author details

¹Department of Anesthesiology, The First Affiliated Hospital, The Children's Hospital, Zhejiang University School of Medicine, Hangzhou, China. ²Department of Anesthesiology, The Children's Hospital, Zhejiang University School of Medicine, Hangzhou, China. ³Department of Cell Biology, Yale University, New Haven, CT, USA. ⁴Department of Neurobiology, The Innovation Center for Brain Science, Institute of Neuroscience, Zhejiang University School of Medicine, Hangzhou, China

Conflict of interest

The authors declare that they have no conflict of interest.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41398-020-00884-5>).

Received: 1 January 2020 Revised: 7 June 2020 Accepted: 8 June 2020
Published online: 20 June 2020

References

- Faraone, S. V. et al. Attention-deficit/hyperactivity disorder. *Nat. Rev. Dis. Prim.* **1**, 15020 (2015).
- Hayward, A., Tomlinson, A. & Neill, J. C. Low attentive and high impulsive rats: a translational animal model of ADHD and disorders of attention and impulse control. *Pharmacol. Therapeutics* **158**, 41–51 (2016).
- Sebastian, A. et al. Frontal dysfunctions of impulse control - a systematic review in borderline personality disorder and attention-deficit/hyperactivity disorder. *Front. Hum. Neurosci.* **8**, 698–698 (2014).
- Schubert, D., Martens, G. J. M. & Kolk, S. M. Molecular underpinnings of prefrontal cortex development in rodents provide insights into the etiology of neurodevelopmental disorders. *Mol. Psychiatry* **20**, 795–809 (2015).
- Hinshaw, S. P. Attention deficit hyperactivity disorder (ADHD): controversy, developmental mechanisms, and multiple levels of analysis. *Annu. Rev. Clin. Psychol.* **14**, 291–316 (2018).
- Xu, L. et al. The association between attention deficit hyperactivity disorder and general anaesthesia - a narrative review. *Anaesthesia* **74**, 57–63 (2019).
- Cubillo, A., Halari, R., Smith, A., Taylor, E. & Rubia, K. A review of fronto-striatal fronto-cortical brain abnormalities in children and adults with Attention Deficit Hyperactivity Disorder (ADHD) and new evidence for dysfunction in adults with ADHD during motivation and attention. *Cortex* **48**, 194–215 (2012).
- Hu, D. et al. Association between exposure of young children to procedures requiring general anesthesia and learning and behavioral outcomes in a population-based birth cohort. *Anesthesiology* **127**, 227–240 (2017).
- Sprung, J. et al. Attention-deficit/hyperactivity disorder after early exposure to procedures requiring general anesthesia. *Mayo Clin. Proc.* **87**, 120–129 (2012).
- Tsai, C.-J. et al. Risk of ADHD after multiple exposures to general anesthesia: a nationwide retrospective cohort study. *J. Atten. Disord.* **22**, 229–239 (2015).
- Fredriksson, A. & Archer, T. Neurobehavioural deficits associated with apoptotic neurodegeneration and vulnerability for ADHD. *Neurotox. Res.* **6**, 435–456 (2004).
- Makaryus, R. et al. Brain maturation in neonatal rodents is impeded by sevoflurane anesthesia. *Anesthesiology* **123**, 557–568 (2015).
- Colon, E. et al. Anesthesia, brain changes, and behavior: insights from neural systems biology. *Prog. Neurobiol.* **153**, 121–160 (2017).
- Alkam, T. et al. Prenatal nicotine exposure decreases the release of dopamine in the medial frontal cortex and induces atomoxetine-responsive neurobehavioral deficits in mice. *Psychopharmacol. (Berl.)* **234**, 1853–1869 (2017).
- Valencia, S. et al. Comparative behavioral correlation of high and low-performing mice in the forced swim test. *Biomol. Ther. (Seoul)* **27**, 349–356 (2019).
- Custodio, R. J. P. et al. Overexpression of the thyroid hormone-responsive (THRSP) gene in the striatum leads to the development of inattentive-like phenotype in mice. *Neuroscience* **390**, 141–150 (2018).
- Yamashita, M. et al. Impaired cliff avoidance reaction in dopamine transporter knockout mice. *Psychopharmacology* **227**, 741–749 (2013).
- Chen, X., Cao, H., Saraf, A., Zweifel, L. S. & Storm, D. R. Overexpression of the type 1 adenylyl cyclase in the forebrain leads to deficits of behavioral inhibition. *J. Neurosci.* **35**, 339–351 (2015).
- Cheng, J. et al. Anterior paraventricular thalamus to nucleus accumbens projection is involved in feeding behavior in a novel environment. *Front. Mol. Neurosci.* **11**, 202 (2018).
- Stein, J. S. et al. Impulsive choice predicts anxiety-like behavior, but not alcohol or sucrose consumption, in male Long-Evans rats. *Alcohol, Clin. Exp. Res.* **39**, 932–940 (2015).
- Joo, J.-Y., Schaukowitz, K., Farbiak, L., Kilaru, G. & Kim, T.-K. Stimulus-specific combinatorial functionality of neuronal c-fos enhancers. *Nat. Neurosci.* **19**, 75–83 (2016).
- Martinon, D. et al. Oxytocin receptors in the dorsolateral bed nucleus of the stria terminalis (BNST) bias fear learning toward temporally predictable cued fear. *Transl. Psychiatry* **9**, 140 (2019).
- Shao, L.-X. et al. Functional coupling of Trm74 and HCN1 channels regulates anxiety-like behavior in BLA neurons. *Mol. Psychiatry* **24**, 1461–1477 (2019).
- Bhattacharjee, A. et al. Cell type-specific transcriptional programs in mouse prefrontal cortex during adolescence and addiction. *Nat. Commun.* **10**, 4169 (2019).
- McCann, M. E. et al. Neurodevelopmental outcome at 5 years of age after general anaesthesia or awake-regional anaesthesia in infancy (GAS): an international, multicentre, randomised, controlled equivalence trial. *Lancet* **393**, 664–677 (2019).
- Flick, R. P. et al. Cognitive and behavioral outcomes after early exposure to anesthesia and surgery. *Pediatrics* **128**, e1053–e1061 (2011).
- Lin, E. P., Soriano, S. G. & Loepke, A. W. Anesthetic neurotoxicity. *Anesthesiol. Clin.* **32**, 133–155 (2014).
- Benveniste, H. & Makaryus, R. Are we moving closer to noninvasive imaging and monitoring of neonatal anesthesia-induced neurotoxicity? *Anesthesiology* **125**, 22–24 (2016).
- Stratmann, G. et al. Effect of general anesthesia in infancy on long-term recognition memory in humans and rats. *Neuropsychopharmacology* **39**, 2275–2287 (2014).
- Ing, C. & Brambrink, A. M. Mayo anesthesia safety in kids continued: two new studies and a potential redirection of the field. *Br. J. Anaesth.* **122**, 716–719 (2019).
- Fitzsimons, M. G., Brookman, J. C., Arnholz, S. H. & Baker, K. Attention-deficit/hyperactivity disorder and successful completion of anesthesia residency: a case report. *Academic Med. J. Assoc. Am. Med. Coll.* **91**, 210–214 (2016).
- Castellheim, A. et al. The role of general anesthesia on traits of neurodevelopmental disorders in a Swedish cohort of twins. *J. Child Psychol. Psychiatry* **59**, 966–972 (2018).
- Liu, J. et al. The claustrum-prefrontal cortex pathway regulates impulsive-like behavior. *J. Neurosci.* **39**, 10071–10080 (2019).
- Euston, D. R., Gruber, A. J. & McNaughton, B. L. The role of medial prefrontal cortex in memory and decision making. *Neuron* **76**, 1057–1070 (2012).
- Warden, M. R. et al. A prefrontal cortex-brainstem neuronal projection that controls response to behavioural challenge. *Nature* **492**, 428–432 (2012).
- Song, R. et al. Maternal sevoflurane exposure causes abnormal development of fetal prefrontal cortex and induces cognitive dysfunction in offspring. *Stem Cells Int.* **2017**, 6158468–6158468 (2017).
- Maltezos, S. et al. Glutamate/glutamine and neuronal integrity in adults with ADHD: a proton MRS study. *Transl. Psychiatry* **4**, e373–e373 (2014).
- Turrigiano, G. Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu. Rev. Neurosci.* **34**, 89–103 (2011).
- Xu, C. et al. Anesthesia with sevoflurane in neonatal rats: Developmental neuroendocrine abnormalities and alleviating effects of the corticosteroid and Cl⁻ importer antagonists. *Psychoneuroendocrinology* **60**, 173–181 (2015).
- Shen, F. Y. et al. Role of T-type calcium channels in generating hyperexcitatory behaviors during emergence from sevoflurane anesthesia in neonatal rats. *Neurosci. Bull.* **36**, 519–529 (2020).
- Zhou, B. et al. Astroglial dysfunctions drive aberrant synaptogenesis and social behavioral deficits in mice with neonatal exposure to lengthy general anesthesia. *PLoS Biol.* **17**, e3000086 (2019).
- Chung, W. et al. Sevoflurane exposure during the critical period affects synaptic transmission and mitochondrial respiration but not long-term behavior in mice. *Anesthesiology* **126**, 288–299 (2017).
- McKvein, J. M., Moloney, R. D., Scheimann, J. R., Myers, B. & Herman, J. P. “Braking” the prefrontal cortex: the role of glucocorticoids and interneurons in stress adaptation and pathology. *Biol. Psychiatry* **86**, 669–681 (2019).
- Riga, D. et al. Optogenetic dissection of medial prefrontal cortex circuitry. *Front. Syst. Neurosci.* **8**, 230 (2014).