



Published in final edited form as:

J Am Chem Soc. 2020 April 22; 142(16): 7362–7367. doi:10.1021/jacs.0c02637.

NHC-Copper Mediated Ligand-Directed Radiofluorination of Aryl Halides

Liam S. Sharninghausen[†], Allen F. Brooks[‡], Wade Winton[‡], Katarina J. Makaravage[†], Peter J. H. Scott^{*‡}, Melanie S. Sanford^{*†}

[†]Department of Chemistry, University of Michigan, 930 North University Avenue, Ann Arbor, Michigan 48109, United States

[‡]Department of Radiology, University of Michigan, 1301 Catherine, Ann Arbor, Michigan 48109, United States

Abstract

[¹⁸F]-labeled aryl fluorides are widely used as radiotracers for positron emission tomography (PET) imaging. Aryl halides (ArX) are particularly attractive precursors to these radiotracers, as they are readily available, inexpensive, and stable. However, to date, the direct preparation of [¹⁸F]-aryl fluorides from aryl halides remains limited to S_NAr reactions between highly activated ArX substrates and K¹⁸F. This report describes an aryl halide radiofluorination reaction in which the C(sp²)-¹⁸F bond is formed via a copper-mediated pathway. Copper N-heterocyclic carbene complexes serve as mediators for this transformation, using aryl halide substrates with directing groups at the ortho position. This reaction is applied to the radiofluorination of electronically diverse aryl halide derivatives, including the bioactive molecules vismodegib and PH-089.

Graphical Abstract



Late-stage methods for constructing ¹⁸F-(hetero)aryl bonds are highly valued for the synthesis of positron emission tomography (PET) radiotracers.^{1,2} Historically, ¹⁸F-labeled aromatic substrates have most commonly been prepared via S_NAr reactions between electron deficient aryl halide precursors and K¹⁸F (Scheme 1A).^{3,4} Aryl halides are particularly attractive radiofluorination precursors because they are abundant, stable, and synthetically accessible. However, the substrate scope of S_NAr (radio)fluorination reactions remains narrow, as resonance electron withdrawing substituents on the aromatic ring are required to stabilize Meisenheimer-type intermediates.^{1,5} Furthermore, even with such highly activated substrates, S_NAr pathways often require long reaction times and forcing conditions, which renders them ill-suited for many late-stage radiofluorination applications.

*Corresponding Author: Peter J. H. Scott. pjhscott@umich.edu. Melanie S. Sanford. mssanfor@umich.edu.


Supporting Information. A listing of the contents of each file supplied as Supporting Information should be included.

^{6,7} As such, a key objective for the field is to develop complementary methods for the radiofluorination of (hetero)aryl-halides and pseudohalides.^{8,9}

Our approach to this challenge has focused on developing Cu-mediated methods for C(*sp*²)-¹⁸F coupling reactions.^{1,2b} Recent studies have shown that Cu salts such as Cu(OTf)₂ and Cu(CH₃CN)₄PF₆ mediate the nucleophilic radiofluorination of aryl stannane,¹⁰ aryl boron,¹¹ diaryliodonium,¹² and aryl C-H substrates¹³ with K¹⁸F. In these systems the key C(*sp*²)-¹⁸F bond is formed via reductive elimination from an organometallic Cu(aryl)(¹⁸F-fluoride) intermediate.^{11f,14} This organometallic pathway is mechanistically distinct from an S_NAr reaction. As such, it enables the radiofluorination of a wide scope of electronically diverse aryl groups.

Despite this progress, analogous Cu mediators have proven ineffective at engaging aryl halide substrates in radiofluorination reactions. Two reports have documented the Cu-promoted nucleophilic ¹⁹F-fluorination of aryl halides (e.g., the work of Liu in Scheme 1B). However, both require superstoichiometric AgF as the fluoride source,^{15,16} and neither has proven translatable to radiolabeling with ¹⁸F⁻ (*vide infra*). This report describes the use of *N*-heterocyclic carbene (NHC) Cu complexes as mediators for ligand-directed aryl halide radiofluorination (Scheme 1C). The discovery of this transformation in the context of ¹⁹F-fluorination and its subsequent translation to radiofluorination are described in detail.

Our initial studies attempted to translate Liu's ¹⁹F-fluorination of 2-(2-bromophenyl)pyridine (Scheme 1B) to a radiolabeling protocol. However, as shown in eq. 1, under the standard conditions (with Cu^I(CH₃CN)₄PF₆, Ag¹⁸F, and NBu₄PF₆ in CH₃CN at 120 °C), no trace of product **1-¹⁸F** was detected by radio-TLC or radio-HPLC after 0.5 h. Furthermore, no improvement was observed upon variation of the ¹⁸F source, solvent, additives, or temperature (Table S7). We note that, in contrast to the ¹⁹F-fluorination, the radiofluorination reaction requires the use of Ag¹⁸F as the limiting reagent at sub-micromolar concentrations. We hypothesize that this renders Cu^I(CH₃CN)₄PF₆-mediated radiofluorination prohibitively slow relative to the decay of the radionuclide (t_{1/2} ~110 min).



Literature reports suggest that aryl-bromide bond activation (via oxidative addition at Cu^I) is likely the slow step in this transformation.^{16,17} We reasoned that the introduction of a strongly electron donating NHC ligand at the Cu^I center would accelerate this key step.^{18,19} Furthermore, since (NHC)Cu^I(F) complexes can be generated directly from KF,²⁰ this approach should eliminate the requirement for excess AgF. Finally, sterically bulky NHC ligands are known to stabilize Cu^I-fluoride complexes to dimerization or disproportionation,^{19,21} which are likely competing decomposition pathways for the Cu mediator.²²

To test this hypothesis, we initially examined the reactivity of a series of (NHC)Cu^I(¹⁹F) complexes with 2-(2-bromophenyl)pyridine (Scheme 2A). As summarized in Table S3, the yield of fluorinated product **1-¹⁹F** varied from 3–65% as a function of the structure of the NHC ligand,^{19,23} with 1,3-bis-(2,6-diisopropylphenyl)imidazol-2-ylidene (IPr) affording the

optimal result. Notably, (IPr)Cu^I(¹⁹F) (**A-¹⁹F**) is available in nearly quantitative yield from the reaction of (IPr)Cu^I(OTf) (**A-OTf**) with K¹⁹F (Scheme 2A),²⁰ thus precluding the requirement for Ag salts in this transformation. Importantly, control studies revealed that other group 11 metal salts including Cu^I(CH₃CN)₄PF₆/KF, CuF₂,²⁴ or AgF afforded 3% of **1-¹⁹F** under otherwise identical conditions (Table S4). Furthermore, no reaction was observed between the aryl bromide substrate and K¹⁹F under these conditions in the absence of copper.

A time study with **A-¹⁹F** shows that the fluorination reaction is complete within 2 h at 140 °C and affords 40% yield after just 30 min (Scheme 2B). This suggests the feasibility of achieving radiofluorination with this system. Finally, a preliminary survey of substrates revealed that **A-¹⁹F**-mediated fluorination has a significantly enhanced scope versus that of Liu's Cu^I(CH₃CN)₄PF₆/Ag¹⁹F system (Scheme 1B). For instance, the sterically hindered pyridine substrate 2-(2-(bromo)phenyl)-6-methylpyridine was unreactive under Liu's conditions, but affords **2-¹⁹F** in 34% yield with **A-¹⁹F** as the Cu mediator (Scheme 2C). Similarly, the oxazoline and imine substrates were unreactive under Liu's conditions, but afford 30% and 37% yield of **3-¹⁹F** and **4-¹⁹F**, respectively, using **A-¹⁹F**.²⁵

We next focused on translating these preliminary results to radiofluorination. The reaction of (IPr)Cu^I(OTf) (**A-OTf**) with 2-(2-bromophenyl)pyridine and K¹⁸F for 30 min at 140 °C in DMF afforded **1-¹⁸F** in 10% radiochemical conversion (RCC) as determined by radio-TLC and radio-HPLC (Table 1, entry 1).^{26,27} The reaction was optimized by exploring additives that have been shown to enhance yields in other Cu-mediated C(*sp*²)-¹⁸F coupling reactions (e.g., phase transfer reagents, nitrogen heterocycles, Table 1, entries 2–5).^{1,11a,13,28} Of the surveyed additives, 1 equiv of 4-dimethylaminopyridine (DMAP) relative to the aryl bromide precursor proved optimal, affording **1-¹⁸F** in 65% RCC.

With these optimized conditions in hand, we next explored the scope of the **A-OTf**-mediated radiofluorination of aryl halides. As shown in Figure 1, the chloro-, bromo-, and iodo-2-phenylpyridine precursors all reacted to afford **1-¹⁸F** in RCCs ranging from 10–65%. In contrast, no ¹⁹F/¹⁸F exchange was detected with **1-¹⁹F** under these conditions. It is currently unclear why **1-I** affords lower yield than **1-Br**; however, this observation is in line with Liu's results for the Cu-catalyzed [¹⁹F]-fluorination of halophenylpyridines.¹⁶ Substitution on either the pyridine or aryl ring was tolerated to afford products such as **2-¹⁸F**, **6-¹⁸F**, and **7-¹⁸F**. Other nitrogen-donors, including oxazoline, pyrazole, cyclohexyl imine, and mesityl imine, served as effective directing groups, affording **3-¹⁸F**, **8-¹⁸F**, **4-¹⁸F**, and **9-¹⁸F**, respectively. The scope of cyclohexyl imine derivatives was most thoroughly explored, as this directing group is straightforward to install and remove starting from readily available benzaldehyde derivatives. Various substitution patterns on the (hetero)arene ring were well tolerated, affording compounds **10–16-¹⁸F** in RCCs ranging from 16–74%. An intramolecular competition reaction between an *ortho*-chloride and bromide resulted in selective radiofluorination of the bromide to form **13-¹⁸F**. This selectivity is consistent with that expected for a metal-mediated activation of a C(*sp*²)-X bond.²⁹

Importantly, a variety of control reactions were conducted in these systems. First, the 4-substituted aryl bromides in the pyridine and cyclohexyl imine series were subjected to the

reaction conditions. These are electronically similar, but do not benefit from the directing effect. As shown in Figure 1, these substrates did not afford detectable **17-¹⁸F** or **18-¹⁸F** under the optimized conditions.³⁰ In addition, all of these reactions were conducted in the absence of Cu to test for background S_NAr reactivity. As shown in Table S12, 1% of compounds **1–16-¹⁸F** were detected under these conditions. Finally, substituting simple Cu^I or Cu^{II} salts for (IPr)Cu^I(OTf) afforded yields of ~5% for representative substrates (Table S11), underscoring the central role of the NHC ligand in these transformations.

¹⁸F-analogues of several bioactive molecules could also be accessed using this approach. In a first example, the bromide analogue of vismodegib (**19-Br**), a basal cell carcinoma treatment,³¹ underwent radiofluorination to afford **19-¹⁸F** (Scheme 3). In a second example, ¹⁸F-labeled PH-089 (**20-¹⁸F** in Scheme 3), an MK-2 inhibitor,³² was synthesized in 5% RCC from the chloride precursor.

A final set of studies focused on automating the radiosynthesis of **1-¹⁸F** using a TRACERLab FX_{FN} synthesis module. Initial automated studies using 241.1 mCi (8.93×10^9 Bq) of K¹⁸F gave 57 ± 8 % radiochemical yield (RCY; n = 2), demonstrating the compatibility of the method with automation. Further investigations coupled automated synthesis with semi-preparative HPLC purification to afford **1-¹⁸F** in 14.3 ± 3.2 % RCY (decay-corrected; 119.9 mCi \pm 28; n = 2) with good molar activity (1614 ± 353 Ci/mmol; n = 2) and radiochemical purity. While unoptimized, this result demonstrates the potential of this method for PET applications.

In conclusion, we have developed a Cu-mediated protocol for the ¹⁹F- and ¹⁸F-fluorination of diverse aryl halide substrates. Strategic design of the Cu mediator was necessary to achieve the reaction rates/yields required for efficient radiofluorination, and an NHC-ligated Cu complex ultimately proved optimal in this system. A wide scope of nitrogen-containing directing groups and substituted aryl halide derivatives underwent ¹⁸F-fluorination, and the reaction proved effective for the synthesis of biologically relevant molecules such as **19-¹⁸F** and **20-¹⁸F**. More broadly, this work demonstrates that NHC-type ligands enable new C(*sp*²)-F coupling reactions at Cu. As such, this work opens up opportunities for designing next-generation Cu mediators for the radiofluorination of currently inert substrates (e.g., aryl halides that lack a directing group).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

This work was supported by the NIH [award number F32GM136022 (LSS) and award number R01EB021155 (MSS and PJHS)]. We acknowledge Dr. Devin M. Ferguson, Isaac M. Blythe, Dr. Yiyang See, Dr. Sean M. Preshlock, Dr. Jay S. Wright and Pronay Roy for helpful discussions. We also acknowledge Dr. Rachel S. Plumb for assistance with data processing.

REFERENCES

1. Preshlock S; Tredwell M; Gouverneur V 18F-Labeling of Arenes and Heteroarenes for Applications in Positron Emission Tomography. Chem. Rev 2016, 116, 719–766. [PubMed: 26751274]


2. (a)Miller PW; Long NJ; Vilar R; Gee AD Synthesis of ¹¹C, ¹⁸F, ¹⁵O, and ¹³N Radiolabels for Positron Emission Tomography. *Angew. Chem. Int. Ed* 2008, 47, 8998–9033.(b)Brooks AF; Topczewski JJ; Ichiishi N; Sanford MS; Scott PJH Late-Stage [¹⁸F]Fluorination: New Solutions to Old Problems. *Chem. Sci* 2014, 5, 4545–4553. [PubMed: 25379166] (c)Coenen HH; Ermert J ¹⁸F-labelling Innovations and Their Potential for Clinical Application. *Clin. Transl. Imaging* 2018, 6, 169–193.(d)Chen W; Huang Z; Tay NES; Giglio B; Wang M; Wang H; Wu Z; Nicewicz DA; Li Z Direct Arene C-H Fluorination with ¹⁸F- via Organic Photoredox Catalysis. *Science* 2019, 364, 1170–1174. [PubMed: 31221856] (e)Xu P; Zhao D; Berger F; Hamad A; Rickmeier J; Petzold R; Kondratiuk M; Bohdan KJ; Ritter T Site-Selective Late-Stage Aromatic [¹⁸F]Fluorination via Aryl Sulfonium Salts. *Angew. Chem. Int. Ed* 2020, 59, 1956–1960.
3. For examples of the SNAr radiofluorination of aryl halides, see:(a)Lemaire C; Guillaume M; Christiaens L; Palmer AJ; Cantineau R A New Route for the Synthesis of [¹⁸F]Fluoroaromatic Substituted Amino Acids: No Carrier Added L-p-[¹⁸F]-Fluorophenylalanine. *Appl. Radiat. Isot* 1987, 38, 1033–1038.(b)Pascali C; Luthra SK; Pike VW; Price GW; Ahier RG; Hume SP; Mers R; Manjil L; Cremer JE The Radiosynthesis of [¹⁸F]PK 14105 as an Alternative Radioligand for Peripheral Type Benzodiazepine Binding Sites. *Appl. Radiat. Isot* 1990, 41, 477–482.(c)Hashizume K; Tamakawa H; Hashimoto N; Miyake Y Single-step Synthesis of [¹⁸F]Haloperidol from the Chloro-precursor and its Application in PET Imaging of a Cat's Brain. *Appl. Radiat. Isot* 1997, 48, 1179–1185. [PubMed: 9418209] (d)Shen B; Loffler D; Zeller K-P; Ubele M; Reischl G; Machulla H-J Decarbonylation of Multi-substituted [¹⁸F]- Benzaldehydes for Modelling Synthesis of ¹⁸F- Labelled Aromatic Amino Acids. *Appl. Radiat. Isot* 2007, 65, 1227–1231. [PubMed: 17681802]
4. For examples of the SNAr radiofluorination of halopyridines, see:(a)Beer H-F; Haeberli M; Ametamey S; Schubiger PA Comparison of Two Synthetic Methods to Obtain [¹⁸F] N-(2-aminoethyl)-5-fluoropyridine-2-carboxamide, a Potential Mao-B Imaging Tracer for PET. *J. Labelled Compd. Radiopharm* 1995, 36, 933–945.(b)Karramkam M; Hinnen F; Vaufrey F; Dollé, F. 2-, 3- and 4- [¹⁸F]Fluoropyridine by No-Carrier-Added Nucleophilic Aromatic Substitution with K[¹⁸F]F-K222 – A Comparative Study. *J. Labelled Compd. Radiopharm* 2003, 46, 979–992. (c)Malik N; Voelter W; Machulla H-J; Solbach C Radiofluorination of 2-Fluoropyridines by Isotopic Exchange with [¹⁸F]Fluoride. *J. Radioanal. Nucl. Chem* 2011, 287, 287–292.
5. (a)Terrier F Modern Nucleophilic Aromatic Substitution; Wiley-VCH, Weinheim, 2013, p. 236–268. (b)Cole EL; Stewart MN; Littich R; Hoareau R; Scott PJH Radiosyntheses using Fluorine-18: the Art and Science of Late Stage Fluorination. *Curr. Top. Med. Chem* 2014, 14, 875–900. [PubMed: 24484425] (c)Jacobson O; Kiesewetter DO; Chen X Fluorine-18 Radiochemistry, Labeling Strategies and Synthetic Routes. *Bioconjugate Chem* 2015, 26, 1–18.
6. In many cases, radiofluorination of simple building blocks followed by subsequent synthetic steps is required. See:(a)Krüll J; Heinrich MR [¹⁸F]Fluorine-Labeled Pharmaceuticals: Direct Aromatic Fluorination Compared to Multi-Step Strategies. *Asian J. Org. Chem* 2019, 8, 576–590.(b)van der Born D; Pees A; Poot AJ; Orru RVA; Windhorst AD; Vugts DJ Fluorine-18 Labelled Building Blocks for PET Tracer Synthesis. *Chem. Soc. Rev* 2017, 46, 4709–4773. [PubMed: 28608906]
7. (a)Jacobson O; Chen X PET Designated Fluoride-18 Production and Chemistry. *Curr. Top. Med. Chem* 2010, 10, 1048–1059. [PubMed: 20388116] (b)Sun H; DiMugno S G Competitive Demethylation and Substitution in N,N,N-Trimethylanilinium Fluorides. *J. Fluor. Chem* 2007, 128, 806–812.(c)Ermert J; Hocke C; Ludwig T; Gail R; Coenen HH Comparison of Pathways to the Versatile Synthone of No-Carrier-Added 1-Bromo-4-[¹⁸F]fluorobenzene. *J. Labelled Compd. Radiopharm* 2004, 47, 429–441.
8. (a)Grushin VV The Organometallic Fluorine Chemistry of Palladium and Rhodium: Studies toward Aromatic Fluorination. *Acc. Chem. Res* 2010, 43, 160–171. [PubMed: 19788304] (b)Campbell MG; Ritter T Modern Carbon–Fluorine Bond Forming Reactions for Aryl Fluoride Synthesis *Chem. Rev* 2015, 115, 612–633. [PubMed: 25474722]
9. For an example of indirect radiofluorination of aryl bromides via isolated NiII-aryl intermediates, see:Lee E; Hooker JM; Ritter T Nickel-Mediated Oxidative Fluorination for PET with Aqueous [¹⁸F] Fluoride. *J. Am. Chem. Soc* 2012, 134, 17456–17458. [PubMed: 23061667]
10. (a)Makaravage KJ; Brooks AF; Mossine AV; Sanford MS; Scott PJH Copper-Mediated Radiofluorination of Arylstannanes with [¹⁸F]KF. *Org. Lett* 2016, 18, 5440–5443. [PubMed: 27718581] (b)Gamache RF; Waldmann C; Murphy JM Copper-Mediated Oxidative Fluorination of Aryl Stannanes with Fluoride. *Org. Lett* 2016, 18, 4522–4525. [PubMed: 27571319] (c)Zarrad

F; Zlatopolskiy BD; Krapf P; Zischler J; Neumaier B A Practical Method for the Preparation of ¹⁸F-Labeled Aromatic Amino Acids from Nucleophilic [¹⁸F]Fluoride and Stannyl Precursors for Electrophilic Radiohalogenation. *Molecules*. 2017, 22, 2231.

11. (a)Mossine AV; Brooks AF; Makaravage KJ; Miller JM; Ichiishi N; Sanford MS; Scott PJH Synthesis of [¹⁸F]Arenes via the Copper-Mediated [¹⁸F]Fluorination of Boronic Acids. *Org. Lett* 2015, 17, 5780–5783. [PubMed: 26568457] (b)Tredwell M; Preshlock SM; Taylor NJ; Gruber S; Huiban M; Passchier J; Mercier J; Génicot C; Gouverneur V A General Copper-Mediated Nucleophilic ¹⁸F Fluorination of Arenes *Angew. Chem. Int. Ed* 2014, 53, 7751–7755. (c)Zlatopolskiy BD Zischler J; Krapf P; Zarrad F; Urusova EA; Kordys E; Endepols H; Neumaier B Copper-mediated Aromatic Radiofluorination Revisited: Efficient Production of PET Tracers on a Preparative Scale. *Chem. Eur. J* 2015, 21, 5972–5979. [PubMed: 25708748] (d)Preshlock S; Calderwood S; Verhoog S; Tredwell M; Huiban M; Hienzsch A; Gruber S; Wilson TC; Taylor NJ; Cailly T; Schedler M; Collier TL; Passchier J; Smits R; Mollitor J; Hoepping A; Mueller M; Genicot C; Mercier J; Gouverneur V Enhanced Copper-Mediated ¹⁸F-Fluorination of Aryl Boronic Esters provides Eight Radiotracers for PET Applications. *Chem. Commun* 2016, 52, 8361–8364. (e)Mossine AV; Brooks AF; Bernard-Gauthier V; Bailey JJ; Ichiishi N; Schirmmacher R; Sanford MS; Scott PJH Automated Synthesis of PET Radiotracers by Copper-mediated ¹⁸F-Fluorination of Organoborons: Importance of the Order of Addition and Competing Protodeborylation. *J. Label. Compd. Radiopharm* 2018, 61, 228–236. (f)Taylor NJ; Emer E; Preshlock S; Schedler M; Tredwell M; Verhoog S; Mercier J; Genicot C; Gouverneur V Derisking the Cu-Mediated ¹⁸F-Fluorination of Heterocyclic Positron Emission Tomography Radioligands. *J. Am. Chem. Soc* 2017, 139, 8267–8276. [PubMed: 28548849] (g)Zischler J; Kolks N; Modemann D; Neumaier B; Zlatopolskiy BD Alcohol-enhanced Cu-mediated Radiofluorination. *Chem. Eur. J* 2017, 23, 3251–3256. [PubMed: 27943464] (h)Wilson TC; Cailly T; Gouverneur V Boron Reagents for Divergent Radiochemistry. *Chem Soc Rev* 2018, 47, 6990–7005. [PubMed: 30140795]
12. Ichiishi N; Brooks AF; Topczewski JJ; Rodnick ME; Sanford MS; Scott PJH Copper- Catalyzed [¹⁸F]Fluorination of (Mesityl)(aryl)iodonium Salts. *Org. Lett* 2014, 16, 3224–3227. [PubMed: 24890658]
13. Lee SJ; Makaravage KJ; Brooks AF; Sanford MS; Scott PJH Cu-Mediated Aminoquinoline-Directed Radiofluorination of Aromatic C–H Bonds with KF. *Angew. Chem. Int. Ed* 2019, 58, 3119–3122.
14. (a)Casitas A; Canta M; Sola M; Costas M; Ribas X Nucleophilic Aryl Fluorination and Aryl Halide Exchange Mediated by a CuI/CuIII Catalytic Cycle. *J. Am. Chem. Soc* 2011, 133, 19386–19392. [PubMed: 22026511] (b)Ichiishi N; Cauty AJ; Yates BF; Sanford MS Mechanistic Investigations of Cu-Catalyzed Fluorination of Diaryliodonium Salts: Elaborating the CuI/CuIII Manifold for Copper Catalysis. *Organometallics* 2014, 33, 5525–5534. [PubMed: 25328272] (c)Ye Y; Schimler SD; Hanley PS; Sanford MS Cu(OTf)₂-Mediated Fluorination of Aryltrifluoroborates with Potassium Fluoride. *J. Am. Chem. Soc* 2013, 135, 16292–16295. [PubMed: 24160267] (d)Yao B.; Wang Z-L; Zhang H; Wang D-X; Zhao L; Wang M-X Cu(ClO₄)₂-Mediated Arene C–H Bond Halogenations of Azacalixaromatics Using Alkali Metal Halides as Halogen Sources. *J. Org. Chem* 2012, 77, 3336–3340. [PubMed: 22420606]
15. Fier PS; Hartwig JF Copper-Mediated Fluorination of Aryl Iodides. *J. Am. Chem. Soc* 2012, 134, 10795–10798. [PubMed: 22709145]
16. Mu X; Zhang H; Chen P; Liu G Copper-Catalyzed Fluorination of 2-Pyridyl Aryl Bromides. *Chem. Sci* 2014, 5, 275–280.
17. Le C; Chen TQ; Liang T; Zhang P; MacMillan DWC A Radical Approach to the Copper Oxidative Addition Problem: Trifluoromethylation of Bromoarenes. *Science*, 2018, 360, 1010–1014. [PubMed: 29853683]
18. (a)Hopkinson MN; Richter C; Schedler M; Glorius F An Overview of N-Heterocyclic Carbenes. *Nature* 2014, 510, 485–496. [PubMed: 24965649] (b)Danopoulos AA; Simler T; Braunstein P N-Heterocyclic Carbene Complexes of Copper, Nickel, and Cobalt. *Chem. Rev* 2019, 119, 3730–3961. [PubMed: 30843688] (c)Díez-Gonzalez S; Marion N; Nolan S[†] N-Heterocyclic Carbenes in Late Transition Metal Catalysis. *Chem. Rev* 2009, 109, 3612–3676. [PubMed: 19588961] (d)Jurkaskas V; Sadighi JP; Buchwald SL Conjugate Reduction of α,β -Unsaturated Carbonyl

Compounds Catalyzed by a Copper Carbene Complex. *Org. Lett* 2003, 5, 2417–2420. [PubMed: 12841744]


19. The reaction of the (NHC)Cu(19F) complex (sICy)Cu(19F) with p-NO₂PhI at 100 °C was reported to afford p-NO₂Ph19F in 16% yield. However, this result was not followed up in further detail. Laitar, D. S. *Synthetic and Catalytic Studies of Group 11 N-Heterocyclic Carbene Complexes*. Ph.D. Thesis, Massachusetts Institute of Technology, Cambridge, MA, 2006.
20. Dang H; Mailig M; Lalic G Mild Copper-Catalyzed Fluorination of Alkyl Triflates with Potassium Fluoride. *Angew. Chem. Int. Ed* 2014, 53, 6473–6476.
21. (a)Herron JR; Ball ZT Synthesis and Reactivity of Functionalized Arylcopper Compounds by Transmetalation of Organosilanes. *J. Am. Chem. Soc* 2008, 130, 16486–16487. [PubMed: 19554685] (b)Fujihara T; Xu T; Semba K; Terao J; Tsuji Y Copper-Catalyzed Hydrocarboxylation of Alkynes Using Carbon Dioxide and Hydrosilanes. *Angew. Chem. Int. Ed* 2011, 50, 523–527.
22. Waddington TC The Lattice Energies and Thermodynamic Properties of the Hypothetical Compounds AuF and CuF. *Trans. Faraday Soc* 1959, 55, 1531–1535.
23. With many of these complexes, a side product resulting from the C–C coupling of the NHC ligand and 2-(2-bromophenyl)pyridine was detected by high resolution mass spectrometry (see SI, section 3.6 for complete details). Notably, both NHC-aryl and NHC-halide coupling products have been observed in other reactions of (NHC)Cu complexes.(a)Williams TJ; Bray JTW; Lake BRM; Willans CE; Rajabi NA; Ariaferd A; Manzini C; Bellina F; Whitwood AC; Fairlamb IJS Mechanistic Elucidation of the Arylation of Non-Spectator N-Heterocyclic Carbenes at Copper Using a Combined Experimental and Computational Approach. *Organometallics* 2015, 34, 3497–3507.(b)Grandbois A; Mayer M-E; Bedard M; Collins SK; Michel T Synthesis of C1-Symmetric BINOLs Employing N-Heterocyclic Carbene–Copper Complexes *Organometallics* 2010, 29, 3683–3685.(c)Cheng J; Wang L; Wang P; Deng L High-Oxidation-State 3d Metal (Ti–Cu) Complexes with N-Heterocyclic Carbene Ligation. *Chem. Rev* 2018, 118, 9930–9987. [PubMed: 30011189]
24. The following patent reported the stoichiometric fluorination of aryl halides with CuF₂ at 180 °C. However, minimal details are provided. Grushin, V. V. (DuPont & Co., USA) Process for Preparing Fluoroarenes from Haloarenes. U.S. Patent 7,202,388, 5 10, 2007.
25. In contrast, aryl halides without directing groups (e.g., 4-bromobiphenyl) afforded no detectable fluorinated products under these conditions (see SI, section 3.3 for complete details).
26. Control reactions (in the absence of Cu or using various other Cu precursors) result in 1% RCC under analogous conditions (Tables S11 and S12).
27. Procedure for radiofluorination involves initial combination of aryl halide, A-OTf, DMAP and DMF in a N₂ glove box. The reaction mixture is removed from the glovebox and a DMF solution of K18F is added to the reaction vial through a septa cap prior to heating. See SI, section 5.3 for full details.
28. Mossine AV; Brooks AF; Ichiishi N; Makaravage KJ; Sanford MS; Scott PJH Development of Customized [18F]Fluoride Elution Techniques for the Enhancement of Copper-Mediated Late-Stage Radiofluorination. *Sci. Rep* 2017, 7, 233. [PubMed: 28331174]
29. Tomashenko OA; Grushin VV Aromatic Trifluoromethylation with Metal Complexes. *Chem. Rev* 2011, 111, 4475–4521. [PubMed: 21456523]
30. Similarly, other aryl halides lacking directing groups were not effective substrates (SI, Table S13)
31. Dlugosz A; Agrawal S; Kirkpatrick P Vismodegib. *Nat. Rev. Drug Discovery* 2012, 11, 437–438. [PubMed: 22653209]
32. The chloride substrate was used (rather than the bromide) based on the availability of a straightforward literature prep for this compound. Anderson DR; Meyers MJ; Vernier WF; Mahoney MW; Kurumbail RG; Caspers N; Poda GI; Schindler JF; Reitz DB; Mourey RJ Pyrrolopyridine Inhibitors of Mitogen-Activated Protein Kinase-Activated Protein Kinase 2 (MK-2). *J. Med. Chem* 2007, 50, 2647–2654. [PubMed: 17480064]

**Figure 1.**


Substrate scope of Cu-mediated radiofluorination of aryl halides.

Conditions: aryl halide (0.005 mmol, 1 equiv), **A-OTf** (1 equiv), DMAP (1 equiv), $K^{18}F$, DMF (0.015 M), N_2 atmosphere, 140 °C for 30 min.²⁷ RCC determined by radio-TLC (n 3). ^aReaction conducted at 160 °C.


A. Radiofluorination of ArX via S_NAr pathway (*ref. 3*)






B. Liu's directed ¹⁹F-fluorination of ArBr via organometallic pathway (*ref. 16*)




C. Directed radiofluorination of ArX via organometallic pathway (*this work*)



Scheme 1.
Strategies for direct fluorination of aryl halides.

A. Synthesis and reactivity of **A-¹⁹F**B. Time study for **A-¹⁹F**-mediated fluorinationC. Unoptimized yields of ¹⁹F fluorinated products with **A-¹⁹F** (after 21 h)**Scheme 2.**NHC-Cu-mediated ¹⁹F-fluorination of aryl bromides


(A) Conditions: **A-OTf** (0.006 mmol, 1 equiv), KF (1.5 equiv), DMF (0.01 M), 140 °C for 30 min, then aryl bromide (0.006 mmol), 140 °C for 21 h. (B) Conditions: **A-¹⁹F** (0.01 mmol, 1 equiv), aryl bromide (1 equiv), DMF (0.015 M), 140 °C for 21 h. (C) Conditions: **A-¹⁹F** (0.006 mmol, 1 equiv), aryl bromide (1 equiv), DMF (0.01 M), 140 °C for 21 h. Yields determined by ¹⁹F NMR spectroscopic analysis of crude reaction mixtures.



Scheme 3.
Radiofluorination of bioactive molecules.

Table 1.

Cu-mediated radiofluorination of aryl halides.



entry	Additive	RCC (%)
1	none	10
2	Kryptofix	26
3	pyridine	23
4	DBU	30
5	DMAP	65

Conditions: aryl bromide (0.005 mmol, 1 equiv), A-OTf(1 equiv), additive (1 equiv), K¹⁸F, DMF (0.015 M), N₂ atmosphere, 140 °C, 30 min.²⁷
 RCC determined by radio-TLC (n = 2).